

SPECIAL PUBLICATION No. 98 CGS Special Publ. 98


U. S. DEPARTMENT OF COMMERCE  
COAST AND GEODETIC SURVEY

---

**MANUAL OF  
HARMONIC ANALYSIS  
AND PREDICTION OF TIDES**

---

SPECIAL PUBLICATION No. 98  
REVISED (1940) EDITION



MLB/MHOI



0 0301 009547 5

**U. S. DEPARTMENT OF COMMERCE**

JESSE H. JONES, Secretary

**COAST AND GEODETIC SURVEY**

LEO OTIS COLBERT, Director

---

**Special Publication No. 98**

Revised (1940) Edition

(Reprinted 1958 with corrections)

---

**MANUAL OF HARMONIC ANALYSIS  
AND PREDICTION OF TIDES**

BY

**PAUL SCHUREMAN**

Senior Mathematician



**Reprinted October 1971**

UNITED STATES  
GOVERNMENT PRINTING OFFICE  
WASHINGTON : 1958

## PREFACE

This volume was designed primarily as a working manual for use in the United States Coast and Geodetic Survey and describes the procedure used in this office for the harmonic analysis and prediction of tides and tidal currents. It is based largely upon the works of Sir William Thomson, Prof. George H. Darwin, and Dr. Rollin A. Harris. In recent years there also has been considerable work done on this subject by Dr. A. T. Doodson, of the Tidal Institute of the University of Liverpool.

The first edition of the present work was published in 1924. In this revised edition there has been a rearrangement of the material in the first part of the volume to bring out more clearly the development of the tidal forces. Tables of astronomical data and other tables to facilitate the computations have been retained with a few revisions and additions and there has been added a list of symbols used in the work.

The collection of tidal harmonic constants for the world that appeared in the earlier edition has been omitted altogether because the work of maintaining such a list has now been taken over by the International Hydrographic Bureau at Monaco. These constants are now published in International Hydrographic Bureau Special Publication No. 26, which consists of a collection of loose sheets which permit the addition of new constants as they become available.

Special acknowledgment is due Walter B. Zerbe, associate mathematician of the Division of Tides and Currents, who reviewed the manuscript of this edition and offered many valuable suggestions.

## CONTENTS

---

	Page
<b>Introduction</b>	1
Historical statement	1
General explanation of tidal movement	2
Harmonic treatment of tidal data	2
Astronomical data	3
Degree of approximation	8
<b>Development of tide-producing force</b>	10
Fundamental formulas	10
Vertical component of force	15
Horizontal components of force	26
Equilibrium tide	28
Terms involving 4th power of moon's parallax	34
Solar tides	39
The $M_1$ tide	41
The $L_2$ tide	43
Lunisolar $K_1$ and $K_2$ tides	44
Meteorological and shallow-water tides	46
<b>Analysis of observations</b>	49
Harmonic constants	49
Observational data	50
Summations for analysis	52
Stencils	53
Secondary stencils	57
Fourier series	62
Augmenting factors	71
Phase lag or epoch	75
Inference of constants	78
Elimination	84
Long period constituents	87
Analysis of high and low waters	100
Forms used for analysis of tides	104
Analysis of tidal currents	118
<b>Prediction of tides</b>	123
Harmonic method	123
Tide-predicting machine	126
Forms used with tide-predicting machine	143
Prediction of tidal currents	147
<b>Tables</b>	153
Explanation of tables	153
1. Fundamental astronomical data	162
2. Harmonic constituents	164
2a. Shallow-water constituents	167
3. Latitude factors	168
4. Mean longitude of lunar and solar elements	170
5. Differences to adapt table 4 to any month, day, and hour	172
6. Values of $I$ , $\nu$ , $\xi$ , $\nu'$ , and $2\nu''$ for each degree of $N$	173
7. Values of $\log R_a$ for amplitude of constituent $L_2$	177
8. Values of $R$ for argument of constituent $L_2$	178
9. Values of $\log Q$ for amplitude of constituent $M_1$	179
10. Values of $Q$ for argument of constituent $M_1$	180
11. Values of $u$ of equilibrium arguments for each degree of $N$	182
12. Values of log factor $F$ for each tenth degree of $I$	186
13. Values of $u$ and log factor $F$ for constituents $L_2$ and $M_1$ for the years 1900 to 2000	192
14. Node factor $f$ for middle of each year 1850 to 1999	199
15. Equilibrium argument $V_o + u$ for beginning of each year 1850 to 2000	204

## Tables—Continued.

16. Differences to adapt table 15 to beginning of each calendar month-	212
17. Differences to adapt table 15 to beginning of each day of month-	213
18. Differences to adapt table 15 to beginning of each hour of day-	216
19. Products for Form 194-	218
20. Augmenting factors-	228
21. Acceleration in epoch of $K_1$ due to $P_1$ -	229
22. Ratio of increase in amplitude of $K_1$ due to $P_1$ -	229
23. Acceleration in epoch of $S_2$ due to $K_2$ -	230
24. Ratio of increase in amplitude of $S_2$ due to $K_2$ -	230
25. Acceleration in epoch of $S_2$ due to $T_2$ -	231
26. Resultant amplitude $S_2$ due to $T_2$ -	232
27. Critical logarithms for Form 245-	233
28. Constituent speed differences ( $b-a$ )-	234
29. Elimination factors-	236
30. Products for Form 245-	266
31. For construction of primary stencils-	268
32. Divisors for primary stencil sums-	288
33. For construction of secondary stencils-	299
34. Assignment of daily page sums for long-period constituents-	302
35. Products for Form 444-	304
36. Angle differences for Form 445-	306
37. Coast and Geodetic Survey tide-predicting machine No. 2—general gears-	307
38. Coast and Geodetic Survey tide-predicting machine No. 2—constituent gears-	308
39. Synodic periods of constituents-	309
40. Day of common year corresponding to day of month-	309
41. Values of $h$ in formula $h = (1 + r^2 + 2r \cos x)^{\frac{1}{2}}$ -	310
42. Values of $k$ in formula $k = \tan^{-1} \frac{r \sin x}{1 + r \cos x}$ -	310
Explanation of symbols-	311
Index-	314

## ILLUSTRATIONS

1. Ecliptic, celestial equator, and moon's orbit-	6
2. Tide-producing force-	11
3. Celestial sphere-	16
4. Longitude relations-	19
5. Equilibrium tide with moon on equator-	29
6. Equilibrium tide with moon at maximum declination-	29
7. Constituent tide curve-	50
8. Phase relations-	77
9. Form 362, hourly heights-	105
10. Stencil for constituent M-	106
11. Application of stencil-	107
12. Form 142, stencil sums-	108
13. Computation of hourly means-	109
14. Form 244, computation of $V_o + u$ -	110
15. Form 244a, log $F$ and arguments for elimination-	111
16. Form 194, harmonic analysis-	112
17. Form 452, $R$ , $\kappa$ , and $\xi$ from analysis and inference, diurnal tides-	115
18. Form 452, $R$ , $\kappa$ , and $\xi$ from analysis and inference, semidiurnal tides-	116
19. Form 245, elimination-	117
20. Form 723, currents, harmonic comparison-	120
21. Coast and Geodetic Survey tide-predicting machine-	128
22. Tide-predicting machine, time side-	128
23. Tide-predicting machine, recording devices-	128
24. Tide-predicting machine, driving gears-	128
25. Tide-predicting machine, dial case from height side-	128
26. Tide-predicting machine, dial case from time side-	128
27. Tide-predicting machine, vertical driving shaft of middle section-	128
28. Tide-predicting machine, forward driving shaft of rear section-	128
29. Tide-predicting machine, rear end-	128
30. Tide-predicting machine, details of releasable gear-	128
31. Tide-predicting machine, details of constituent crank-	128
32. Form 444, standard harmonic constants for predictions-	143
33. Form 445, settings for tide-predicting machine-	145
34. Graphic solution of formulas (470) and (471)-	149

# MANUAL OF HARMONIC ANALYSIS AND PREDICTION OF TIDES

## INTRODUCTION

### HISTORICAL STATEMENT

1. Sir William Thomson (Lord Kelvin) devised the method of reduction of tides by harmonic analysis about the year 1867. The principle upon which the system is based—which is that any periodic motion or oscillation can always be resolved into the sum of a series of simple harmonic motions—is said to have been discovered by Eudoxas as early as 356 B. C., when he explained the apparently irregular motions of the planets by combinations of uniform circular motions.<sup>1</sup> In the early part of the nineteenth century Laplace recognized the existence of partial tides that might be expressed by the cosine of an angle increasing uniformly with the time, and also applied the essential principles of the harmonic analysis to the reduction of high and low waters. Dr. Thomas Young suggested the importance of observing and analyzing the entire tidal curve rather than the high and low waters only. Sir George B. Airy also had an important part in laying the foundation for the harmonic analysis of the tides. To Sir William Thomson, however, we may give the credit for having placed the analysis on a practical basis.

2. In 1867 the British Association for the Advancement of Science appointed a committee for the purpose of promoting the extension, improvement, and harmonic analysis of tidal observations. The report on the subject was prepared by Sir William Thomson and was published in the Report of the British Association for the Advancement of Science in 1868. Supplementary reports were made from time to time by the tidal committee and published in subsequent reports of the British association. A few years later a committee, consisting of Profs. G. H. Darwin and J. C. Adams, drew up a very full report on the subject, which was published in the Report of the British Association for the Advancement of Science in 1883.

3. Among the American mathematicians who have had an important part in the development of this subject may be named Prof. William Ferrel and Dr. Rollin A. Harris, both of whom were associated with the U. S. Coast and Geodetic Survey. The Tidal Researches, by Professor Ferrel, was published in 1874, and additional articles on the harmonic analysis by the same author appeared from time to time in the annual reports of the Superintendent of the Coast and Geodetic Survey. The best known work of Doctor Harris is his Manual of Tides, which was published in several parts as appendices to the annual reports of the Superintendent of the Coast and Geodetic Survey. The subject of the harmonic analysis was treated principally in Part II of the Manual which appeared in 1897.

<sup>1</sup> Nautical Science, p. 279, by Charles Lane Poor.

## GENERAL EXPLANATION OF TIDAL MOVEMENT

4. That the tidal movement results from the gravitational attraction of the moon and sun acting upon the rotating earth is now a well-established scientific fact. The movement includes both the vertical rise and fall of the tide and the horizontal flow of the tidal currents. It will be shown later that the tide-producing force due to this attraction, when taken in connection with the attraction between the particles of matter which constitute the earth, can be expressed by mathematical formulas based upon the well-known laws of gravitation.

5. Although the acting forces are well understood, the resultant tidal movement is exceedingly complicated because of the irregular distribution of land and water on the earth and the retarding effects of friction and inertia. Contrary to the popular idea of a progressive tidal wave following the moon around the earth, the basic tidal movement as evidenced by observations at numerous points along the shores of the oceans consists of a number of oscillating areas, the movement being somewhat similar to that in a pan of water that has been tilted. Such oscillations are technically known as stationary waves. The complex nature of the movement can be appreciated when consideration is given to the fact that such stationary waves may overlap or be superimposed upon each other and may be accompanied by a progressive wave movement.

6. Any basin of water has its natural free period of oscillation depending upon its size and depth. The usual formula for the period of oscillation in a rectangular tank of uniform depth is  $2L/\sqrt{gd}$ , in which  $L$  is the length and  $d$  the depth of the tank and  $g$  is the acceleration of gravity. When a disturbing force is applied periodically at intervals corresponding to the free period of a body of water, it tends to build up an oscillation of much greater magnitude than would be possible with a single application of the force. The major tidal oscillations have periods approximating the half and the whole lunar day.

## HARMONIC TREATMENT OF TIDAL DATA

7. The harmonic analysis of tides is based upon an assumption that the rise and fall of the tide in any locality can be expressed mathematically by the sum of a series of harmonic terms having certain relations to astronomical conditions. A simple harmonic function is a quantity that varies as the cosine of an angle that increases uniformly with time. In the equation  $y = A \cos at$ ,  $y$  is an harmonic function of the angle  $at$  in which  $a$  is a constant and  $t$  represents time as measured from some initial epoch. The general equation for the height ( $h$ ) of the tide at any time ( $t$ ) may be written

$$h = H_0 + A \cos (at + \alpha) + B \cos (bt + \beta) + C \cos (ct + \gamma) + \text{etc.} \quad (1)$$

in which  $H_0$  is the height of the mean water level above the datum used. Other symbols are explained in the following paragraph.

8. Each cosine term in equation (1) is known as a *constituent* or *component* tide. The coefficients  $A$ ,  $B$ ,  $C$ , etc. are the *amplitudes* of the constituents and are derived from observed tidal data in each locality. The expression in parentheses is a uniformly-varying angle and its value at any time is called its *phase*. Any constituent term has its maximum positive value when the phase of the angle is zero and a maximum negative value when the phase equals  $180^\circ$ , and the

term becomes zero when the phase equals  $90^\circ$  or  $270^\circ$ . The coefficient of  $t$  represents the rate of change in the phase and is called the *speed* of the constituent and is usually expressed in degrees per hour. The time required for a constituent to pass through a complete cycle is known as its *period* and may be obtained by dividing  $360^\circ$  by its speed. The periods and corresponding speeds of the constituents are derived from astronomical data and are independent of the locality of the tide station. The symbols  $\alpha$ ,  $\beta$ ,  $\gamma$ , etc. refer to the initial phases of the constituent angles at the time when  $t$  equals zero. The initial phases depend upon locality as well as the instant from which the time is reckoned and their values are derived from tidal observations. *Harmonic analysis* as applied to tides is the process by which the observed tidal data at any place are separated into a number of harmonic constituents. The quantities sought are known as *harmonic constants* and consist of the amplitudes and certain phase relations which will be more fully explained later. *Harmonic prediction* is accomplished by reuniting the elementary constituents in accordance with astronomical relations prevailing at the time for which the predictions are being made.

#### ASTRONOMICAL DATA

9. In tidal work the only celestial bodies that need be considered are the moon and sun. Although every other celestial body whose gravitational influence reaches the earth creates a theoretical tide-producing force, the greater distance or smaller size of such body renders negligible any effect of this force upon the tides of the earth. In deriving mathematical expressions for the tide-producing forces of the moon and sun, the principal factors to be taken into consideration are the rotation of the earth, the revolution of the moon around the earth, the revolution of the earth around the sun, the inclination of the moon's orbit to the earth's equator, and the obliquity of the ecliptic. Numerical values pertaining to these factors will be found in table 1.

10. The earth rotates on its axis once each day. There are, however, several kinds of days—the sidereal day, the solar day, the lunar day, and the constituent day—depending upon the object used as a reference for the rotation. The *sidereal day* is defined by astronomers as the time required for the rotation of the earth with respect to the vernal equinox. Because of the precession of the equinox, this day differs slightly from the time of rotation with respect to a fixed star, the difference being less than the hundredth part of a second. The *solar day* and *lunar day* are respectively the times required for rotation with respect to the sun and moon. Since the motions of the earth and moon in their orbits are not uniform, the solar and lunar days vary a little in length and their average or mean values are taken as standard units of time. A *constituent day* is the time of the rotation of the earth with respect to a fictitious satellite representing one of the periodic elements in the tidal forces. It approximates in length the lunar or solar day and corresponds to the period of a diurnal constituent or twice the period of a semidiurnal constituent.

11. A *calendar day* is a mean solar day commencing at midnight. Such a calendar day is known also as a *civil day* to distinguish it from the *astronomical day* which commences at noon of the same date.

Prior to the year 1925, the astronomical day was in general use by astronomers for the recording of astronomical data, but beginning with the Ephemeris and Nautical Almanac published in 1925 the civil day has been adopted for the calculations. Each day of whatever kind may be divided into 24 equal parts known as hours which are qualified by the name of the kind of day of which they are a part, as *sidereal hour, solar hour, lunar hour, or constituent hour*.

12. The moon revolves around the earth in an elliptical orbit. Although the average eccentricity of this orbit remains approximately constant for long periods of time, there are a number of perturbations in the moon's motion due, primarily, to the attractive force of the sun. Besides the revolution of the line of apsides and the regression of the nodes which take place more or less slowly, the principal inequalities in the moon's motion which affect the tides are the evection and variation. The evection depends upon the alternate increase and decrease of the eccentricity of the moon's orbit, which is always a maximum when the sun is passing the moon's line of apsides, and a minimum when the sun is at right angles to it. The variation inequality is due mainly to the tangential component of the disturbing force. The period of the revolution of the moon around the earth is called a month. The month is designated as sidereal, tropical, anomalistic, nodical, or synodical, according to whether the revolution is relative to a fixed star, the vernal equinox, the perigee, the ascending node, or the sun. The calendar month is a rough approximation to the synodical month.

13. It is customary to refer to the revolution of the earth around the sun, although it may be more accurately stated that they both revolve around their common center of gravity; but if we imagine the earth as fixed, the sun will describe an apparent path around the earth which is the same in size and form as the orbit of the earth around the sun, and the effect upon the tides would be the same. This orbit is an ellipse with an eccentricity that changes so slowly that it may be considered as practically constant. The period of the revolution of the earth around the sun is a year, but there are several kinds of years. The *sidereal year* is a revolution with respect to a fixed star, the *tropical year* is a revolution with respect to the vernal equinox, the *eclipse year* is a revolution with respect to the moon's ascending node, and the *anomalistic year* is a revolution with respect to the solar perigee.

14. A *calendar year* consists of an integral number of mean solar days and may be a *common year* of 365 days or a *leap year* of 366 days, these years being selected according to the calendars described below so that the average length will agree as nearly as practicable with the length of the tropical year which fixes the periodic changes in the seasons. The average length of the calendar year by the Julian calendar is exactly 365.25 days and by the Gregorian calendar 365.2425 days and these may be designated respectively as a *Julian year* and a *Gregorian year*.

15. The two principal kinds of calendars in use by most of the civilized world since the beginning of the Christian era are the Julian and the Gregorian calendars, the latter being the modern calendar in which the dates are sometimes referred to as "new style" to distinguish them from the dates of the older calendars. Prior to the year 45 B. C. there was more or less confusion in the calendars, inter-

calations of months and days being arbitrarily made by the priesthood and magistrates to bring the calendar into accord with the seasons and for other purposes.

16. The Julian calendar received its name from Julius Cæsar, who introduced it in the year 45 B. C. This calendar provided that the common year should consist of 365 days and every fourth year of 366 days, each year to begin on January 1. As proposed by Julius Cæsar, the 12 months beginning with January were to be alternately 31 days and 30 days in length with the exception that February should have only 29 days in the common years. When Augustus succeeded Julius Cæsar a few years later, he slightly modified this arrangement by transferring one day from February to the month of Sextilis, or August as it was then renamed, and also transferred the 31st day of September and November to October and December to avoid having three 31-day months in succession.

17. The Gregorian calendar received its name from Pope Gregory, who introduced it in the year 1582. It was immediately adopted by the Catholic countries but was not accepted by England until 1752. This calendar differs from the Julian calendar in having the century years not exactly divisible by 400 to consist of only 365 days, while in the Julian calendar every century year as well as every other year divisible by 4 is taken as a leap year with 366 days. For dates before Christ the year number must be diminished by 1 before testing its divisibility by 4 or 400 since the year 1 B. C. corresponds to the year 0 A. D. The Gregorian calendar will gain on the Julian calendar three days in each 400 years. When originally adopted, in order to adjust the Gregorian calendar so that the vernal equinox should fall upon March 21, as it had at the time of the Council of Nice in 325 A. D., 10 days were dropped and it was ordered that the day following October 4, 1582 of the Julian calendar should be designated as October 15, 1582 of the Gregorian calendar. This difference of 10 days between the dates of the two calendars continued until 1700, which was a leap year according to the Julian calendar and a common year by the Gregorian calendar. The difference between the two then became 11 days and in 1800 was increased to 12 days. Since 1900 the difference has been 13 days and will remain the same until the year 2100.

18. Dates of the Christian era prior to October 4, 1582, will, in general, conform to the Julian calendar. Since that time both calendars have been used. The Gregorian calendar was adopted in England by an act of Parliament passed in 1751, which provided that the day following September 2, 1752, should be called September 14, 1752, and also that the year 1752 and subsequent years should commence on the 1st day of January. Previous to this the legal year in England commenced on March 25. Except for this arbitrary beginning of the year, the old English calendar was the same as the Julian calendar. When Alaska was purchased from Russia by the United States, its calendar was altered by 11 days, one of these days being necessary because of the difference between the Asiatic and American dates when compared across the one hundred and eightieth meridian. Dates in the tables at the back of this volume refer to the Gregorian calendar.

19. The three great circles formed by the intersections of the planes of the earth's equator, the ecliptic, and the moon's orbit with the

celestial sphere are represented in figure 1. These circles intersect in six points, three of them being marked by symbols in the figure, namely, the *vernal equinox*  $\tau$  at the intersection of the celestial equator and ecliptic, the ascending *lunar node*  $\Omega$  at the intersection of the ecliptic and the projection of the moon's orbit, and the *lunar intersection*  $A$  at the intersection of the celestial equator and the projection of the moon's orbit. For brevity these three points are sometimes called respectively "the equinox," "the node," and "the intersection." The vernal equinox, although subject to a slow westward motion of about  $50''$  per year, is generally taken as a fixed point of reference for the motion of other parts of the solar system. The moon's node has a westward motion of about  $19^\circ$  a year, which is sufficient to carry it entirely around a great circle in a little less than 19 years.

**20.** The angle  $\omega$  between the ecliptic and the celestial equator is known as the obliquity of the ecliptic and has a nearly constant value of  $23\frac{1}{2}^\circ$ . The angle  $i$  between the ecliptic and the plane of the moon's orbit is also constant with a value of about  $5^\circ$ .




FIGURE 1.

The angle  $I$  which measures the inclination of the moon's orbit to the celestial equator might appropriately be called the obliquity of the moon's orbit. Its magnitude changes with the position of the moon's node. When the moon's ascending node coincides with the vernal equinox, the angle  $I$  equals the sum of  $\omega$  and  $i$ , or about  $28\frac{1}{2}^\circ$ , and when the descending node coincides with the vernal equinox, the angle  $I$  equals the difference between  $\omega$  and  $i$ , or about  $18\frac{1}{2}^\circ$ . This variation in the obliquity of the moon's orbit with its period of approximately 18.6 years introduces an important inequality in the tidal movement which must be taken into account.

**21.** In the celestial sphere the terms "latitude" and "longitude" apply especially to measurements referred to the ecliptic and vernal equinox, but the terms may with propriety also be applied to measurements referred to other great circles and origins, provided they are sufficiently well defined to prevent any ambiguity. For example, we may say "longitude in the moon's orbit measured from the moon's

node." Celestial longitude is always understood to be measured toward the east entirely around the circle. Longitude in the celestial equator reckoned from the vernal equinox is called right ascension, and the angular distance north or south of the celestial equator is called declination.

**22.** The true longitude of any point referred to any great circle in the celestial sphere may be defined as the arc of that circle intercepted between the accepted origin and the projection of the point on the circle, the measurement being always eastward from the origin to the projection of the point. The true longitude of any point will generally be different when referred to different circles, although reckoned from a common origin; and the longitude of a body moving at a uniform rate of speed in one great circle will not have a uniform rate of change when referred to another great circle.

**23.** The mean longitude of a body moving in a closed orbit and referred to any great circle may be defined as the longitude that would be attained by a point moving uniformly in the circle of reference at the same average angular velocity as that of the body and with the initial position of the point so taken that its mean longitude would be the same as the true longitude of the body at a certain selected position of that body in its orbit. With a common initial point, the mean longitude of a moving body will be the same in whatever circle it may be reckoned. Longitude in the ecliptic and in the celestial equator are usually reckoned from the vernal equinox  $\Upsilon$ , which is common to both circles. In order to have an equivalent origin in the moon's orbit, we may lay off an arc  $\Omega \Upsilon'$  (fig. 1) in the moon's orbit equal to  $\Omega \Upsilon$  in the ecliptic and for convenience call the point  $\Upsilon'$  the referred equinox. The mean longitude of any body, if reckoned from either the equinox or the referred equinox, will be the same in any of the three orbits represented. This will, of course, not be the case for the true longitude.

**24.** Let us now examine more closely the spherical triangle  $\Omega \Upsilon A$  in figure 1. The angles  $\omega$  and  $i$  are very nearly constant for long periods of time and have already been explained. The side  $\Omega \Upsilon$ , usually designated by  $N$ , is the longitude of the moon's node and is undergoing a constant and practically uniform change due to the regression of the moon's nodes. This westward movement of the node, by which it is carried completely around the ecliptic in a period of approximately 18.6 years, causes a constant change in the form of the triangle, the elements of which are of considerable importance in the present discussion. The value of the angle  $I$ , the supplement of the angle  $\Omega A \Upsilon$ , has an important effect upon both the range and time of the tide, which will be noted later. The side  $A \Upsilon$ , designated by  $\nu$ , is the right ascension or longitude in the celestial equator of the intersection  $A$ . The arc designated by  $\xi$  is equal to the side  $\Omega \Upsilon$ —side  $\Omega A$  and is the longitude in the moon's orbit of the intersection  $A$ . Since the angles  $i$  and  $\omega$  are assumed to be constant, the values of  $I$ ,  $\nu$ , and  $\xi$  will depend directly upon  $N$ , the longitude of the moon's node, and may be readily obtained by the ordinary solution of the spherical triangle  $\Omega \Upsilon A$ . Table 6 give the values of  $I$ ,  $\nu$ , and  $\xi$  for each degree of  $N$ . In the computation of this table the value of  $\omega$  for the beginning of the twentieth century was used. However, the secular change in the obliquity of the ecliptic is so slow that a difference of a century in

the epoch taken as the basis of the computation would have resulted in differences of less than 0.02 of a degree in the tabular values. The table may therefore be used without material error for reductions pertaining to any modern time.

25. Looking again at figure 1, it will be noted that when the longitude of the moon's node is zero the value of the inclination  $I$  will equal the sum of  $\omega$  and  $i$  and will be at its maximum. In this position the northern portion of the moon's orbit will be north of the ecliptic. When the longitude of the moon's node is  $180^\circ$ , the moon's orbit will be between the Equator and ecliptic, and the angle  $I$  will be equal to angle  $\omega$ —angle  $i$ . The angle  $I$  will be always positive and will vary from  $\omega-i$  to  $\omega+i$ . When the longitude of the moon's node equals zero or  $180^\circ$ , the values of  $\nu$  and  $\xi$  will each be zero. For all positions of the moon's node north of the Equator as its longitude changes from  $180$  to  $0^\circ$ ,  $\nu$  and  $\xi$  will have positive values, as indicated in the figure, these arcs being considered as positive when reckoned eastward from  $\Upsilon$  and  $\Upsilon'$ , respectively. For all positions of the node south of the Equator, as the longitude changes from  $360$  to  $180^\circ$ ,  $\nu$  and  $\xi$  will each be negative, since the intersection  $A$  will then lay to the westward of  $\Upsilon$  and  $\Upsilon'$ .

#### DEGREE OF APPROXIMATION

26. The problem of finding expressions for tidal forces and the equilibrium height of the tide in terms of time and place does not admit of a strict solution, but approximate expressions can be obtained which may be carried to as high an order of precision as desired. In ordinary numerical computations exact results are seldom obtained, the degree of precision depending upon the number of decimal places used in the computations, which, in turn, will be determined largely by the magnitude of the quantity sought. In general, the degree of approximation to the value of any quantity expressed numerically will be determined by the number of significant figures used. With a quantity represented by a single significant figure, the error may be as great as  $33\frac{1}{3}$  percent of the quantity itself, while the use of two significant figures will reduce the maximum error to less than 5 percent of the true value of the quantity. The large possible error in the first case renders it of little value, but in the latter case the approximation is sufficiently close to be useful when only rough results are necessary. The distance of the sun from the earth is popularly expressed by two significant figures as 93,000,000 miles.

27. With three or four significant figures fairly satisfactory approximations may be represented, and with a greater number very precise results may be expressed. For theoretical purposes the highest attainable precision is desirable, but for practical purposes, because of the increase in the labor without a corresponding increase in utility, it will be usually found advantageous to limit the degree of precision in accordance with the prevailing conditions.

28. Frequently a quantity that is to be used as a factor in an expression may be expanded into a series of terms. If the approximate value of such a series is near unity, terms which would affect the third decimal place, if expressed numerically, should usually be retained. The retention of the smaller terms will depend to some ex-

tent upon the labor involved since their rejection would not seriously affect the final results.

29. The formulas for the moon's true longitude and parallax on pages 19-20 are said to be given to the second order of approximation, a fraction of the first order being considered as one having an approximate value of  $1/20$  or  $0.05$ , a fraction of the second order having an approximate value of  $(0.05)^2$  or  $0.0025$ , a fraction of third order having an approximate value of  $(0.05)^3$  or  $0.000125$ , etc. As these formulas provide important factors in the development of the equations representing the tide-producing forces, they determine to a large extent the degrees of precision to be expected in the results.

## DEVELOPMENT OF TIDE-PRODUCING FORCE

### FUNDAMENTAL FORMULAS

30. The tide-producing forces exerted by the moon and sun are similar in their action and mathematical expressions obtained for one may therefore by proper substitutions be adapted to the other. Because of the greater importance of the moon in its tide-producing effects, the following development will apply primarily to that body, the necessary changes to represent the solar tides being afterwards indicated.

31. The tide-producing force of the moon is that portion of its gravitational attraction which is effective in changing the water level on the earth's surface. This effective force is the difference between the attraction for the earth as a whole and the attraction for the different particles which constitute the yielding part of the earth's surface; or, if the entire earth were considered to be a plastic mass, the tide-producing force at any point within the mass would be the force that tended to change the position of a particle at that point relative to a particle at the center of the earth. That part of the earth's surface which is directly under the moon is nearer to that body than is the center of the earth and is therefore more strongly attracted since the force of gravity varies inversely as the square of the distance. For the same reason the center of the earth is more strongly attracted by the moon than is that part of the earth's surface which is turned away from the moon.

32. The tide-producing force, being the difference between the attraction for particles situated relatively near together, is small compared with the attraction itself. It may be interesting to note that, although the sun's attraction on the earth is nearly 200 times as great as that of the moon, its tide-producing force is less than one-half that of the moon. If the forces acting upon each particle of the earth were equal and parallel, no matter how great those forces might be, there would be no tendency to change the relative positions of those particles, and consequently there would be no tide-producing force.

33. The tide-producing force may be graphically represented as in figure 2.

Let  $O$ =the center of the earth,

$C$ =the center of the moon,

$P$ =any point within or on the surface of the earth.

Then  $OC$  will represent the direction of the attractive force of the moon upon a particle at the center of the earth and  $PC$  the direction of the attractive force of the moon upon a particle at  $P$ . Now, let the magnitude of the moon's attraction at  $P$  be represented by the length of the line  $PC$ . Then, since the attraction of gravitation varies inversely as the square of the distance, it is necessary, in order to represent the attraction at  $O$  on the same scale, to take a line  $CQ$  of such length that  $CQ : CP = \overline{CP^2} : \overline{CO^2}$ .

**34.** The line  $PQ$ , joining  $P$  and  $Q$ , will then represent the direction and magnitude of the resultant force that tends to disturb the position of  $P$  relative to  $O$ , for it represents the difference between the force  $PC$  and a force through  $P$  equal and parallel to the force  $QC$  which acts upon  $O$ . This last statement may be a little clearer to the reader if he will consider the force  $PC$  as being resolved into a force  $PD$  equal and parallel to  $QC$ , and the force  $PQ$ . The force  $PD$ , acting upon the particle at  $P$ , being equal and parallel to the force  $QC$ , acting upon a particle at  $O$ , will have no tendency to change the position of  $P$  relative to  $O$ . The remaining force  $PQ$  will tend to alter the position of  $P$  relative to  $O$  and is the tide-producing force of the moon at  $P$ . The force  $PQ$  may be resolved into a vertical component  $PR$ , which tends to raise the water at  $P$ , and the horizontal component  $PT$ , which tends to move the water horizontally.




FIGURE 2.

**35.** If the point  $P'$  is taken so that the distance  $CP'$  is greater than the distance  $CO$ , the tide-producing force  $P'Q'$  will be directed away from the moon. While at first sight this may appear paradoxical, it will be noted that the moon tends to separate  $O$  from  $P'$ , but as  $O$  is taken as the point of reference, this resulting force that tends to separate the points is considered as being applied at the point  $P'$  only.

**36.** To express the tide-producing force by mathematical equations, refer to figure 2 and let

$$\begin{aligned} r &= OP = \text{distance of particle } P \text{ from center of earth,} \\ b &= PC = \text{distance of particle } P \text{ from center of moon,} \\ d &= OC = \text{distance from center of earth to center of moon,} \\ z &= COP = \text{angle at center of earth between } OP \text{ and } OC. \end{aligned}$$

Also let

$$\begin{aligned} M &= \text{mass of moon,} \\ E &= \text{mass of earth,} \\ a &= \text{mean radius of earth,} \\ \mu &= \text{attraction of gravitation between unit masses at unit distance.} \\ g &= \text{mean acceleration of gravity on earth's surface.} \end{aligned}$$

Since the force of gravitation varies directly as the mass and inversely as the square of the distance,

$$\text{Attraction of moon for unit mass at point } O \text{ in direction } OC = \frac{\mu M}{d^2} \quad (2)$$

$$\text{Attraction of moon for unit mass at point } P \text{ in direction } PC = \frac{\mu M}{b^2} \quad (3)$$

37. Let each of these forces be resolved into a vertical component along the radius  $OP$  and a horizontal component perpendicular to the same in the plane  $OPC$ , and consider the direction from  $O$  toward  $P$  as positive for the vertical component and the direction corresponding to the azimuth of the moon as positive for the horizontal component. We then have from (2) and (3)

$$\text{Attraction at } O \text{ in direction } O \text{ to } P = \frac{\mu M}{d^2} \cos z \quad (4)$$

$$\text{Attraction at } O \text{ perpendicular to } OP = \frac{\mu M}{d^2} \sin z \quad (5)$$

$$\text{Attraction at } P \text{ in direction } O \text{ to } P = \frac{\mu M}{b^2} \cos CPR \quad (6)$$

$$\text{Attraction at } P \text{ perpendicular to } OP = \frac{\mu M}{b^2} \sin CPR \quad (7)$$

38. The tide-producing force of the moon at any point  $P$  is measured by the difference between the attraction at  $P$  and at the center of the earth. Letting

$F_v$ =vertical component of tide-producing force, and

$F_a$ =horizontal component in azimuth of moon,

and taking the differences between (6) and (4) and between (7) and (5), we obtain the following expressions for these component forces in terms of the unit  $\mu$ :

$$F_v / \mu = M \left( \frac{\cos CPR}{b^2} - \frac{\cos z}{d^2} \right) \quad (8)$$

$$F_a / \mu = M \left( \frac{\sin CPR}{b^2} - \frac{\sin z}{d^2} \right) \quad (9)$$

39. From the plane triangle  $COP$  the following relations may be obtained:

$$b^2 = r^2 + d^2 - 2rd \cos z = d^2[1 - 2(r/d) \cos z + (r/d)^2] \quad (10)$$

$$\sin CPR = \sin CPO = (d/b) \sin z = \frac{\sin z}{[1 - 2(r/d) \cos z + (r/d)^2]^{\frac{1}{2}}} \quad (11)$$

$$\cos CPR = (1 - \sin^2 CPR)^{\frac{1}{2}} = \frac{\cos z - r/d}{[1 - 2(r/d) \cos z + (r/d)^2]^{\frac{1}{2}}} \quad (12)$$

40. In figure 2 it will be noted that the value of  $z$ , being reckoned in any plane from the line  $OC$ , may vary from zero to  $180^\circ$ , and also that the angle  $CPR$  increases as  $z$  increases within the same limits.  $\sin z$  and  $\sin CPR$  will therefore always be positive. As the angle  $OCP$  is always very small, the angle  $CPR$  will differ by only a very small amount from the angle  $z$  and will usually be in the same quadrant. In obtaining the square root for the numerator of (12) it was therefore necessary to use only that sign which would preserve this

relationship. The denominators of (11) and (12) are to be considered as positive.

41. Substituting in equations (8) and (9) the equivalents for  $b$ ,  $\sin CPR$ , and  $\cos CPR$  from equations (10) to (12), the following basic formulas are obtained for the vertical and horizontal components of the tide-producing force at any point  $P$  at  $r$  distance from the center of the earth:

$$F_v/\mu = \frac{M}{d^2} \left[ \frac{\cos z - r/d}{\{1 - 2(r/d) \cos z + (r/d)^2\}^{3/2}} - \cos z \right] \quad (13)$$

$$F_a/\mu = \frac{M}{d^2} \left[ \frac{\sin z}{\{1 - 2(r/d) \cos z + (r/d)^2\}^{3/2}} - \sin z \right] \quad (14)$$

42. To express these forces in their relation to the mean acceleration of gravity on the earth's surface, represented by the symbol  $g$ , we have

$$g/\mu = E/a^2, \text{ or } \mu/g = a^2/E \quad (15)$$

in which  $E$  is the mass and  $a$  is the mean radius of the earth. Substituting the above in formulas (13) and (14), we may write

$$F_v/g = (M/E) (a/d)^2 \left[ \frac{\cos z - r/d}{\{1 - 2(r/d) \cos z + (r/d)^2\}^{3/2}} = \cos z \right] \quad (16)$$

$$F_a/g = (M/E) (a/d)^2 \left[ \frac{\sin z}{\{1 - 2(r/d) \cos z + (r/d)^2\}^{3/2}} - \sin z \right] \quad (17)$$

43. Formulas (16) and (17) represent completely the vertical and horizontal components of the lunar tide-producing force at any point in the earth. If  $r$  is taken equal to the mean radius  $a$ , the formulas will involve the constant ratio  $M/E$  and two variable quantities—the angle  $z$  which is the moon's zenith distance, and the ratio  $a/d$  which is the sine of the moon's horizontal parallax in respect to the mean radius of the earth. Because of the smallness of the ratio  $a/d$  it may also be taken as the parallax itself expressed as a fraction of a radian. The parallax is largest when the moon is in perigee and at this time the tide-producing force will reach its greatest magnitude. A more rapid change in the tidal force at any point on the earth's surface is caused by the continuous change in the zenith distance of the moon resulting from the earth's rotation. The vertical component attains its maximum value when  $z$  equals zero, and the horizontal component has its maximum value when  $z$  is a little less than  $45^\circ$ . Substituting numerical values in formulas (16) and (17) and in similar formulas for the tide-producing force of the sun, the following are obtained as the approximate extreme component forces when the moon and sun are nearest the earth:

$$\text{Greatest } F_v/g = .144 \times 10^{-6} \text{ for moon, or } .054 \times 10^{-6} \text{ for sun} \quad (18)$$

$$\text{Greatest } F_a/g = .107 \times 10^{-6} \text{ for moon, or } .041 \times 10^{-6} \text{ for sun} \quad (19)$$

The horizontal component of the tide-producing force may be measured by its deflection of the plumb line, the relation of this component to gravity as expressed by the above formula being the tangent of the angle of deflection. Under the most favorable conditions the

greatest deflection due to the moon is about  $0.022''$  and the greatest deflection due to the sun is less than  $0.009''$  of arc.

44. To simplify the preceding formulas, the quantity involving the fractional exponent may be developed by Maclaurin's theorem into a series arranged according to the ascending powers of  $r/d$ , this being a small fraction with an approximate maximum value of 0.018. Thus

$$\frac{1}{\{1-2(r/d) \cos z + (r/d)^2\}^{\frac{1}{2}}} = 1 + 3 \cos z (r/d) \\ + 3/2 (5 \cos^2 z - 1) (r/d)^2 \\ + 5/2 (7 \cos^3 z - 3 \cos z) (r/d)^3 + \text{etc.} \quad (20)$$

45. Substituting (20) in formulas (16) and (17) and neglecting the higher powers of  $r/d$ , we obtain the following formulas:

$$F_v/g = 3 (M/E)(a/d)^2 (\cos^2 z - 1/3) (r/d) \\ + 3/2 (M/E)(a/d)^2 (5 \cos^3 z - 3 \cos z) (r/d)^2 \quad (21)$$

$$F_a/g = 3/2 (M/E)(a/d)^2 (\sin 2z) (r/d) \\ + 3/2 (M/E)(a/d)^2 \sin z (5 \cos^2 z - 1) (r/d)^2 \quad (22)$$

46. If  $r$ , which represents the distance of the point of observation for the center of the earth, is replaced by the mean radius  $a$ , it will be noted that the first term of each of the above formulas involves the cube of the ratio  $a/d$  while the second term involves the fourth power of this quantity. This ratio is essentially the moon's parallax expressed in the radian unit. These terms may now be written as separate formulas and for convenience of identification the digits "3" and "4" will be annexed to the formula symbol to represent respectively the terms involving the cube and fourth power of the parallax. Thus

$$F_{v3}/g = 3 (M/E)(a/d)^3 (\cos^2 z - 1/3) \quad (23)$$

$$F_{v4}/g = 3/2 (M/E)(a/d)^4 (5 \cos^3 z - 3 \cos z) \quad (24)$$

$$F_{a3}/g = 3/2 (M/E)(a/d)^3 \sin 2z \quad (25)$$

$$F_{a4}/g = 3/2 (M/E)(a/d)^4 \sin z (5 \cos^2 z - 1) \quad (26)$$

Formulas (23) and (25) involving the cube of the parallax represent the principal part of the tide-producing force. For the moon this is about 98 per cent of the whole and for the sun a higher percentage. The part of the tide-producing force represented by formulas (24) and (26) and involving the fourth power of the parallax is of very little practical importance but as a matter of theoretical interest will be later given further attention.

47. An examination of formulas (23) and (25) shows that the principal part of the tide-producing force is symmetrically distributed over the earth's surface with respect to a plane through the center of the earth and perpendicular to a line joining the centers of the earth and moon. The vertical component (23) has a maximum positive value when the zenith distance  $z=0$  or  $180^\circ$  and a maximum negative value when  $z=90^\circ$ , the maximum negative value being one-half as great as the maximum positive value. The vertical component be-

comes zero when  $z = \cos^{-1} \pm \sqrt{1/3}$  (approx.  $54.74^\circ$  and  $125.26^\circ$ ). The horizontal component (25) has its maximum value when  $z = 45^\circ$  and an equal maximum negative value when  $z = 135^\circ$ . The horizontal component becomes zero when  $z = 0, 90^\circ$ , or  $180^\circ$ .

48. If numerical values applicable to the mean parallax of the moon are substituted in (23) and (25), these component forces may be written

$$F_{v3}/g \text{ at mean parallax} = 0.000,000,167 (\cos^2 z - 1/3) \quad (27)$$

$$F_{a3}/g \text{ at mean parallax} = 0.000,000,084 \sin 2z \quad (28)$$

For the corresponding components of the solar tide-producing force, the numerical coefficients will be 0.46 times as great as those in the above formulas.

49. For the extreme values of the components represented by (23) and (25), with the moon and sun nearest the earth, the following may be obtained by suitable substitutions:

$$\text{Greatest } F_{v3}/g = .140 \times 10^{-6} \text{ for moon, or } .054 \times 10^{-6} \text{ for sun} \quad (29)$$

$$\text{Greatest } F_{a3}/g = .105 \times 10^{-6} \text{ for moon, or } .041 \times 10^{-6} \text{ for sun} \quad (30)$$

Comparing the above with (18) and (19), it will be noted that the maximum values of the lunar components involving the cube of the moon's parallax are only slightly less than the corresponding maximum values for the entire lunar force, while for the solar components the differences are too small to be shown with the number of decimal places used.

#### VERTICAL COMPONENT OF FORCE

50. It is now proposed to expand into a series of harmonic terms formula (23) which represents the principal vertical component of the lunar tide-producing force. In figure 3 let  $O$  represent the center of the earth and let projections on the celestial sphere be as follows:

- $C$ , the north pole
- $I M' P'$ , the earth's equator
- $I M$ , the moon's orbit
- $M$ , the position of the moon
- $P$ , the place of observation
- $C' M M'$ , the hour circle of the moon
- $C P P'$ , the meridian of place of observation
- $I$ , the intersection of moon's orbit and equator

Also let

- $I = \text{angle } M I M' = \text{inclination of moon's orbit to earth's equator}$
- $t = \text{arc } P' M' \text{ or angle } PCM = \text{hour angle of moon}$
- $X = IP' = \text{longitude of } P \text{ measured in celestial equator from intersection } I$
- $j = IM = \text{longitude of moon in orbit reckoned from intersection } I$
- $z = PM = \text{zenith distance of moon}$
- $D = M'M = \text{declination of moon}$
- $Y = P'P = \text{latitude cf } P$

The solution of a number of the spherical triangles represented in figure 3 will provide certain relations needed in the development of the formulas for the tide-producing force.

51. In spherical triangle  $MCP$ , the angle  $C$  equals  $t$  and the sides  $MC$  and  $PC$  are the complements of  $D$  and  $Y$ , respectively. We may therefore write

$$\cos z = \sin Y \sin D + \cos Y \cos D \cos t \quad (31)$$

Substituting this value in formula (23), we obtain

$$\begin{aligned} F_{v3}/g = & 3/2 (M/E)(a/d)^3 (1/2 - 3/2 \sin^2 Y) (2/3 - 2 \sin^2 D) \dots F_{v30}/g \\ & + 3/2 (M/E)(a/d)^3 \sin 2Y \sin 2D \cos t \dots F_{v31}/g \\ & + 3/2 (M/E)(a/d)^3 \cos^2 Y \cos^2 D \cos 2t \dots F_{v32}/g \end{aligned} \quad (32)$$




FIGURE 3.

52. In formula (32) the vertical component of the tide-producing force has been separated into three parts. The first term is independent of the rotation of the earth but is subject to variations arising from changes in declination and distance of the moon. It includes what are known as the *long-period constituents*, that is to say, constituents with periods somewhat longer than a day and in general a half month or longer. The second term involves the cosine of the hour angle ( $t$ ) of the moon and this includes the *diurnal constituents* with periods approximating the lunar day. The last term involves the cosine of twice the hour angle of the moon and includes the *semidiurnal constituents* with periods approximating the half lunar day. The grouping of the tidal constituents according to their approximate periods affords an important classification in the further development of the tidal forces and these groups will be called *classes* or *species*. Symbols pertaining to a particular species are often identified by a subscript indicating the number of periods in a day,

the subscript o being used for the long-period constituents. In formula (32) the individual terms are identified by the annexation of the species subscript to the general symbol for the formula.

53. As written, all of the three terms of formula (32) have the same coefficient  $3/2 (M/E) (a/d)^3$ . In each case the latitude ( $Y$ ) factor has a maximum value of unity, this maximum being negative for the first term. For the long-period term ( $F_{e30}/g$ ), the latitude factor has a maximum positive value of  $\frac{1}{2}$  at the equator, becomes zero in latitude  $35.26^\circ$  (approximately), and reaches a maximum negative value of  $-1$  at the poles, the factor being the same for corresponding latitudes in both northern and southern hemispheres. For the diurnal term ( $F_{e31}/g$ ), the latitude factor is positive for the northern hemisphere and negative for the southern hemisphere. It has a maximum value of unity in latitude  $45^\circ$  and is zero at the equator and poles. For the semidiurnal terms ( $F_{e32}/g$ ), the latitude factor is always positive and has a maximum value of unity at the equator and equals zero at the poles.

54. For extreme values attainable for the declinational ( $D$ ) factors, consideration must be given to the greatest declination which can be reached by the tide-producing body. The periodic maximum declination reached by the moon in its 18.6 year node-cycle is  $28.6^\circ$  but this may be slightly increased by other inequalities in the moon's motion. The maximum declination for the sun, taken the same as the obliquity of the ecliptic, is  $23.45^\circ$ . The declinational factor of the long-period term ( $F_{e30}/g$ ) has a maximum value of  $2/3$  when the declination is zero. It diminishes with increasing north or south declination but must always remain positive because of the limits of the declination. For the diurnal term ( $F_{e31}/g$ ) the declinational factor has its greatest value when the declination is greatest. For the moon the maximum value of this factor is approximately 0.841 and for the sun 0.730. This factor is positive for the north declination and negative for the south declination. For the semidiurnal term ( $F_{e32}/g$ ) the declinational factor for both moon and sun is always positive and has a maximum value of unity at zero declination.

55. The greatest numerical values for the several terms of the vertical component of the tide-producing force as represented by formula (32) and applicable to the time when the moon and sun are nearest the earth, are as follows:

$$\text{Greatest } F_{e30}/g = -0.070 \times 10^{-6} \text{ for moon, or } -0.027 \times 10^{-6} \text{ for sun} \quad (33)$$

$$\text{Greatest } F_{e31}/g = \pm 0.088 \times 10^{-6} \text{ for moon, or } \pm 0.030 \times 10^{-6} \text{ for sun} \quad (34)$$

$$\text{Greatest } F_{e32}/g = +0.105 \times 10^{-6} \text{ for moon, or } +0.041 \times 10^{-6} \text{ for sun} \quad (35)$$

For the long-period term (33) the greatest value applies to either pole and is negative. For the diurnal term (34) the greatest value applies in latitude  $45^\circ$  and may be positive or negative according to whether the latitude and declinational factors have the same or opposite signs. For the semidiurnal term (35) the greatest value applies to the equator and is positive.

56. Referring to formula (32), let  $a/c$  equal the mean value of parallax  $a/d$ . Then  $a/d$  may be replaced by its equivalent  $(a/c)(c/d)$ , in which the fraction  $c/d$  expresses the relation between the true and, the mean parallax. Also let  $U = (M/E) (a/c)^3$ , the numerical value of which will be found in table 1. Expressing separately the three terms of formula (32), we then have

$$F_{v30}/g = 3/2 U (c/d)^3 (1/2 - 3/2 \sin^2 Y) (2/3 - 2 \sin^2 D) \quad (36)$$

$$F_{v31}/g = 3/2 U (c/d)^3 \sin 2Y \sin 2D \cos t \quad (37)$$

$$F_{v32}/g = 3/2 U (c/d)^3 \cos^2 Y \cos^2 D \cos 2t \quad (38)$$

57. Referring to figure 3, the following relations may be obtained from the right spherical triangles  $MIM'$  and  $MP'M'$  and the oblique spherical triangle  $MP'I$ :

$$\sin D = \sin I \sin j \quad (39)$$

$$\cos D \cos t = \cos MP' \quad (40)$$

$$\cos MP' = \cos X \cos j + \sin X \sin j \cos I \quad (41)$$

$$\begin{aligned} \cos D \cos t &= \cos X \cos j + \sin X \sin j \cos I \\ &= \cos^2 \frac{1}{2}I \cos (X-j) + \sin^2 \frac{1}{2}I \cos (X+j) \end{aligned} \quad (42)$$

58. Replacing the functions of  $D$  and  $t$  in formulas (36) to (38) by their equivalents derived from equations (39) and (42), there are obtained the following:

$$F_{v30}/g = 3/2 U (c/d)^3 (1/2 - 3/2 \sin^2 Y) \times [2/3 - \sin^2 I + \sin^2 I \cos 2j] \quad (43)$$

$$F_{v31}/g = 3/2 U (c/d)^3 \sin 2Y \times [\sin I \cos^2 \frac{1}{2}I \cos (X+90^\circ-2j) + 1/2 \sin 2I \cos (X-90^\circ) + \sin I \sin^2 \frac{1}{2}I \cos (X-90^\circ+2j)] \quad (44)$$

$$F_{v32}/g = 3/2 U (c/d)^3 \cos^2 Y \times [\cos^4 \frac{1}{2}I \cos (2X-2j) + 1/2 \sin^2 I \cos 2X + \sin^4 \frac{1}{2}I \cos (2X+2j)] \quad (45)$$

The above formulas involve the moon's actual distance  $d$  and its true longitude  $j$  as measured in its orbit from the intersection. While these are functions of time, they do not vary uniformly because of certain inequalities in the motion of the moon, and it is now desired to replace these quantities by elements that do change uniformly.

59. Referring to paragraphs 23-24 and to figure 1, it will be noted that longitude measured from intersection  $A$  in the moon's orbit equals the longitude measured from the referred equinox  $\Upsilon'$  less arc  $\xi$ , and longitude measured from intersection  $A$  in the celestial equator equals the longitude measured from the equinox  $\Upsilon$  less arc  $\nu$ . Now let

$$s' = \text{true longitude of moon in orbit referred to equinox}$$

$$s = \text{mean longitude of moon referred to equinox}$$

$$k = \text{difference } (s' - s)$$

Then

$$j = s' - \xi = s - \xi + k \quad (46)$$

60. In figure 4 let  $S'$  and  $P'$  be the points where the hour circles of the mean sun and place of observation intersect the celestial equator,  $\Upsilon$  the vernal equinox, and  $I$  the lunar intersection. Then  $X$  will equal the arc  $P'I$  and  $\nu$  the arc  $I\Upsilon$ . Now let

$$h = \text{mean longitude of sun}$$

$$T = \text{hour angle of mean sun}$$

Then

$$X = T + h - \nu \quad (47)$$

61. Substituting the values of  $j$  and  $X$  from (46) and (47) in formulas (43) to (45), these may be written

$$\begin{aligned} F_{v30}/g = & 3/2 U(1/2 - 3/2 \sin^2 Y) \times \\ & [(c/d)^3 (2/3 - \sin^2 I) \\ & + (c/d)^3 \sin^2 I \cos (2s - 2\xi + 2k)] \end{aligned} \quad (48)$$

$$\begin{aligned} F_{v31}/g = & 3/2 U \sin 2Y \times \\ & [(c/d)^3 \sin I \cos^2 \frac{1}{2}I \cos (T - 2s + h + 2\xi - \nu + 90^\circ - 2k) \\ & + 1/2 (c/d)^3 \sin 2I \cos (T + h - \nu - 90^\circ) \\ & + (c/d)^3 \sin I \sin^2 \frac{1}{2}I \cos (T + 2s + h - 2\xi - \nu - 90^\circ + 2k)] \end{aligned} \quad (49)$$

$$\begin{aligned} F_{v32}/g = & 3/2 U \cos^2 Y \times \\ & [(c/d)^3 \cos^4 \frac{1}{2}I \cos (2T - 2s + 2h + 2\xi - 2\nu - 2k) \\ & + 1/2 (c/d)^3 \sin^2 I \cos (2T + 2h - 2\nu) \\ & + (c/d)^3 \sin^4 \frac{1}{2}I \cos (2T + 2s + 2h - 2\xi - 2\nu + 2k)] \end{aligned} \quad (50)$$




FIGURE 4.

Disregarding at this time the slow change in the function of  $I$ , the variable part of each term of the above formulas may be expressed in one of the following forms— $(c/d)^3$ ,  $(c/d)^3 \cos A$ ,  $(c/d)^3 \cos (A+2k)$ , or  $(c/d)^3 \cos (A-2k)$ , in which  $A$  includes all the elements of the variable angular function excepting the multiple of  $k$ .

62. The following equations for the motion of the moon were adapted from Godfray's Elementary Treatise on the Lunar Theory:

$s'$  = true longitude of moon (in radians)

$$\begin{aligned} &= s - \dots \quad \text{(mean longitude)} \\ &+ 2e \sin(s-p) + 5/4 e^2 \sin 2(s-p) \dots \quad \text{(elliptic inequality)} \\ &+ 15/4 m e \sin(s-2h+p) \dots \quad \text{(evectional inequality)} \\ &+ 11/8 m^2 \sin 2(s-h) \dots \quad \text{(variational inequality)} \end{aligned} \quad (51)$$

$$\begin{aligned}
 c/d &= (\text{true parallax of moon})/(\text{mean parallax of moon}) \\
 &= \text{unity} \\
 &+ e \cos(s-p) + e^2 \cos 2(s-p) \quad \text{(elliptic inequality)} \\
 &+ 15/8 me \cos(s-2h+p) \quad \text{(evectional inequality)} \\
 &+ m^2 \cos 2(s-h) \quad \text{(variational inequality)} \quad (52)
 \end{aligned}$$

in which

$s'$  = true longitude of moon in orbit (referred to equinox)

$s$  = mean longitude of moon

$h$  = mean longitude of sun

$p$  = mean longitude of lunar perigee

$e$  = eccentricity of moon's orbit = 0.0549

$m$  = ratio of mean motion of sun to that of moon = 0.0748

The elements  $e$  and  $m$  are small fractions of the first order and the square of either or the product of both may be considered as being of the second order. In the following development the higher powers of these elements will be omitted.

**63.** Since  $k$  has been taken as the difference between the true and the mean longitude of the moon, we may obtain from (51)

$$\begin{aligned}
 k &= 2e \sin(s-p) + 5/4 e^2 \sin 2(s-p) \\
 &\quad + 15/4 me \sin(s-2h+p) + 11/8 m^2 \sin 2(s-h) \quad (53)
 \end{aligned}$$

The value of  $k$  is always small, its maximum value being about 0.137 radian. It may therefore be assumed without material error that the sine of  $k$  or the sine of  $2k$  is equal to the angle itself. Then

$$\begin{aligned}
 \sin 2k &= 2k = 4e \sin(s-p) + 5/2 e^2 \sin 2(s-p) \\
 &\quad + 15/2 me \sin(s-2h+p) + 11/4 m^2 \sin 2(s-h) \quad (54)
 \end{aligned}$$

$$\begin{aligned}
 \cos 2k &= 1 - 2 \sin^2 k = 1 - 2k^2 \\
 &= 1 - 4e^2 + 4e^2 \cos 2(s-p) \quad (55)
 \end{aligned}$$

terms smaller than those of the second order being omitted.

**64.** Cubing (52) and neglecting the smaller terms, we obtain

$$\begin{aligned}
 (c/d)^3 &= 1 + 3/2 e^2 + 3e \cos(s-p) + 9/2 e^2 \cos 2(s-p) \\
 &\quad + 45/8 me \cos(s-2h+p) + 3 m^2 \cos 2(s-h) \quad (56)
 \end{aligned}$$

Multiplying (54) and (55) by (56)

$$\begin{aligned}
 (c/d)^3 \sin 2k &= 4e \sin(s-p) + 17/2 e^2 \sin 2(s-p) \\
 &\quad + 15/2 me \sin(s-2h+p) + 11/4 m^2 \sin 2(s-h) \quad (57)
 \end{aligned}$$

$$\begin{aligned}
 (c/d)^3 \cos 2k &= 1 - 5/2 e^2 + 3e \cos(s-p) + 17/2 e^2 \cos 2(s-p) \\
 &\quad + 45/8 me \cos(s-2h+p) + 3 m^2 \cos 2(s-h) \quad (58)
 \end{aligned}$$

**65.** From (56), (57), and (58), we may obtain the following general expressions applicable to the further development of formulas (48) to (50). Negative coefficients have been avoided by the introduction of  $180^\circ$  in the angle when necessary.

$$\begin{aligned}
 (c/d)^3 \cos(A-2k) &= (c/d)^3 \cos 2k \cos A + (c/d)^3 \sin 2k \sin A \\
 &= (1 - 5/2 e^2) \cos A \\
 &\quad + 7/2 e \cos(A-s+p) + 1/2e \cos(A+s-p+180^\circ) \\
 &\quad + 17/2 e^2 \cos(A-2s+2p) \\
 &\quad + 105/16 me \cos(A-s+2h-p) + 15/16 me \cos(A+s-2h+p+180^\circ) \\
 &\quad + 23/8 m^2 \cos(A-2s+2h) + 1/8 m^2 \cos(A+2s-2h) \quad (59)
 \end{aligned}$$

$$(c/d)^3 \cos A = (1 + 3/2 e^2) \cos A + 3/2 e \cos (A - s + p) + 3/2 e \cos (A + s - p) + 9/4 e^2 \cos (A - 2s + 2p) + 9/4 e^2 \cos (A + 2s - 2p) + 45/16 me \cos (A - s + 2h - p) + 45/16 me \cos (A + s - 2h + p) + 3/2 m^2 \cos (A - 2s + 2h) + 3/2 m^2 \cos (A + 2s - 2h) \quad (60)$$

$$(c/d)^3 \cos (A + 2k) = (c/d)^3 \cos 2k \cos A - (c/d)^3 \sin 2k \sin A = (1 - 5/2 e^2) \cos A + 7/2 e \cos (A + s - p) + 1/2 e \cos (A - s + p + 180^\circ) + 17/2 e^2 \cos (A + 2s - 2p) + 105/16 me \cos (A + s - 2h + p) + 15/16 me \cos (A - s + 2h - p + 180^\circ) + 23/8 m^2 \cos (A + 2s - 2h) + 1/8 m^2 \cos (A - 2s + 2h) \quad (61)$$

**66.** After suitable substitutions for  $A$  have been made in the three preceding equations they are immediately applicable to the final expansion of the several terms in formulas (48) to (50), excepting the first term of (48) for which formula (56) may be used directly. Each term in the expanded formulas given below represents a constituent of the lunar tide-producing force and for convenience of reference is designated by the letter  $A$  with a subscript. There are also given the generally recognized symbols for the principal constituents, and when such a symbol is enclosed in brackets it signifies that the term given only partially represents the constituent so named.

**67.** Formula for long-period constituents of vertical component of principal lunar tide-producing force:

$$F_{v30} / g = 3/2 U(1/2 - 3/2 \sin^2 Y) \times \\ (A_1) [2(3 - \sin^2 I)\{(1 + 3/2 e^2) \dots \dots \text{permanent term} \\ (A_2) \quad + 3 e \cos (s - p) \dots \dots \text{Mm} \\ (A_3) \quad + 9/2 e^2 \cos (2s - 2p) \\ (A_4) \quad + 45/8 me \cos (s - 2h + p) \\ (A_5) \quad + 3 m^2 \cos (2s - 2h)\} \dots \dots \text{MSf} \\ (A_6) + \sin^2 I\{(1 - 5/2 e^2) \cos (2s - 2\xi) \dots \dots \text{Mf} \\ (A_7) \quad + 7/2 e \cos (3s - p - 2\xi) \\ (A_8) \quad + 1/2 e \cos (s + p + 180^\circ - 2\xi) \\ (A_9) \quad + 17/2 e^2 \cos (4s - 2p - 2\xi) \\ (A_{10}) \quad + 105/16 me \cos (3s - 2h + p - 2\xi) \\ (A_{11}) \quad + 15/16 me \cos (s + 2h - p + 180^\circ - 2\xi) \\ (A_{12}) \quad + 23/8 m^2 \cos (4s - 2h - 2\xi) \\ (A_{13}) \quad + 1/8 m^2 \cos (2h - 2\xi)\}] \quad (62)$$

**68.** Formula for diurnal constituents of vertical component of principal lunar tide-producing force:

$$F_{v31} / g = 3/2 U \sin 2Y \times \\ (A_{14}) [\sin I \cos^2 1/2 I \{(1 - 5/2 e^2) \cos (T - 2s + h + 90^\circ + 2\xi - \nu) \dots \dots O_1 \\ (A_{15}) \quad + 7/2 e \cos (T - 3s + h + p + 90^\circ + 2\xi - \nu) \dots \dots Q_1 \\ (A_{16}) \quad + 1/2 e \cos (T - s + h - p - 90^\circ + 2\xi - \nu) \dots \dots [M_1] \\ (A_{17}) \quad + 17/2 e^2 \cos (T - 4s + h + 2p + 90^\circ + 2\xi - \nu) \dots \dots 2Q_1 \\ (A_{18}) \quad + 105/16 me \cos (T - 3s + 3h - p + 90^\circ + 2\xi - \nu) \dots \dots \rho_1 \\ (A_{19}) \quad + 15/16 me \cos (T - s - h + p - 90^\circ + 2\xi - \nu) \\ (A_{20}) \quad + 23/8 m^2 \cos (T - 4s + 3h + 90^\circ + 2\xi - \nu) \dots \dots \sigma_1 \\ (A_{21}) \quad + 1/8 m^2 \cos (T - h + 90^\circ + 2\xi - \nu)\}]$$

(Formula continued next page)

$$\begin{aligned}
 (A_{22}) & +\sin 2I\{(1/2+3/4 e^2) \cos(T+h-90^\circ-\nu) \dots [K_1] \\
 (A_{23}) & +3/4 e \cos(T-s+h+p-90^\circ-\nu) \dots [M_1] \\
 (A_{24}) & +3/4 e \cos(T+s+h-p-90^\circ-\nu) \dots J_1 \\
 (A_{25}) & +9/8 e^2 \cos(T-2s+h+2p-90^\circ-\nu) \\
 (A_{26}) & +9/8 e^2 \cos(T+2s+h-2p-90^\circ-\nu) \\
 (A_{27}) & +45/32 me \cos(T-s+3h-p-90^\circ-\nu) \dots \chi_1 \\
 (A_{28}) & +45/32 me \cos(T+s-h+p-90^\circ-\nu) \dots \theta_1 \\
 (A_{29}) & +3/4 m^2 \cos(T-2s+3h-90^\circ-\nu) \dots MP_1 \\
 (A_{30}) & +3/4 m^2 \cos(T+2s-h-90^\circ-\nu)\} \dots SO_1 \\
 & +\sin I \sin^2 \frac{\gamma}{2} I \\
 (A_{31}) & \{(1-5/2 e^2) \cos(T+2s+h-90^\circ-2\xi-\nu) \dots OO_1 \\
 (A_{32}) & +7/2 e \cos(T+3s+h-p-90^\circ-2\xi-\nu) \dots KQ_1 \\
 (A_{33}) & +1/2 e \cos(T+s+h+p+90^\circ-2\xi-\nu) \\
 (A_{34}) & +17/2 e^2 \cos(T+4s+h-2p-90^\circ-2\xi-\nu) \\
 (A_{35}) & +105/16 me \cos(T+3s-h+p-90^\circ-2\xi-\nu) \\
 (A_{36}) & +15/16 me \cos(T+s+3h-p+90^\circ-2\xi-\nu) \\
 (A_{37}) & +23/8 m^2 \cos(T+4s-h-90^\circ-2\xi-\nu) \\
 (A_{38}) & +1/8 m^2 \cos(T+3h-90^\circ-2\xi-\nu)\} \quad (63)
 \end{aligned}$$

69. Formula for semidiurnal constituents of vertical component of principal lunar tide-producing force:

$$\begin{aligned}
 F_{v32} /g = & 3/2 U \cos^2 Y \times \\
 (A_{39}) & [\cos^4 \frac{1}{2} I \{(1-5/2 e^2) \cos(2T-2s+2h+2\xi-2\nu) \dots M_2 \\
 (A_{40}) & +7/2 e \cos(2T-3s+2h+p+2\xi-2\nu) \dots N_2 \\
 (A_{41}) & +1/2 e \cos(2T-s+2h-p+180^\circ+2\xi-2\nu) \dots [L_2] \\
 (A_{42}) & +17/2 e^2 \cos(2T-4s+2h+2p+2\xi-2\nu) \dots 2N_2 \\
 (A_{43}) & +105/16 me \cos(2T-3s+4h-p+2\xi-2\nu) \dots \nu_2 \\
 (A_{44}) & +15/16 me \cos(2T-s+p+180^\circ+2\xi-2\nu) \dots \lambda_2 \\
 (A_{45}) & +23/8 m^2 \cos(2T-4s+4h+2\xi-2\nu) \dots \mu_2 \\
 (A_{46}) & +1/8 m^2 \cos(2T+2\xi-2\nu)\} \\
 (A_{47}) & +\sin^2 I \{(1/2+3/4 e^2) \cos(2T+2h-2\nu) \dots [K_2] \\
 (A_{48}) & +3/4 e \cos(2T-s+2h+p-2\nu) \dots [L_2] \\
 (A_{49}) & +3/4 e \cos(2T+s+2h-p-2\nu) \dots KJ_2 \\
 (A_{50}) & +9/8 e^2 \cos(2T-2s+2h+2p-2\nu) \\
 (A_{51}) & +9/8 e^2 \cos(2T+2s+2h-2p-2\nu) \\
 (A_{52}) & +45/32 me \cos(2T-s+4h-p-2\nu) \\
 (A_{53}) & +45/32 me \cos(2T+s+p-2\nu) \\
 (A_{54}) & +3/4 m^2 \cos(2T-2s+4h-2\nu) \\
 (A_{55}) & +3/4 m^2 \cos(2T+2s-2\nu)\} \\
 (A_{56}) & +\sin^4 \frac{1}{2} I \{(1-5/2 e^2) \cos(2T+2s+2h-2\xi-2\nu) \\
 (A_{57}) & +7/2 e \cos(2T+3s+2h-p-2\xi-2\nu) \\
 (A_{58}) & +1/2 e \cos(2T+s+2h+p+180^\circ-2\xi-2\nu) \\
 (A_{59}) & +17/2 e^2 \cos(2T+4s+2h-2p-2\xi-2\nu) \\
 (A_{60}) & +105/16 me \cos(2T+3s+p-2\xi-2\nu) \\
 (A_{61}) & +15/16 me \cos(2T+s+4h-p+180^\circ-2\xi-2\nu) \\
 (A_{62}) & +23/8 m^2 \cos(2T+4s-2\xi-2\nu) \\
 (A_{63}) & +1/8 m^2 \cos(2T+4h-2\xi-2\nu)\} \quad (64)
 \end{aligned}$$

70. *Arguments.*—Except for the slow changes in the values of  $I$ ,  $\xi$ , and  $\nu$  which result from the revolution of the moon's node, each term other than the permanent one in the three preceding formulas is an harmonic function of an angle that changes uniformly with time. This angle is known as the *argument* of the constituent, also as the *equilibrium argument* when obtained in connection with the develop-

ment of the equilibrium tide. By analogy, the argument of the permanent term may be considered as zero, the cosine of zero being unity.

71. The argument serves to identify the constituent by determining its speed and period and fixing the times of the maxima and minima of the corresponding tidal force. It usually consists of two parts represented by the symbols  $V$  and  $u$ . When referring to a particular instant of time such as the beginning of a series of observations, the  $V$  is written with a subscript as  $V_0$ . The first part of the argument includes any constant and multiples of one or more of the following astronomical elements— $T$ , the hour angle of the mean sun at the place of observation;  $s$ , the mean longitude of the moon;  $h$  the mean longitude of the sun; and  $p$ , the longitude of the lunar perigee. The second part  $u$  includes multiples of one or both of the elements  $\xi$  and  $\nu$ , which are functions of the longitude of the moon's node and vary slowly between small positive and negative limits throughout a 19-year cycle. In a series of observations covering a year or less they are treated as constants with values pertaining to the middle of the series. They do not affect the average speed or period of the constituent. Their values corresponding to each degree of  $N$ , the longitude of the moon's node, are included in table 6, formulas for their computation being given on p. 156.

72. The hourly speed of a constituent may be obtained by adding the hourly speeds of the elements included in the  $V$  of the argument. These elementary speeds will be found in table 1. The period of a constituent is obtained by dividing  $360^\circ$  by its speed. The approximate period is determined by the element of greatest speed contained in the argument. Thus, the hour angle  $T$  has a speed of  $15^\circ$  per mean solar hour and all constituents with a single  $T$  in their arguments have periods approximating one day, while constituents with arguments containing the multiple  $2T$  have periods approximating the half day. Next to  $T$ , the element of greatest speed is  $s$  the mean longitude of the moon, and long-period constituents with a single  $s$  in their arguments will have periods approximating the month and with any multiple of  $s$  the corresponding fraction of a month. The arguments and speeds of the constituents are listed in table 2. Numerical values of the arguments for the beginning of each calendar year from 1850 to 2000 are given in table 15 for constituents used in the Coast and Geodetic Survey tide-predicting machine. Tables 16 to 18 provide differences for referring these arguments to any day and hour of the year.

73. In order to visualize the arguments of the constituents depending primarily upon the rotation of the earth, some have found it convenient to conceive of a system of fictitious stars, or "astres fictifs" as they are sometimes called, which move at a uniform rate in the celestial equator, each constituent being represented by a separate star. Thus, for the principal lunar constituent we have the mean moon and for the principal solar constituent the mean sun, while the various inequalities in the motions of these bodies are served by imaginary stars which reach the meridian of the place of observation at times corresponding to the zero value of the constituent argument. For the diurnal constituents the argument equals the hour angle of the star but for the semidiurnal constituents the argument is double the hour angle of the star.

**74.** *Coefficients.*—The complete coefficient of each term of formulas (62) to (64) includes several important factors. First, the *basic factor*  $U$ , which equals the ratio of the mass of the moon to that of the earth multiplied by the cube of the mean parallax of the moon, is common to all of the terms. This together with the common numerical coefficient may be designated as the *general coefficient*. Next, the function involving the latitude  $Y$  is known as the *latitude factor*, each formula having a different latitude factor. Following the latitude factor is a function of  $I$ , the inclination of the moon's orbit to the plane of the earth's equator, which may appropriately be called the *obliquity factor*, each factor applying to a group of terms. Lastly, we have an individual term coefficient which includes a numerical factor and involves the quantity  $e$  or  $m$ . Since these factors are derived from the equations of elliptic motion, they will here be referred to as *elliptic factors*. The product of the elliptic factor by the mean value of the obliquity factor is known as the *mean constituent coefficient* ( $C$ ). Numerical values for these coefficients are given in table 2. Since all terms in any one of the formulas have the same general coefficient and latitude factor, their relative magnitudes will be proportional to their constituent coefficients. Terms of different formulas, however, have different latitude factors and their constituent coefficients are not directly comparable without taking into account the latitude of the place of observation.

**75.** The obliquity factors are subject to variations throughout an 18.6-year cycle because of the revolution of the moon's node. During this period the value of  $I$  varies between the limits of  $\omega - i$  and  $\omega + i$ , or from  $18.3^\circ$  to  $28.6^\circ$  approximately, and the functions of  $I$  change accordingly. In order that tidal data pertaining to different years may be made comparable, it is necessary to adopt certain standard mean values for the obliquity factors to which results for different years may be reduced. While there are several systems of means which would serve equally well as standard values, the system adopted by Darwin in the early development of the harmonic analysis of tides has the sanction of long usage and is therefore followed. By the Darwin method, the mean for the obliquity factor is obtained from the product of the obliquity factor and the cosine of the elements  $\xi$  and  $\nu$  appearing in the argument. This may be expressed as the mean value of the product  $J \cos u$ , in which  $J$  is the function of  $I$  in the coefficient and  $u$  the function of  $\xi$  and  $\nu$  in the argument. Since  $u$  is relatively small and its cosine differs little from unity, the resulting mean will not differ greatly from the mean of  $J$  alone or from the function of  $I$  when given its mean value.

**76.** Using Darwin's system as described in section 6 of his paper on the Harmonic Analysis of Tidal Observations published in volume I of his collection of Scientific Papers (also in Report of the British Association for the Advancement of Science in 1883), the following mean values are obtained for the obliquity factors in formulas (62) to (64). These values were used in the computation of the corresponding constituent coefficients in table 2. The subscript  $\circ$  is here used to indicate the mean value of the function.

For terms  $A_1$  to  $A_5$  in formula (62)

$$[2/3 - \sin^2 I]_0 = (2/3 - \sin^2 \omega)(1 - 3/2 \sin^2 i) = 0.5021 \quad (65)$$

For terms  $A_6$  to  $A_{13}$  in formula (62)

$$[\sin^2 I \cos 2\xi]_0 = \sin^2 \omega \cos^4 \frac{1}{2}i = 0.1578 \quad (66)$$

For terms  $A_{14}$  to  $A_{21}$  in formula (63)

$$[\sin I \cos^2 \frac{1}{2}I \cos (2\xi - \nu)]_0 = \sin \omega \cos^2 \frac{1}{2}\omega \cos^4 \frac{1}{2}i = 0.3800 \quad (67)$$

For terms  $A_{22}$  to  $A_{30}$  in formula (63)

$$[\sin 2I \cos \nu]_0 = \sin 2\omega (1 - 3/2 \sin^2 i) = 0.7214 \quad (68)$$

For terms  $A_{31}$  to  $A_{38}$  in formula (63)

$$[\sin I \sin^2 \frac{1}{2}I \cos (2\xi + \nu)]_0 = \sin \omega \sin^2 \frac{1}{2}\omega \cos^4 \frac{1}{2}i = 0.0164 \quad (69)$$

For terms  $A_{39}$  to  $A_{46}$  in formula (64)

$$[\cos^4 \frac{1}{2}I \cos (2\xi - 2\nu)]_0 = \cos^4 \frac{1}{2}\omega \cos^4 \frac{1}{2}i = 0.9154 \quad (70)$$

For terms  $A_{47}$  to  $A_{55}$  in formula (64)

$$[\sin^2 I \cos 2\nu]_0 = \sin^2 \omega (1 - 3/2 \sin^2 i) = 0.1565 \quad (71)$$

For terms  $A_{56}$  to  $A_{63}$  in formula (64)

$$[\sin^4 \frac{1}{2}I \cos (2\xi + 2\nu)]_0 = \sin^4 \frac{1}{2}\omega \cos^4 \frac{1}{2}i = 0.0017 \quad (72)$$

**77.** The ratio obtained by dividing the true obliquity factor for any value of  $I$  by its mean value may be called a *node factor* since it is a function of the longitude of the moon's node. The symbol generally used for the node factor is the small  $f$ . The node factor may be used with a mean constituent coefficient to obtain the true coefficient corresponding to a given longitude of the moon's node. Node factors for the several terms of formulas (62) to (64) may be expressed by the following ratios:

$$f(A_1) \text{ to } f(A_5) = f(Mm) = (2/3 - \sin^2 I) / 0.5021 \quad (73)$$

$$f(A_6) \text{ to } f(A_{13}) = f(Mf) = \sin^2 I / 0.1578 \quad (74)$$

$$f(A_{14}) \text{ to } f(A_{21}) = f(O_1) = \sin I \cos^2 \frac{1}{2}I / 0.3800 \quad (75)$$

$$f(A_{22}) \text{ to } f(A_{30}) = f(J_1) = \sin 2I / 0.7214 \quad (76)$$

$$f(A_{31}) \text{ to } f(A_{38}) = f(OO_1) = \sin I \sin^2 \frac{1}{2}I / 0.0164 \quad (77)$$

$$f(A_{39}) \text{ to } f(A_{46}) = f(M_2) = \cos^4 \frac{1}{2}I / 0.9154 \quad (78)$$

$$f(A_{47}) \text{ to } f(A_{55}) = \sin^2 I / 0.1565 \quad (79)$$

$$f(A_{56}) \text{ to } f(A_{63}) = \sin^4 \frac{1}{2}I / 0.0017 \quad (80)$$

Node factors for the middle of each calendar year from 1850 to 1999 are given in table 14 for the constituents used in the Coast and Geodetic Survey tide-predicting machine. These include all the factors above excepting formulas (79) and (80). However, since formula (79) represents an increase of only about one per cent over formula (74), the tabular values for the latter are readily adapted to formula (79). Node factors change slowly and interpolations can be made in table 14 for any desired part of the year. For practical purposes, however, the values for the middle of the year are generally taken as constant for the entire year.

**78.** The reciprocal of the node factor is called the *reduction factor* and is usually represented by the capital  $F$ . Applied to tidal coefficients pertaining to any particular year, the reduction factors serve to reduce them to a uniform standard in order that they may be comparable. Logarithms of the reduction factors for every tenth of a degree of  $I$  are given in table 12 for the constituents used on the tide-predicting machine of this office.

**79.** Formulas (62), (63), and (64), for the long-period, diurnal, and semidiurnal constituents of the vertical component of the tide-producing force may now be summarized as follows:

Let  $E$  = constituent argument from table 2

$C$  = mean constituent coefficient from table 2

$f$  = node factor from table 14

Then

$$F_{v30}/g = 3/2 U(1/2 - 3/2 \sin^2 Y) \Sigma fC \cos E \quad (81)$$

$$F_{v31}/g = 3/2 U \sin 2Y \Sigma fC \cos E \quad (82)$$

$$F_{v32}/g = 3/2 U \cos^2 Y \Sigma fC \cos E \quad (83)$$

Latitude factors for each degree of  $Y$  are given in table 3. The column symbol in this table is  $Y$  with annexed letter and digits corresponding to those in the designation of the tidal forces. Thus,  $Y_{v30}$  represents the latitude factor to be used with force  $F_{v30}$ , its value being equal to the function  $(1/2 - 3/2 \sin^2 Y)$ . Taking the numerical value for the basic factor  $U$  from table 1, the general coefficient  $3/2 U$  is found to be  $0.8373 \times 10^{-7}$ .

#### HORIZONTAL COMPONENTS OF FORCE

**80.** The horizontal component of the principal part of the tide-producing force as expressed by formula (25), page 14, is in the direction of the azimuth of the tide-producing body. This component may be further resolved into a north-and-south and an east-and-west direction. In the following discussion the south and west will be considered as the positive directions for these components. Now let

$F_{s3}/g$  = south component of principal tide-producing force

$F_{w3}/g$  = west component of principal tide-producing force

$A$  = azimuth of moon reckoned from the south through the west.

From formula (25), we then have

$$F_{s3}/g = 3/2 (M/E)(a/d)^3 \sin 2z \cos A \quad (84)$$

$$F_{w3}/g = 3/2 (M/E)(a/d)^3 \sin 2z \sin A \quad (85)$$

**81.** Referring to figure 3, page 16, the angle  $P'PM$  equals  $A$ , the azimuth of the moon. Now, keeping in mind that the angle  $MPC$  is the supplement of  $A$ , the angle  $PCM$  equals  $t$ , and the arcs  $MC$  and  $PC$  are the respective complements of  $D$  and  $Y$ , we may obtain from the spherical triangle  $MPC$  the following relations:

$$\sin z \cos A = -\cos Y \sin D + \sin Y \cos D \cos t \quad (86)$$

$$\sin z \sin A = \cos D \sin t \quad (87)$$

Multiplying each of the above equations by the value of  $\cos z$  from formula (31), the following equations may be derived:

$$\begin{aligned} \sin 2z \cos A &= 2 \sin z \cos z \cos A \\ &= 3/4 \sin 2Y (2/3 - 2 \sin^2 D) \\ &\quad - \cos 2Y \sin 2D \cos t \\ &\quad + 1/2 \sin 2Y \cos^2 D \cos 2t \end{aligned} \quad (88)$$

$$\begin{aligned} \sin 2z \sin A &= 2 \sin z \cos z \sin A \\ &= \sin Y \sin 2D \sin t \\ &\quad + \cos Y \cos^2 D \sin 2t \end{aligned} \quad (89)$$

**82.** Substituting in (84) and (85) the quantities from equations (88) and (89), we have

$$\begin{aligned} F_{s3}/g &= 9/8 (M/E)(a/d)^3 \sin 2Y (2/3 - 2 \sin^2 D) \dots & F_{s30}/g \\ &\quad - 3/2 (M/E)(a/d)^3 \cos 2Y \sin 2D \cos t \dots & F_{s31}/g \\ &\quad + 3/4 (M/E)(a/d)^3 \sin 2Y \cos^2 D \cos 2t \dots & F_{s32}/g \end{aligned} \quad (90)$$

$$\begin{aligned} F_{w3}/g &= 3/2 (M/E)(a/d)^3 \sin Y \sin 2D \sin t \dots & F_{w31}/g \\ &\quad + 3/2 (M/E)(a/d)^3 \cos Y \cos^2 D \sin 2t \dots & F_{w32}/g \end{aligned} \quad (91)$$

The south component is expressed by three terms representing respectively the long-period, diurnal, and semidiurnal constituents. For the west component there are only two terms—the diurnal and semidiurnal, there being no long-period constituents in the west component. Each term has been marked separately by a symbol with annexed digits analogous to those used for the vertical component to indicate the class to which the term belongs.

**83.** Comparing formula (90) for the south component with formula (32) for the vertical component, it will be noted that the same functions of  $D$  and  $t$  are involved in the corresponding terms of both formulas, and that the terms differ only in their numerical coefficient and the latitude factor. Allowing for these differences, summarized formulas analogous to those given for the vertical component (page 26) may be readily formed. In order to eliminate the negative sign of the coefficient of the middle term,  $180^\circ$  will be applied to the arguments of that term. With all symbols as before, we then have

$$F_{s30}/g = 9/8 U \sin 2Y \Sigma fC \cos E \quad (92)$$

$$F_{s31}/g = 3/2 U \cos 2Y \Sigma fC \cos (E + 180^\circ) \quad (93)$$

$$F_{s32}/g = 3/4 U \sin 2Y \Sigma fC \cos E \quad (94)$$

**84.** Comparing the two terms in formula (91) for the west component with the corresponding terms in formula (32) for the vertical component, it will be noted that the  $D$  functions are the same but that in (91) the sine replaces the cosine for the functions of  $t$ . It may be shown that the corresponding development of these terms will be the same as for the vertical component except that in the developed series each argument will be represented by its sine instead of cosine. In order that the summarized formulas may be expressed in cosine functions,  $90^\circ$  will be subtracted from each argument. With the same symbols as before and allowing for differences in the latitude factors, we obtain

$$F_{w31}/g = 3/2 U \sin Y \Sigma fC \cos (E - 90^\circ) \quad (95)$$

$$F_{w32}/g = 3/2 U \cos Y \Sigma fC \cos (E - 90^\circ) \quad (96)$$

**85.** Formulas for the horizontal component of tide-producing force in any given direction may be derived as follows: Let  $A$  equal the azimuth (measured from south through west) of given direction, and let  $F_{a30}/g$ ,  $F_{a31}/g$ , and  $F_{a32}/g$ , respectively, represent the long-period, diurnal, and semidiurnal terms of the component in this direction. Then

$$F_{a30}/g = F_{s30}/g \times \cos A \quad (97)$$

$$F_{a31}/g = F_{s31}/g \times \cos A + F_{w31}/g \times \sin A \quad (98)$$

$$F_{a32}/g = F_{s32}/g \times \cos A + F_{w32}/g \times \sin A \quad (99)$$

As the long-period term has no west component, the summarized formula for the azimuth  $A$  may be derived by simply introducing the factor  $\cos A$  into the coefficient of formula (92). For the diurnal and semidiurnal terms it is necessary to combine the resolved elements from the south and west components.

**86.** Referring to formulas (93) to (96) and considering a single constituent in each species we obtain the following:

Diurnal constituent,

$$\begin{aligned} & 3/2 UfC [\cos 2Y \cos A \cos (E+180^\circ) + \sin Y \sin A \cos (E-90^\circ)] \\ & = 3/2 UfC (-\cos 2Y \cos A \cos E + \sin Y \sin A \sin E) \\ & = 3/2 UfC P_1 \cos (E-X_1) \end{aligned} \quad (100)$$

in which

$$P_1 = (\cos^2 2Y \cos^2 A + \sin^2 Y \sin^2 A)^{\frac{1}{2}} \quad (101)$$

$$X_1 = \tan^{-1} \frac{\sin Y \sin A}{-\cos 2Y \cos A} \quad (102)$$

Semidiurnal constituent,

$$\begin{aligned} & 3/2 UfC [\sin Y \cos Y \cos A \cos E + \cos Y \sin A \cos (E-90^\circ)] \\ & = 3/2 UfC \cos Y (\sin Y \cos A \cos E + \sin A \sin E) \\ & = 3/2 UfC P_2 \cos (E-X_2) \end{aligned} \quad (103)$$

in which

$$P_2 = \cos Y (\sin^2 Y \cos^2 A + \sin^2 A)^{\frac{1}{2}} \quad (104)$$

$$X_2 = \tan^{-1} \frac{\sin A}{\sin Y \cos A} \quad (105)$$

87. Summarized formulas for the horizontal component of the tide-producing force in any direction  $A$  may now be written as follows:

$$F_{a30}/g = 9/8 U \sin 2Y \cos A \Sigma fC \cos E \quad (106)$$

$$F_{a31}/g = 3/2 UP_1 \Sigma fC \cos (E-X_1) \quad (107)$$

$$F_{a32}/g = 3/2 UP_2 \Sigma fC \cos (E-X_2) \quad (108)$$

the values for  $P_1$ ,  $P_2$ ,  $X_1$  and  $X_2$  being obtained by formulas in the preceding paragraph.  $P_1$  and  $P_2$  are to be taken as positive and the following table will be found convenient in determining the proper quadrant for  $X_1$  and  $X_2$ .

A quadrant	North latitude		South latitude	
	$X_1$ quadrant	$X_2$ quadrant	$X_1$ quadrant	$X_2$ quadrant
1	2 or 1	1	3 or 4	2
2	1 or 2	2	4 or 3	1
3	4 or 3	3	1 or 2	4
4	3 or 4	4	2 or 1	3

For the  $X_1$  quadrant the first value of each pair is applicable when the latitude does not exceed  $45^\circ$  north or south. Otherwise the second value is applicable.

#### EQUILIBRIUM TIDE

88. The *equilibrium theory* of the tides is a hypothesis under which it is assumed that the waters covering the face of the earth instantly respond to the tide-producing forces of the moon and the sun and form a surface of equilibrium under the action of these forces. The theory disregards friction and inertia and the irregular distribution of the land masses of the earth. Although the actual tidal movement




FIGURE 5.




FIGURE 6.

of nature does not even approximate to that which might be expected under the assumed conditions, the theory is of value as an aid in visualizing the distribution of the tidal forces over the surface of the earth. The theoretical tide formed under these conditions is known as the *equilibrium tide*, and sometimes as the *astronomical* or *gravitational* tide.

99. Under the equilibrium theory, the moon would tend to draw the earth into the shape of a prolate spheroid with the longest axis in line with the moon, thus producing one high water directly under the moon and another one on the opposite side of the earth with a low water belt extending entirely around the earth in a great circle midway between the high water points. It may be shown mathematically, however, that the total effect of the moon at its mean distance would be to raise the high water points about 14 inches above the mean surface of the earth and depress the low water belt about 7 inches below this surface, giving a maximum range of tide of about 21 inches. The corresponding range due to the sun is about 10 inches. Figures 5 and 6 illustrate on an exaggerated scale the theoretical disturbing effect of the moon on the earth. In the first figure the moon is assumed to be directly over the equator and in the last figure the moon is approximately at its greatest north declination.

90. With the moon over the equator (fig. 5), the range of the equilibrium tide will be at a maximum at the equator and diminish to zero at the poles and at any point there will be two high and low waters of equal range with each rotation of the earth. With the moon north or south of the equator (fig. 6), a declinational inequality is introduced and the two high and low waters of the day for any given latitude would no longer be equal except at the equator. This inequality would increase with the latitude and near the poles only one high and low water would occur with each rotation of the earth. Although latitude is an important factor in determining the range of the equilibrium tide, it is to be kept in mind that in the actual tide of nature the latitude of a place has no direct effect upon the rise and fall of the water.

91. A surface of equilibrium is a surface at every point of which the sum of the potentials of all the forces is a constant. On such a surface the resultant of all the forces at each point must be in the direction of the normal to the surface at that point. If the earth were a homogeneous mass with gravity as the only force acting, the surface of equilibrium would be that of a sphere. Each additional force will tend to disturb this spherical surface, and the total deformation will be represented by the sum of the disturbances of each of the forces acting separately. In the following investigation we need not be especially concerned with the more or less permanent deformation due to the centrifugal force of the earth's rotation, since we may assume that the disturbances of this spheroidal surface due to the tidal forces will not differ materially from the disturbances in a true spherical surface due to the same cause.

92. The potential at any point due to a force is the amount of work that would be required to move a unit of matter from that point, against the action of the force, to a position where the force is zero. This amount of work will be independent of the path along which the unit of matter is moved. If the force being considered is the gravity of the earth the potential at any point will be the amount

of work required to move a unit mass against the force of gravity from the point to an infinite distance from the earth's center. For the tide-producing force, the potential at any point will be measured by the amount of work necessary to move the unit of mass to the earth's center where this force is zero.

93. Referring to formula (21) for the vertical component of the tide-producing force, if the unit  $g$  is replaced by the unit  $\mu$  from equation (15), the formula may be written as follows:

$$F_v = \frac{3\mu M}{d^3} (\cos^2 z - 1/3)r + \frac{3\mu M}{2d^4} (5 \cos^3 z - 3 \cos z)r^2 \quad (109)$$

94. Considering separately the tide-producing potential due to the two terms in the above formula, let the potential for the first term involving the cube of the moon's distance be represented by  $V_3$  and the potential for the second term involving the 4th power of the moon's distance by  $V_4$ . In each case the work required to move a unit mass against the force through an infinitesimal distance  $-dr$  toward the center of the earth is the product of the force by  $-dr$ , and the potential or total work required to move the particle to the center of the earth may be obtained by integrating between the limits  $r$  and zero. Thus

$$\begin{aligned} V_3 &= -\frac{3\mu M}{d^3} (\cos^2 z - 1/3) \int_r^0 r dr \\ &= \frac{3\mu M}{2d^3} (\cos^2 z - 1/3)r^2 \end{aligned} \quad (110)$$

$$\begin{aligned} V_4 &= -\frac{3\mu M}{2d^4} (5 \cos^3 z - 3 \cos z) \int_r^0 r^2 dr \\ &= \frac{\mu M}{2d^4} (5 \cos^3 z - 3 \cos z)r^3 \end{aligned} \quad (111)$$

95. At any instant of time the tide-producing potential at different points on the earth's surface will depend upon the zenith distance ( $z$ ) of the moon and may be either positive or negative. It will now be shown that the average tide-producing potential for all points on the earth's surface, assuming it to be a sphere, is zero. Assume a series of right conical surfaces with common apex at center of earth and axis coinciding with the line joining centers of earth and moon, the angle between the generating line and the axis being  $z$ . These conical surfaces separated by infinitesimal angle  $dz$  will cut the surface of the sphere into a series of equipotential rings, the surface area of any ring being equal to a  $2\pi r^2 \sin z dz$ . The average potential for the entire spherical surface may then be obtained by summing the products of the ring areas and corresponding potentials and dividing the sum by the total surface area of the sphere. Thus

$$\begin{aligned} \text{Average } V_3 &= \frac{3\mu Mr^2}{4d^3} \int_0^\pi (\cos^2 z - 1/3) \sin z dz \\ &= \frac{3\mu Mr^2}{4d^3} \left[ -1/3 \cos^3 z + 1/3 \cos z \right]_0^\pi = 0 \end{aligned} \quad (112)$$

$$\text{Average } V_4 = \frac{\mu M r^3}{4d^4} \int_0^\pi (5 \cos^3 z - 3 \cos z) \sin z \, dz \\ = \frac{\mu M r^3}{4d^4} \left[ -5/4 \cos^4 z + 3/2 \cos^2 z \right]_0^\pi = 0 \quad (113)$$

**96.** Let  $V_o$  represent the potential due to gravity at any point on the earth's surface. Since the force of gravity at any point on or above the earth's surface equals  $\mu E/r^2$ , the corresponding potential becomes

$$V_o = \mu E \int_r^\infty \frac{dr}{r^2} = \frac{\mu E}{r} \quad (114)$$

If the earth is assumed to be a sphere with radius  $a$ , the gravitational potential at each point will equal  $\mu E/a$ , which may be taken as the average gravitational potential over the surface of the earth.

**97.** For a surface of equilibrium under the combined action of gravity and that part of the tide-producing force involving the cube of the moon's distance the sum of the corresponding potentials must be a constant, and since the average tide-producing potential for the entire surface of the earth is zero (par. 95), the constant will be the average gravitational potential or  $\mu E/a$ . Then from (110) and (114) we have

$$V_3 + V_o = \frac{3\mu M}{2d^3} (\cos^2 z - 1/3)r^2 + \frac{\mu E}{r} = \frac{\mu E}{a} \quad (115)$$

Transposing and omitting common factor  $\mu$ , we may obtain

$$\frac{(r-a)a^2}{r^3} = 3/2(M/E)(a/d)^3(\cos^2 z - 1/3) \quad (116)$$

Let

$$r = a + h \quad (117)$$

so that  $h$  represents the height of the equilibrium surface as referred to the undisturbed spherical surface of an equivalent sphere. Then

$$\frac{(r-a)a^2}{r^3} = \frac{ha^2}{(a+h)^3} = h/a - 3(h/a)^2 + 6(h/a)^3 - \text{etc.} \quad (118)$$

As fraction  $h/a$  is very small, its greatest value being less than 0.000001, the powers above the first may be neglected. Substituting in (116) and writing  $h$  with subscript  $_3$  to identify it with the principal tide-producing force, we have

$$h_3/a = 3/2(M/E)(a/d)^3(\cos^2 z - 1/3) \quad (119)$$

**98.** Similarly, for a surface of equilibrium under the combined action of gravity and the part of the tide-producing force involving the 4th power of the moon's distance, we have from (111) and (114)

$$V_4 + V_o = \frac{\mu M}{2d^4} (5 \cos^3 z - 3 \cos z)r^3 + \frac{\mu E}{r} = \frac{\mu E}{a} \quad (120)$$

$$\frac{(r-a)a^3}{r^4} = 1/2 (M/E)(a/d)^4(5 \cos^3 z - 3 \cos z) \quad (121)$$

Letting  $r=a+h_4$  and expanding the first member of the above formula, it becomes equal to  $h_4/a$  after the rejection of the higher powers of this small fraction. The formula may then be written

$$h_4/a = 1/2 (M/E)(a/d)^4(5 \cos^3 z - 3 \cos z) \quad (122)$$

**99.** Formulas (119) and (122) involving the cube and 4th power of the moon's parallax, respectively, represent the equilibrium heights of the tide due to the corresponding forces, the heights being expressed in respect to the mean radius ( $a$ ) of the earth as the unit. In deriving these formulas the centrifugal force of the earth's rotation was disregarded and the resulting heights represent the disturbances in a true spherical surface due to the action of the tide-producing force. It may be inferred that in a condition of equilibrium the tidal forces would produce like disturbances in the spheroidal surface of the earth and the  $h$  of the formulas may therefore be taken as being referred to the earth's surface as defined by the mean level of the sea.

**100.** The extreme limits of the equilibrium tide, applicable to the time when the tide-producing body is nearest the earth, may be obtained by substituting the proper numerical values in formulas (119) and (122). They are given below for both moon and sun.

From formula (119) involving the cube of parallax—

$$\text{Greatest rise} = 1.46 \text{ feet for moon, or } 0.57 \text{ foot for sun} \quad (123)$$

$$\text{Lowest fall} = 0.73 \text{ foot for moon, or } 0.28 \text{ foot for sun} \quad (124)$$

$$\text{Extreme range} = 2.19 \text{ feet for moon, or } 0.85 \text{ foot for sun.} \quad (125)$$

From formula (122) involving the 4th power of parallax—

$$\text{Greatest rise} = 0.026 \text{ foot for moon, or } 0.000025 \text{ foot for sun} \quad (126)$$

$$\text{Lowest fall} = 0.026 \text{ foot for moon, or } 0.000025 \text{ foot for sun} \quad (127)$$

$$\text{Extreme range} = 0.052 \text{ foot for moon, or } 0.00005 \text{ foot for sun.} \quad (128)$$

**101.** A comparison of formulas (23) and (119), the first expressing the relation of the vertical component of the principal tide-producing force to the acceleration of gravity ( $g$ ) and the other the relation of the height of the corresponding equilibrium tide to the mean radius ( $a$ ) of the earth, will show that they are identical with the single exception that the coefficient of the height formula is one-half that of the force formula. Therefore the development of the force formula into a series of harmonic constituents is immediately applicable in obtaining similar expressions for the equilibrium height of the tide. Using a notation for the height terms corresponding to that used for the force terms, let  $h_{30}/a$ ,  $h_{31}/a$ , and  $h_{32}/a$  represent, respectively, the long-period, diurnal, and semidiurnal terms of the equilibrium tide involving the cube of the moon's parallax. Then referring to formulas (81) to (83) we may write

$$h_{30}/a = 3/4 U(1/2 - 3/2 \sin^2 Y) \sum fC \cos E \quad (129)$$

$$h_{31}/a = 3/4 U \sin 2Y \sum fC \cos E \quad (130)$$

$$h_{32}/a = 3/4 U \cos^2 Y \sum fC \cos E \quad (131)$$

the symbols having the same significance as in the preceding discussion of the tidal forces.

## TERMS INVOLVING 4TH POWER OF MOON'S PARALLAX

**102.** Formulas (24) and (26) represent the vertical and horizontal components of the part of the tide-producing force involving the 4th power of the moon's parallax. This part of the force constitutes only about 2 percent of the total tide-producing force of the moon and for brevity will be called the *lesser* force to distinguish it from the principal or primary part involving the cube of the parallax. The vertical component  $F_{v4}/g$  has its maximum value when  $z$  equals zero and, if numerical values pertaining to the moon and sun when nearest the earth are substituted in formula (24), the extreme values for this component are found to be  $0.37 \times 10^{-8}$  for the moon and  $0.35 \times 10^{-11}$  for the sun. The horizontal component  $F_{a4}/g$  has its greatest value when  $z$  equals about  $31.09^\circ$  and the substitution of numerical values in formula (26) gives the extreme value of this component as  $0.26 \times 10^{-8}$  for the moon or  $0.24 \times 10^{-11}$  for the sun.

**103.** Substituting in (24) the value of  $\cos z$  from (31), the vertical component of the lesser force is expanded into four terms as follows:

$$\begin{aligned} F_{v4}/g = & 15/4 (M/E)(a/d)^4 \sin Y (\cos^2 Y - 2/5) \sin D (5 \cos^2 D - 2) \quad F_{v40}/g \\ & + 45/8 (M/E)(a/d)^4 \cos Y (\cos^2 Y - 4/5) \cos D (5 \cos^2 D - 4) \cos t \quad F_{v41}/g \\ & + 45/4 (M/E)(a/d)^4 \sin Y \cos^2 Y \sin D \cos^2 D \cos 2t \quad F_{v42}/g \\ & + 15/8 (M/E)(a/d)^4 \cos^3 Y \cos^3 D \cos 3t \quad F_{v43}/g \end{aligned} \quad (132)$$

These four terms represent, respectively, long-period, diurnal, semi-diurnal, and terdiurnal constituents, according to the multiple of the hour angle  $t$  involved in the term. Each term is followed by a symbol which is analogous to those used in the development of the principal force.

**104.** Each term in formula (132) may be further expanded by means of the relations given in formulas (39) and (42). Expressing these terms separately we have—

$$\begin{aligned} F_{v40}/g = & 15/4 (M/E)(a/d)^4 \sin Y (\cos^2 Y - 2/5) \times \\ & [3(\sin I - 5/4 \sin^3 I) \cos (j - 90^\circ) \\ & + 5/4 \sin^3 I \cos (3j - 90^\circ)] \end{aligned} \quad (133)$$

$$\begin{aligned} F_{v41}/g = & 45/8 (M/E)(a/d)^4 \cos Y (\cos^2 Y - 4/5) \times \\ & [5/4 \sin^2 I \cos^2 \frac{1}{2}I \cos (X - 3j) \\ & + (1 - 10 \sin^2 \frac{1}{2}I + 15 \sin^4 \frac{1}{2}I) \cos^2 \frac{1}{2}I \cos (X - j) \\ & + (1 - 10 \cos^2 \frac{1}{2}I + 15 \cos^4 \frac{1}{2}I) \sin^2 \frac{1}{2}I \cos (X + j) \\ & + 5/4 \sin^2 I \sin^2 \frac{1}{2}I \cos (X + 3j)] \end{aligned} \quad (134)$$

$$\begin{aligned} F_{v42}/g = & 45/8 (M/E)(a/d)^4 \sin Y \cos^2 Y \times \\ & [\sin I \cos^4 \frac{1}{2}I \cos (2X - 3j + 90^\circ) \\ & + 3(\cos^2 \frac{1}{2}I - 2/3) \sin I \cos^2 \frac{1}{2}I \cos (2X - j - 90^\circ) \\ & + 3(\cos^2 \frac{1}{2}I - 1/3) \sin I \sin^2 \frac{1}{2}I \cos (2X + j - 90^\circ) \\ & + \sin I \sin^4 \frac{1}{2}I \cos (2X + 3j - 90^\circ)] \end{aligned} \quad (135)$$

$$\begin{aligned} F_{v43}/g = & 15/8 (M/E)(a/d)^4 \cos^3 Y \times \\ & [\cos^6 \frac{1}{2}I \cos (3X - 3j) \\ & + 3 \cos^4 \frac{1}{2}I \sin^2 \frac{1}{2}I \cos (3X - j) \\ & + 3 \cos^2 \frac{1}{2}I \sin^4 \frac{1}{2}I \cos (3X + j) \\ & + \sin^6 \frac{1}{2}I \cos (3X + 3j)] \end{aligned} \quad (136)$$

**105.** If the common factor  $(a/d)^4$  in formulas (133) to (136) is replaced by its equivalent  $(a/c)^4 \times (c/d)^4$ , these formulas may be de-

veloped into numerous constituent terms by a method similar to that already described in the development of the principal lunar force (paragraphs 59-69). In the following development constituents of very small magnitude are omitted. Those given are numbered consecutively with the constituent terms of the principal lunar force.

$$\begin{aligned} F_{v40}/g = & 15/4 (M/E)(a/c)^4 \sin Y (\cos^2 Y - 2/5) \times \\ (A_{64}) \quad & [(\sin I - 5/4 \sin^3 I) \{ 3(1+2e^2) \cos(s-90^\circ-\xi) \\ (A_{65}) \quad & + 9e \cos(2s-p-90^\circ-\xi) \\ (A_{66}) \quad & + 3e \cos(p-90^\circ-\xi) \} \\ (A_{67}) \quad & + \sin^3 I \{ 5/4(1-6e^2) \cos(3s-90^\circ-3\xi) \\ (A_{68}) \quad & + 25/4 e \cos(4s-p-90^\circ-3\xi) \}] \end{aligned} \quad (137)$$

$$\begin{aligned} F_{v41}/g = & 45/8 (M/E)(a/c)^4 \cos Y (\cos^2 Y - 4/5) \times \\ (A_{69}) \quad & [\sin^2 I \cos^2 \frac{1}{2}I \{ 5/4(1-6e^2) \cos(T-3s+h+3\xi-\nu) \\ (A_{70}) \quad & + 25/4 e \cos(T-4s+h+p+3\xi-\nu) \} \\ (A_{71}) \quad & + (1-10 \sin^2 \frac{1}{2}I + 15 \sin^4 \frac{1}{2}I) \cos^2 \frac{1}{2}I \\ (A_{72}) \quad & \{ (1+2e^2) \cos(T-s+h+\xi-\nu) \dots [M_1] \\ (A_{73}) \quad & + 3e \cos(T-2s+h+p+\xi-\nu) \\ (A_{74}) \quad & + (1-10 \cos^2 \frac{1}{2}I + 15 \cos^4 \frac{1}{2}I) \sin^2 \frac{1}{2}I \\ (A_{75}) \quad & \{ (1+2e^2) \cos(T+s+h-\xi-\nu) \\ & + 3e \cos(T+2s+h-p-\xi-\nu) \}] \end{aligned} \quad (138)$$

$$\begin{aligned} F_{v42}/g = & 45/8 (M/E)(a/c)^4 \sin Y \cos^2 Y \times \\ (A_{76}) \quad & [\sin I \cos^4 \frac{1}{2}I \{ (1-6e^2) \cos(2T-3s+2h+90^\circ+3\xi-2\nu) \\ (A_{77}) \quad & + 5e \cos(2T-4s+2h+p+90^\circ+3\xi-2\nu) \\ (A_{78}) \quad & + e \cos(2T-2s+2h-p-90^\circ+3\xi-2\nu) \} \\ (A_{79}) \quad & + (\cos^2 \frac{1}{2}I - 2/3) \sin I \cos^2 \frac{1}{2}I \\ (A_{80}) \quad & \{ 3(1+2e^2) \cos(2T-s+2h-90^\circ+\xi-2\nu) \\ (A_{81}) \quad & + 9e \cos(2T-2s+2h+p-90^\circ+\xi-2\nu) \} \\ (A_{82}) \quad & + (\cos^2 \frac{1}{2}I - 1/3) \sin I \sin^2 \frac{1}{2}I \\ (A_{83}) \quad & \{ 3(1+2e^2) \cos(2T+s+2h-90^\circ-\xi-2\nu) \}] \end{aligned} \quad (139)$$

$$\begin{aligned} F_{v43}/g = & 15/8 (M/E)(a/c)^4 \cos^3 Y \times \\ (A_{82}) \quad & [\cos^6 \frac{1}{2}I \{ (1-6e^2) \cos(3T-3s+3h+3\xi-3\nu) \dots M_3 \\ (A_{83}) \quad & + 5e \cos(3T-4s+3h+p+3\xi-3\nu) \\ (A_{84}) \quad & + e \cos(3T-2s+3h-p+180^\circ+3\xi-3\nu) \\ (A_{85}) \quad & + 127/8 e^2 \cos(3T-5s+3h+2p+3\xi-3\nu) \\ (A_{86}) \quad & + 75/8 me \cos(3T-4s+5h-p+3\xi-3\nu) \} \\ (A_{87}) \quad & + \cos^4 \frac{1}{2}I \sin^2 \frac{1}{2}I \\ (A_{88}) \quad & \{ 3(1+2e^2) \cos(3T-s+3h+\xi-3\nu) \\ & + 9e \cos(3T-2s+3h+p+\xi-3\nu) \}] \end{aligned} \quad (140)$$

**106.** All of the constituent terms in formulas (137) to (140) are relatively unimportant but they are listed in table 1 because of their theoretical interest. The only one of these terms now used in the prediction of tides is (A<sub>82</sub>) representing the constituent M<sub>3</sub> which has a speed exactly three-halves that of the principal lunar constituent M<sub>2</sub>. Term (A<sub>71</sub>) is of interest in having a speed exactly one-half that of M<sub>2</sub> and is sometimes called the true M<sub>1</sub> to distinguish it from the composite M<sub>1</sub> which is used in the prediction of tides and which will be described later.

**107.** For simplicity and the purposes of this publication, the mean values of the obliquity factors in the terms of the lesser tide-producing force will be taken as the values pertaining to the time when  $I$  equals  $\omega$  or  $23.452^\circ$ , excepting that for constituent  $M_3$  and associated terms the mean has been obtained in accord with the system described in paragraph 75. The corresponding node factors (paragraph 77) may then be expressed by the following formulas in which the denominators are the accepted means of the obliquity factors:

$$f(A_{64}) \text{ to } f(A_{66}) = (\sin I - 5/4 \sin^3 I) / 0.3192 \quad (141)$$

$$f(A_{67}) \text{ to } f(A_{68}) = \sin^3 I / 0.0630 \quad (142)$$

$$f(A_{69}) \text{ to } f(A_{70}) = \sin^2 I \cos^{2\frac{1}{2}} I / 0.1518 \quad (143)$$

$$f(A_{71}) \text{ to } f(A_{73}) = (1 - 10 \sin^{2\frac{1}{2}} I + 15 \sin^{4\frac{1}{2}} I) \cos^{2\frac{1}{2}} I / 0.5873 \quad (144)$$

$$f(A_{74}) \text{ to } f(A_{75}) = (1 - 10 \cos^{2\frac{1}{2}} I + 15 \cos^{4\frac{1}{2}} I) \sin^{2\frac{1}{2}} I / 0.2147 \quad (145)$$

$$f(A_{76}) \text{ to } f(A_{78}) = \sin I \cos^{4\frac{1}{2}} I / 0.3658 \quad (146)$$

$$f(A_{79}) \text{ to } f(A_{80}) = (\cos^{2\frac{1}{2}} I - 2/3) \sin I \cos^{2\frac{1}{2}} I / 0.1114 \quad (147)$$

$$f(A_{81}) = (\cos^{2\frac{1}{2}} I - 1/3) \sin I \sin^{2\frac{1}{2}} I / 0.0103 \quad (148)$$

$$f(A_{82}) \text{ to } f(A_{86}) = f(M_3) = \cos^{6\frac{1}{2}} I / 0.8758 \quad (149)$$

$$f(A_{87}) \text{ to } f(A_{88}) = \cos^{4\frac{1}{2}} I \sin^{2\frac{1}{2}} I / 0.0380 \quad (150)$$

Comparing formulas (149) and (78), it will be noted that the node factor for  $M_3$  is equal to the node factor for  $M_2$  raised to the  $3/2$  power. Computed values applicable to terms  $A_{82}$  to  $A_{86}$  are included in table 14 for years 1850 to 1999, inclusive.

**108.** For the tabulated constituent coefficients of the terms in formulas (137) to (140) there are included not only the elliptic and mean obliquity factors but also such other factors as may be necessary to permit the use of the general coefficient ( $3/2 U$ ) of formulas (81) to (83) for the vertical component of the principal tide-producing force. The common coefficient ( $M/E$ ) ( $a/c$ ) $^4$  of formulas (137) to (140) is equal to  $U$  multiplied by the parallax  $a/c$ , and the latter together with the necessary numerical factors is included in the constituent coefficients in table 2. Formulas (137) to (140) may then be summarized as follows:

$$F_{v40} / g = 3/2 U \sin Y (\cos^2 Y - 2/5) \Sigma fC \cos E \quad (151)$$

$$F_{v41} / g = 3/2 U \cos Y (\cos^2 Y - 4/5) \Sigma fC \cos E \quad (152)$$

$$F_{v42} / g = 3/2 U \sin Y \cos^2 Y \Sigma fC \cos E \quad (153)$$

$$F_{v43} / g = 3/2 U \cos^3 Y \Sigma fC \cos E \quad (154)$$

**109.** It is to be noted that in formulas (151), (152), and (153), the maximum value of the latitude factor in each is less than unity, being

0.4, 0.2754, and 0.3849, respectively, if the sign of the function is disregarded. In formula (154), as in the corresponding formulas for the principal tide-producing force, the maximum value of this factor is unity. In comparing the relative importance of the various constituents of the tide-producing force the latitude factor should be included with the mean coefficient. Attention is also called to the fact that the relative importance of the constituents involving the 4th power of the moon's parallax is greater in respect to the vertical component of the tide-producing force than in respect to the height of the equilibrium tide. In table 2 the mean coefficients are taken comparable in respect to the vertical component of the tide-producing force and the constituent coefficients pertaining to the lesser force are therefore 50 percent greater than they would be if taken comparable in respect to the equilibrium tide.

**110.** The south and west horizontal components of the lesser tide-producing force may be obtained by multiplying formula (26) by  $\cos A$  and  $\sin A$ , respectively. Using the same system of notation as before, we then have

$$F_{s4}/g = 3/2 (M/E)(a/d)^4 \sin z (5 \cos^2 z - 1) \cos A \quad (155)$$

$$F_{w4}/g = 3/2 (M/E)(a/d)^4 \sin z (5 \cos^2 z - 1) \sin A \quad (156)$$

**111.** By means of the relations expressed in formulas (31), (86), and (87), the above component forces may be separated into long-period, diurnal, semidiurnal, and terdiurnal terms as follows:

South component,

$$F_{s40}/g = -15/4 (M/E)(a/d)^4 \cos Y (\cos^2 Y - 4/5) \sin D (5 \cos^2 D - 2) \quad (157)$$

$$F_{s41}/g = 45/8 (M/E)(a/d)^4 \sin Y (\cos^2 Y - 4/15) \cos D (5 \cos^2 D - 4) \cos t \quad (158)$$

$$F_{s42}/g = -45/4 (M/E)(a/d)^4 \cos Y (\cos^2 Y - 2/3) \sin D \cos^2 D \cos 2t \quad (159)$$

$$F_{s43}/g = 15/8 (M/E)(a/d)^4 \sin Y \cos^2 Y \cos^3 D \cos 3t \quad (160)$$

West component,

$$F_{w41}/g = 15/8 (M/E)(a/d)^4 (\cos^2 Y - 4/5) \cos D (5 \cos^2 D - 4) \sin t \quad (161)$$

$$F_{w42}/g = 15/4 (M/E)(a/d)^4 \sin 2Y \sin D \cos^2 D \sin 2t \quad (162)$$

$$F_{w43}/g = 15/8 (M/E)(a/d)^4 \cos^2 Y \cos^3 D \sin 3t \quad (163)$$

**112.** Comparing formulas (157) to (160) for the south component force with the corresponding terms of (132) for the vertical component, it will be noted that they differ only in the latitude factors and in sign for two of the terms. With adjustments for these differences the summarized formulas (151) to (154) are directly applicable for expressing the corresponding terms in the south component. Thus

$$F_{s40}/g = 3/2 U \cos Y (\cos^2 Y - 4/5) \Sigma fC \cos(E + 180^\circ) \quad (164)$$

$$F_{s41}/g = 3/2 U \sin Y (\cos^2 Y - 4/15) \Sigma fC \cos E \quad (165)$$

$$F_{s42}/g = 3/2 U \cos Y (\cos^2 Y - 2/3) \Sigma fC \cos(E + 180^\circ) \quad (166)$$

$$F_{s43}/g = 3/2 U \sin Y \cos^2 Y \Sigma fC \cos E \quad (167)$$

**113.** For the west component there is no long-period term. Comparing (161) to (163) with the corresponding terms of (132), it will be noted that the  $t$ -functions are expressed as sines instead of cosines but they may be changed to the latter by subtracting  $90^\circ$  from each

argument. With this change and allowing for differences in the latitude factors and numerical coefficients, the summarized formulas for the west component will be similar to those for the vertical component and may be written as follows:

$$F_{w41}/g = 1/2 U (\cos^2 Y - 4/5) \Sigma fC \cos (E - 90^\circ) \quad (168)$$

$$F_{w42}/g = 1/2 U \sin 2Y \Sigma fC \cos (E - 90^\circ) \quad (169)$$

$$F_{w43}/g = 3/2 U \cos^2 Y \Sigma fC \cos (E - 90^\circ) \quad (170)$$

**114.** To obtain the horizontal component of the lesser force in any direction, the same procedure may be followed as was used for the principal tide-producing force (paragraphs 85 to 87). With the same system of notation we then have

$$F_{a40}/g = 3/2 U \cos Y (\cos^2 Y - 4/5) \cos A \Sigma fC \cos (E + 180^\circ) \quad (171)$$

$$F_{a41}/g = 3/2 U P_1 \Sigma fC \cos (E - X_1) \quad (172)$$

$$F_{a42}/g = 3/2 U P_2 \Sigma fC \cos (E - X_2) \quad (173)$$

$$F_{a43}/g = 3/2 U P_3 \Sigma fC \cos (E - X_3) \quad (174)$$

in which

$$P_1 = [\sin^2 Y (\cos^2 Y - 4/15)^2 \cos^2 A + 1/9 (\cos^2 Y - 4/5)^2 \sin^2 A]^{1/2} \quad (175)$$

$$P_2 = \cos Y [(\cos^2 Y - 2/3)^2 \cos^2 A + 4/9 \sin^2 Y \sin^2 A]^{1/2} \quad (176)$$

$$P_3 = \cos^2 Y (\sin^2 Y \cos^2 A + \sin^2 A)^{1/2} \quad (177)$$

$$X_1 = \tan^{-1} \frac{(\cos^2 Y - 4/5) \sin A}{3 \sin Y (\cos^2 Y - 4/15) \cos A} \quad (178)$$

$$X_2 = \tan^{-1} \frac{2 \sin Y \sin A}{-3 (\cos^2 Y - 2/3) \cos A} \quad (179)$$

$$X_3 = \tan^{-1} \frac{\sin A}{\sin Y \cos A} \quad (180)$$

The proper quadrants for  $X_1$ ,  $X_2$ , and  $X_3$  will be determined by the signs of the numerators and denominators in the above expressions, these signs being respectively the same as for the sine and cosine of the corresponding angles.

**115.** Comparing formula (122) for the equilibrium height of the tide due to the lesser tide-producing force with formula (24) for the vertical component of the force, it will be noted that they are the same with the exception that the numerical coefficient of the former is one-third that of the latter. With this change, the summarized formulas (151) to (154) for the vertical force may be used to express the corresponding equilibrium heights. Following the same system of notation as before, we have

$$h_{40}/a = 1/2 U \sin Y (\cos^2 Y - 2/5) \Sigma fC \cos E \quad (181)$$

$$h_{41}/a = 1/2 U \cos Y (\cos^2 Y - 4/5) \Sigma fC \cos E \quad (182)$$

$$h_{42}/a = 1/2 U \sin Y \cos^2 Y \Sigma fC \cos E \quad (183)$$

$$h_{43}/a = 1/2 U \cos^3 Y \Sigma fC \cos E \quad (184)$$

It is to be noted that the equilibrium height of the tide due to the principal tide-producing force when measured by the mean radius of the earth as a unit is one-half as great as the corresponding vertical component force referred to the mean acceleration of gravity as a unit, while the equilibrium height due to the lesser tide producing force similarly expressed is only one-third as great as the corresponding force. In table 2, the coefficients ( $C$ ) of the constituents derived

from the lesser force are made comparable with the others in respect to the vertical component force rather than in respect to the equilibrium height.

### SOLAR TIDES

**116.** Since the tide-producing force of the sun is similar in action to that of the moon, the formulas derived for the latter are applicable, with suitable substitutions, to the solar forces. Referring to formulas (62), (63), and (64), let  $U$  be replaced by  $U_1$  representing the product  $(S/E)(a/c_1)^3$  in which  $S$  is the mass of the sun and  $(a/c_1)$  its mean parallax. Also replace  $e$  by  $e_1$ , the eccentricity of the earth's orbit;  $I$  by  $\omega$ , the obliquity of the ecliptic;  $s$  by  $h$ , the mean longitude of the sun; and  $p$  by  $p_1$ , the longitude of the solar perigee. For the solar forces the arcs  $\xi$  and  $\nu$  become zero and all terms representing the evectional and variational inequalities are omitted.

**117.** Making the changes indicated the solar constituents are now expressed in the following formulas. Each term is marked for identification by the letter  $B$  with the same subscript used for the corresponding term in the lunar tide. The usual constituent symbol is also given for the more important terms. Using the same system of notation as before,

$$\begin{aligned} \text{Solar } F_{v30}/g = & 3/2 U_1 (1/2 - 3/2 \sin^2 Y) \times \\ (B_1) \quad & [(2/3 - \sin^2 \omega) \{ (1 + 3/2 e_1^2) \dots \dots \text{permanent term} \\ (B_2) \quad & + 3 e_1 \cos(h - p_1) \\ (B_3) \quad & + 9/2 e_1^2 \cos(2h - 2p_1) \} \\ (B_6) \quad & + \sin^2 \omega \{ (1 - 5/2 e_1^2) \cos 2h \dots \dots \text{Ssa} \\ (B_7) \quad & + 7/2 e_1 \cos(3h - p_1) \\ (B_8) \quad & + 1/2 e_1 \cos(h + p_1 + 180^\circ) \\ (B_9) \quad & + 17/2 e_1^2 \cos(4h - 2p_1) \}] \end{aligned} \quad (185)$$

$$\begin{aligned} \text{Solar } F_{v31}/g = & 3/2 U_1 \sin 2Y \times \\ (B_{14}) \quad & [\sin \omega \cos^2 \frac{1}{2}\omega \{ (1 - 5/2 e_1^2) \cos(T - h + 90^\circ) \dots \dots \text{P}_1 \\ (B_{15}) \quad & + 7/2 e_1 \cos(T - 2h + p_1 + 90^\circ) \dots \dots \pi_1 \\ (B_{16}) \quad & + 1/2 e_1 \cos(T - p_1 - 90^\circ) \\ (B_{17}) \quad & + 17/2 e_1^2 \cos(T - 3h + 2p_1 + 90^\circ) \} \\ (B_{22}) \quad & + \sin 2\omega \{ (1/2 + 3/4 e_1^2) \cos(T + h - 90^\circ) \dots \dots [\text{K}_1] \\ (B_{23}) \quad & + 3/4 e_1 \cos(T + p_1 - 90^\circ) \\ (B_{24}) \quad & + 3/4 e_1 \cos(T + 2h - p_1 - 90^\circ) \dots \dots \psi_1 \\ (B_{25}) \quad & + 9/8 e_1^2 \cos(T - h + 2p_1 - 90^\circ) \\ (B_{26}) \quad & + 9/8 e_1^2 \cos(T + 3h - 2p_1 - 90^\circ) \} \\ (B_{31}) \quad & + \sin \omega \sin^2 \frac{1}{2}\omega \{ (1 - 5/2 e_1^2) \cos(T + 3h - 90^\circ) \dots \dots \phi_1 \\ (B_{32}) \quad & + 7/2 e_1 \cos(T + 4h - p_1 - 90^\circ) \\ (B_{33}) \quad & + 1/2 e_1 \cos(T + 2h + p_1 + 90^\circ) \\ (B_{34}) \quad & + 17/2 e_1^2 \cos(T + 5h - 2p_1 - 90^\circ) \}] \end{aligned} \quad (186)$$

$$\begin{aligned} \text{Solar } F_{v32}/g = & 3/2 U_1 \cos^2 Y \times \\ (B_{39}) \quad & [\cos^4 \frac{1}{2}\omega \{ (1 - 5/2 e_1^2) \cos(2T) \dots \dots \text{S}_2 \\ (B_{40}) \quad & + 7/2 e_1 \cos(2T - h + p_1) \dots \dots \text{T}_2 \\ (B_{41}) \quad & + 1/2 e_1 \cos(2T + h - p_1 + 180^\circ) \dots \dots \text{R}_2 \\ (B_{42}) \quad & + 17/2 e_1^2 \cos(2T - 2h + 2p_1) \} \\ (B_{47}) \quad & + \sin^2 \omega \{ (1/2 + 3/4 e_1^2) \cos(2T + 2h) \dots \dots [\text{K}_2] \\ (B_{48}) \quad & + 3/4 e_1 \cos(2T + h + p_1) \\ (B_{49}) \quad & + 3/4 e_1 \cos(2T + 3h - p_1) \} \end{aligned}$$

(Formula continued on next page)

$$\begin{aligned}
 (B_{50}) & +9/8 e_1^2 \cos (2T+2p_1) \\
 (B_{51}) & +9/8 e_1^2 \cos (2T+4h-2p_1) \} \\
 (B_{56}) & +\sin^4 \frac{1}{2}\omega \{ (1-5/2 e_1^2) \cos (2T+4h) \\
 (B_{57}) & +7/2 e_1 \cos (2T+5h-p_1) \\
 (B_{58}) & +1/2 e_1 \cos (2T+3h+p_1+180^\circ) \\
 (B_{59}) & +17/2 e_1^2 \cos (2T+6h-2p_1) \} \\
 \end{aligned} \tag{187}$$

**118.** The general coefficient for the solar tide-producing force differs from that of the lunar force in the basic factor. From the fundamental data in table 1, the ratio of  $U_1/U$  is found to be 0.4602. This ratio, which will be designated as the *solar factor* with symbol  $S'$ , represents the theoretical relation between the principal solar and lunar tide-producing forces. In computing the constituent coefficients of the solar terms for use in table 2, the solar factor was included in order that the same general coefficient may be applicable to both lunar and solar terms. All of the summarized formulas involving the coefficients and arguments of table 2 are therefore applicable to both lunar and solar constituents. For the solar constituents, however, the node factor ( $f$ ) is always unity since  $\omega$ , the obliquity of the ecliptic, may be considered as a constant.

**119.** By substituting solar elements in formulas (137) to (140) the corresponding solar constituents pertaining to the 4th power of the sun's parallax are readily obtained. Since the theoretical magnitude of the lesser solar tide-producing force is less than 0.00002 part of the total tide-producing force of moon and sun, it is usually disregarded altogether. However, certain interest is attached to three of the constituents which are considered in connection with shallow water and meteorological tides (p. 46). These are constituents  $S_a$ ,  $S_1$ , and  $S_3$ , corresponding respectively to terms  $A_{64}$ ,  $A_{71}$ , and  $A_{82}$  of the lunar series. They are listed in table 2 with reference letter  $B$  and corresponding subscripts.  $S_a$  has a speed one-half that of constituent  $Ssa$  represented by term  $B_6$  of formula (185). Its theoretical argument as derived from term  $A_{64}$  contains the constant  $90^\circ$ , but being considered as a meteorological rather than an astronomical constituent, this constant is omitted from the argument. Constituents  $S_1$  and  $S_3$  have speeds respectively one-half and three-halves that of the principal solar constituent  $S_2$ .

**120.** The arguments of a number of the solar constituents include the element  $p_1$  which represents the longitude of the solar perigee. As this changes less than  $2^\circ$  in a century, it may be considered as practically constant for the entire century. Referring to table 4 it will be noted that  $p_1$  changes from  $281.22^\circ$  in 1900 to  $282.94^\circ$  in 2000. The value of  $282^\circ$  may therefore be adopted without material error for all work relating to the present century. With  $p_1$  taken as a constant, it will be found that a number of terms in table 2 have the same speeds and may therefore be expected to merge into single constituents. Thus, constituents receiving contributions from more than one term are as follows:  $S_a$  from terms  $B_2$ ,  $B_8$ , and  $B_{64}$ ;  $Ssa$  from terms  $B_3$  and  $B_6$ ;  $P_1$  from terms  $B_{14}$  and  $B_{25}$ ;  $S_1$  from terms  $B_{16}$ ,  $B_{23}$ , and  $B_{71}$ ;  $\psi_1$  from terms  $B_{24}$  and  $B_{33}$ ;  $\phi_1$  from terms  $B_{26}$  and  $B_{31}$ ;  $S_2$  from terms  $B_{39}$  and  $B_{50}$ ; and  $R_2$  from terms  $B_{41}$  and  $B_{48}$ . A few other solar terms also merge.

THE M<sub>1</sub> TIDE

**121.** The separation of constituents from each other by the process of the analysis depends upon the differences in their speeds. Constituents with nearly equal speeds are not readily separated unless the analysis covers a very long series of observations but they tend to merge and form a single composite constituent. In formula (63), terms  $A_{16}$  and  $A_{23}$  have nearly equal speeds, one being a little less and the other a little greater than one-half the speed of the principal lunar constituent  $M_2$ . These two terms are usually considered as a single constituent and represented by the symbol  $M_1$ . Neglecting for the present the general coefficient and common latitude factor, the two terms may be written as follows:

$$\text{term } A_{16} = 1/2 e \sin I \cos^2 \frac{1}{2}I \cos (T-s+h-p-90^\circ+2\xi-\nu) \quad (188)$$

$$\text{term } A_{23} = 3/2 e \sin I \cos I \cos (T-s+h+p-90^\circ-\nu) \quad (189)$$

The latter term, having a coefficient nearly three times as great as that of the first term, will predominate and determine the speed and period of the composite tide while the first term introduces certain inequalities in the coefficient and argument.

**122.** For brevity, let  $A$  and  $B$  represent the respective coefficients of terms  $A_{16}$  and  $A_{23}$  and let

$$\theta = T-s+h+p-90^\circ-\nu \quad (190)$$

Also let  $P$  equal the mean longitude of the lunar perigee reckoned from the lunar intersection. Then

$$p = P + \xi \quad (191)$$

We then have

$$\begin{aligned} \text{term } A_{16} &= A \cos (\theta - 2P) \\ &= A \cos 2P \cos \theta + A \sin 2P \sin \theta \end{aligned} \quad (192)$$

$$\text{term } A_{23} = B \cos \theta \quad (193)$$

$$\begin{aligned} M_1 &= A_{16} + A_{23} = (A \cos 2P + B) \cos \theta + A \sin 2P \sin \theta \\ &= (A^2 + 2AB \cos 2P + B^2)^{\frac{1}{2}} \cos \left[ \theta - \tan^{-1} \frac{A \sin 2P}{A \cos 2P + B} \right] \\ &= \frac{e \sin I \cos^2 \frac{1}{2}I}{Q_a} \cos (T-s+h+p-90^\circ-\nu-Q_u) \end{aligned} \quad (194)$$

in which

$$1/Q_a = \left[ 1/4 + 3/2 \frac{\cos I}{\cos^2 \frac{1}{2}I} \cos 2P + 9/4 \frac{\cos^2 I}{\cos^4 \frac{1}{2}I} \right]^{\frac{1}{2}} \quad (195)$$

$$Q_u = \tan^{-1} \frac{\sin 2P}{3 \cos I / \cos^2 \frac{1}{2}I + \cos 2P} \quad (196)$$

If  $I$  is given its mean value corresponding to  $\omega$ , formula (195) may be reduced to the form

$$1/Q_a = (2.310 + 1.435 \cos 2P)^{\frac{1}{2}} \quad (197)$$

Values of  $\log Q_a$  for each degree of  $P$  based upon formula (197) are given in table 9.

**123.** The period of the composite constituent  $M_1$  is very nearly an exact multiple of the period of the principal lunar constituent  $M_2$ , and for this reason the summations which are necessary for the analysis of the latter may be conveniently adapted to the analysis of the former. With other symbols as before, let

$$\theta = T - s + h - 90^\circ + \xi - \nu \quad (198)$$

Terms  $A_{16}$  and  $A_{23}$  may then be combined as follows:

$$\begin{aligned} \text{term } A_{16} &= A \cos (\theta - P) \\ &= A \cos P \cos \theta + A \sin P \sin \theta \end{aligned} \quad (199)$$

$$\begin{aligned} \text{term } A_{23} &= B \cos (\theta + P) \\ &= B \cos P \cos \theta - B \sin P \sin \theta \end{aligned} \quad (200)$$

$$\begin{aligned} M_1 &= A_{16} + A_{23} = (A+B) \cos P \cos \theta + (A-B) \sin P \sin \theta \\ &= (A^2 + 2AB \cos 2P + B^2)^{\frac{1}{2}} \cos \left[ \theta - \tan^{-1} \left( \frac{A-B}{A+B} \tan P \right) \right] \\ &= \frac{e \sin I \cos^{\frac{1}{2}} I}{Q_a} \cos(T - s + h - 90^\circ + \xi - \nu + Q) \end{aligned} \quad (201)$$

in which

$$Q = \tan^{-1} \left( \frac{5 \cos I - 1}{7 \cos I + 1} \tan P \right) \quad (202)$$

If  $I$  is given its mean value corresponding to  $\omega$ , formula (202) may be reduced to the following form which was used for computing the values of  $Q$  in table 10.

$$\tan Q = 0.483 \tan P \quad (203)$$

**124.** Formulas (194) and (201) are the same except in the method of representing the argument. The elements  $+p - Q_u$  in the first formula are replaced by  $+\xi + Q$  in the latter, but it may be shown from (196) and (202) that

$$Q_u + Q = P = p - \xi \quad (204)$$

$$p - Q_u = \xi + Q \quad (205)$$

The complete arguments are therefore equal but in formula (201) the uniformly varying element  $p$  has been transferred from the  $V$  of the argument and included in the value of  $Q$  where it is treated as a constant for a series of observations being analyzed. The speed of the argument as determined by the remaining part of the  $V$  is then exactly one-half that of the principal constituent  $M_2$  and with this assumption the summations for the latter may be adapted to the analysis of the former. It is to be noted, however, that the  $u$  in this case has a progressive forward change of nearly  $41^\circ$  each year. The true average speed of this constituent is determined by the  $V$  of formula (194) which includes the element  $p$ .

**125.** The obliquity factor for the composite  $M_1$  constituent may be expressed by the formula  $\sin I \cos^{\frac{1}{2}} I \times 1/Q_a$ . According to the work of Darwin (Scientific Papers by Sir George H. Darwin, vol. 1, p. 39) the

mean value of this factor is represented by the product  $\sin \omega \cos^2 \frac{1}{2} \omega \cos^4 \frac{1}{2} i \times \sqrt{2.307}$ , which equals  $0.3800 \times 1.52$ , or 0.5776. When deriving the node-factor formula for  $M_1$ , Darwin inadvertently omitted the factor  $\sqrt{2.307}$  and obtained the approximate equivalent of the following:

$$j(M_1) = \frac{\sin I \cos^2 \frac{1}{2} I}{\sin \omega \cos^2 \frac{1}{2} \omega \cos^4 \frac{1}{2} i} \times 1/Q_a = \frac{\sin I \cos^2 \frac{1}{2} I}{0.3800} \times 1/Q_a \quad (206)$$

Comparing the above with formula (75), it will be noted that

$$f(M_1) = f(O_1) \times 1/Q_a \quad (207)$$

Factors pertaining to constituent  $M_1$  in tables 13 and 14 are based upon the above formulas.

**126.** Because of the omission of the factor  $\sqrt{2.307}$  from formula (206), the node factors for  $M_1$  which have been in general use since this system of tidal reductions was adopted are about 50 percent greater than was originally intended, while the reciprocal reduction factors are correspondingly too small. This constituent is relatively unimportant and no practical difficulties have resulted from the omission. The  $M_1$  amplitudes as reduced from the observational data are comparable among themselves but should be increased by 50 percent to be on the same basis as the amplitudes of other constituents. The predicted tides have not been affected in the least since the node factors and reduction factors are reciprocal and compensating. The theoretical mean coefficient for this constituent with the factor  $\sqrt{2.307}$  included is 0.0317; but in order that this coefficient may be adapted for use with the tabular node factors when computing tidal forces or the equilibrium height of the tide, the coefficient 0.0209 with the factor  $\sqrt{2.307}$  excluded should be used.

**127.** Although  $M_1$  is one of the relatively unimportant constituents and the error in the node factor has caused no serious difficulties, it may be questionable whether it should be perpetuated. It is obvious, however, that any change in the present procedure would lead to much confusion unless undertaken by general agreement among all the principal organizations engaged in tidal work. By making any change applicable to the analysis of all series of observations beginning after a certain specified date it would be possible to interpret the results on the basis of the period covered by the observations without the necessity of revising all previously published amplitudes for this constituent.

#### THE $L_2$ TIDE

**128.** The composite  $L_2$  constituent is formed by combining terms  $A_{41}$  and  $A_{48}$  of formula (64). Neglecting the general coefficient and common latitude factor these terms may be written

$$\text{term } A_{41} = 1/2 e \cos^4 \frac{1}{2} I \cos (2T - s + 2h - p + 180^\circ + 2\xi - 2\nu) \quad (208)$$

$$\text{term } A_{48} = 3/4 e \sin^2 I \cos (2T - s + 2h + p - 2\nu) \quad (209)$$

A reference to table 2 will show that the mean coefficient of the first term is about four times as great as that of the latter term. The first

term will therefore predominate and determine the speed of the composite constituent.

**129.** With other symbols as before, let  $A$  and  $B$  represent the respective coefficients of the two terms and  $\theta$  the argument of the first term. We then have

$$A_{41}=A \cos \theta \quad (210)$$

$$A_{48}=B \cos (\theta+2P-180^\circ)=-B \cos (\theta+2P) \quad (211)$$

$$\begin{aligned} L_2 &= A_{41} + A_{48} = (A - B \cos 2P) \cos \theta + B \sin 2P \sin \theta \\ &= (A^2 - 2AB \cos 2P + B^2)^{\frac{1}{2}} \cos \left[ \theta - \tan^{-1} \frac{B \sin 2P}{A - B \cos 2P} \right] \\ &= 1/2 e^{-\frac{\cos^4 \frac{1}{2}I}{R_a}} I \cos (2T - s + 2h - p + 180^\circ + 2\xi - 2\nu - R) \end{aligned} \quad (212)$$

in which

$$1/R_a = (1 - 12 \tan^2 \frac{1}{2}I \cos 2P + 36 \tan^4 \frac{1}{2}I)^{\frac{1}{2}} \quad (213)$$

$$R = \tan^{-1} \frac{\sin 2P}{1/6 \cot^2 \frac{1}{2}I - \cos 2P} \quad (214)$$

Values of  $\log R_a$  and  $R$  computed from the above formulas are given in tables 7 and 8, respectively.

**130.** The obliquity factor for the composite  $L_2$  constituent may be expressed by the formula  $\cos^4 \frac{1}{2}I \times 1/R_a$ . The mean value of  $1/R_a$  is approximately unity, and in accord with the Darwinian system the mean for the entire obliquity factor is taken as the product  $\cos^4 \frac{1}{2}\omega \cos^4 \frac{1}{2}i$ , which equals 0.9154 and is the same as the mean value of the obliquity factor for the principal constituent  $M_2$ . Multiplying this by the elliptic factor  $\frac{1}{2}e$  gives 0.0251 as the mean constituent coefficient.

**131.** The node factor formula for constituent  $L_2$  based upon the above mean for the obliquity factor is as follows:

$$f(L_2) = \frac{\cos^4 \frac{1}{2}I}{0.9145} \times 1/R_a = f(M_2) \times 1/R_a \quad (215)$$

Node factors for constituent  $L_2$  based upon the above formula are included in table 14 for the middle of each year from 1850 to 1999, inclusive. The logarithms of the reciprocal reduction factors covering the period 1900 to 2000 are contained in table 13.

#### LUNISOLAR K<sub>1</sub> AND K<sub>2</sub> TIDES

**132.** Lunar diurnal term  $A_{22}$  of formula (63) and solar diurnal term  $B_{22}$  of formula (186) have the same speed. Together they form the lunisolar  $K_1$  constituent. Also, lunar semidiurnal term  $A_{47}$  of formula (64) and solar semidiurnal term  $B_{47}$  of formula (187) have speeds exactly twice that of constituent  $K_1$  and together form the lunisolar  $K_2$  constituent. In order that the solar terms may have the same general coefficient as the lunar terms, the solar factor  $U_1/U$ , which will be designated by the symbol  $S'$ , will be transferred from the general coefficient of the solar terms and included in the constituent coefficients. Then, neglecting the general coefficient and

latitude factors common to the terms combined, we have the following formulas in which numerical values from table 1 have been substituted for constant quantities.

$$\begin{aligned} \text{term } A_{22} &= (1/2 + 3/4e^2) \sin 2I \cos (T+h-90^\circ - \nu) \\ &= 0.5023 \sin 2I \cos (T+h-90^\circ - \nu) \end{aligned} \quad (216)$$

$$\begin{aligned} \text{term } B_{22} &= (1/2 + 3/4e_1^2) S' \sin 2\omega \cos (T+h-90^\circ) \\ &= 0.1681 \cos (T+h-90^\circ) \end{aligned} \quad (217)$$

$$\begin{aligned} \text{term } A_{47} &= (1/2 + 3/4e^2) \sin^2 I \cos (2T+2h-2\nu) \\ &= 0.5023 \sin^2 I \cos (2T+2h-2\nu) \end{aligned} \quad (218)$$

$$\begin{aligned} \text{term } B_{47} &= (1/2 + 3/4e_1^2) S' \sin^2 \omega \cos (2T+2h) \\ &= 0.0365 \cos (T+2h) \end{aligned} \quad (219)$$

**133.** Taking first the diurnal terms, let  $A$  represent the lunar coefficient  $0.5023 \sin 2I$  and let  $B$  represent the solar coefficient  $0.1681$ . We then have

$$\begin{aligned} A_{22} &= A \cos (T+h-90^\circ - \nu) \\ &= A \cos \nu \cos (T+h-90^\circ) + A \sin \nu \sin (T+h-90^\circ) \end{aligned} \quad (220)$$

$$B_{22} = B \cos (T+h-90^\circ) \quad (221)$$

$$\begin{aligned} K_1 &= (A \cos \nu + B) \cos (T+h-90^\circ) + A \sin \nu \sin (T+h-90^\circ) \\ &= (A^2 + 2AB \cos \nu + B^2)^{\frac{1}{2}} \cos \left[ T+h-90^\circ - \tan^{-1} \frac{A \sin \nu}{A \cos \nu + B} \right] \\ &= C_1 \cos (T+h-90^\circ - \nu') \end{aligned} \quad (222)$$

in which

$$\begin{aligned} C_1 &= (A^2 + 2AB \cos \nu + B^2)^{\frac{1}{2}} \\ &= (0.2523 \sin^2 2I + 0.1689 \sin 2I \cos \nu + 0.0283)^{\frac{1}{2}} \end{aligned} \quad (223)$$

$$\nu' = \tan^{-1} \frac{A \sin \nu}{A \cos \nu + B} = \tan^{-1} \frac{\sin 2I \sin \nu}{\sin 2I \cos \nu + 0.3347} \quad (224)$$

Values of  $\nu'$  for each degree of  $N$ , which is the longitude of the moon's node, are included in table 6.

**134.** The obliquity factor for  $K_1$  will be taken to include the entire coefficient  $(A^2 + 2AB \cos \nu + B^2)^{\frac{1}{2}}$  and its mean value will be taken as the mean of the product  $(A^2 + 2AB \cos \nu + B^2)^{\frac{1}{2}} \cos \nu'$ .

From (224) we may obtain

$$\cos \nu' = (A \cos \nu + B) / (A^2 + 2AB \cos \nu + B^2)^{\frac{1}{2}} \quad (225)$$

Then for mean value of coefficient of  $K_1$

$$\begin{aligned} [(A^2 + 2AB \cos \nu + B^2)^{\frac{1}{2}} \cos \nu']_0 &= [A \cos \nu + B]_0 \\ &= [0.5023 \sin 2I \cos \nu + 0.1681]_0 = 0.5305 \end{aligned} \quad (226)$$

the numerical mean for  $\sin 2I \cos \nu$  being obtained from formula (68). For the node factor of  $K_1$  divide the coefficient of (222) by its mean value and obtain

$$\begin{aligned} f(K_1) &= (0.2523 \sin^2 2I + 0.1689 \sin 2I \cos \nu + 0.0283)^{\frac{1}{2}} / 0.5305 \\ &= (0.8965 \sin^2 2I + 0.6001 \sin 2I \cos \nu + 0.1006)^{\frac{1}{2}} \end{aligned} \quad (227)$$

The node factors for the middle of each year 1850 to 1999 are included in table 14. Logarithms of the reciprocal reduction factors for each tenth of a degree of  $I$  are given in table 12.

135. The semidiurnal terms  $A_{47}$  and  $B_{47}$  may be combined in a similar manner. Letting  $A$  represent the lunar coefficient 0.5023  $\sin^2 I$  and  $B$  the solar coefficient 0.0365, we have

$$\begin{aligned} A_{47} &= A \cos (2T+2h-2\nu) \\ &= A \cos 2\nu \cos (2T+2h) + A \sin 2\nu \sin (2T+2h) \end{aligned} \quad (228)$$

$$B_{47} = B \cos (2T+2h) \quad (229)$$

$$\begin{aligned} K_2 &= (A \cos 2\nu + B) \cos (2T+2h) + A \sin 2\nu \sin (2T+2h) \\ &= (A^2 + 2AB \cos 2\nu + B^2)^{\frac{1}{2}} \cos \left[ 2T+2h - \tan^{-1} \frac{A \sin 2\nu}{A \cos 2\nu + B} \right] \\ &= C_2 \cos (2T+2h-2\nu'') \end{aligned} \quad (230)$$

in which

$$\begin{aligned} C_2 &= (A^2 + 2AB \cos 2\nu + B^2)^{\frac{1}{2}} \\ &= (0.2523 \sin^4 I + 0.0367 \sin^2 I \cos 2\nu + 0.0013)^{\frac{1}{2}} \end{aligned} \quad (231)$$

$$2\nu'' = \tan^{-1} \frac{A \sin 2\nu}{A \cos 2\nu + B} = \tan^{-1} \frac{\sin^2 I \sin 2\nu}{\sin^2 I \cos 2\nu + 0.0727} \quad (232)$$

Values for  $2\nu''$  for each degree of  $N$  are included in table 6.

136. The obliquity factor for  $K_2$  will be taken to include the entire coefficient  $(A^2 + 2AB \cos 2\nu + B^2)^{\frac{1}{2}}$  and its mean value will be taken as the mean of the product  $(A^2 + 2AB \cos 2\nu + B^2)^{\frac{1}{2}} \cos 2\nu''$ . From (232)

$$\cos 2\nu'' = (A \cos 2\nu + B) / (A^2 + 2AB \cos 2\nu + B^2)^{\frac{1}{2}} \quad (233)$$

Then for the mean value of coefficient of  $K_2$

$$\begin{aligned} [(A^2 + 2AB \cos 2\nu + B^2)^{\frac{1}{2}} \cos 2\nu'']_0 &= [A \cos 2\nu + B]_0 \\ &= [0.5023 \sin^2 I \cos 2\nu + 0.0365]_0 = 0.1151 \end{aligned} \quad (234)$$

the numerical mean for  $\sin^2 I \cos 2\nu$  being obtained from formula (71). For the node factor of  $K_2$  divide the coefficient of (230) by its mean value and obtain

$$\begin{aligned} f(K_2) &= (0.2523 \sin^4 I + 0.0367 \sin^2 I \cos 2\nu + 0.0013)^{\frac{1}{2}} / 0.1151 \\ &= (19.0444 \sin^4 I + 2.7702 \sin^2 I \cos 2\nu + 0.0981)^{\frac{1}{2}} \end{aligned} \quad (235)$$

See table 14 for node factors and table 12 for reciprocal reduction factors.

#### METEOROLOGICAL AND SHALLOW-WATER TIDES

137. In addition to the elementary constituents obtained from the development of the tide-producing forces of the moon and the sun, there are a number of harmonic terms that have their origin in meteorological changes or in shallow-water conditions. Variations in temperature, barometric pressure, and in the direction and force of the wind may be expected to cause fluctuations in the water level. Although in general such fluctuations are very irregular, there are some seasonal and daily variations which occur with a rough periodicity that admit of being expressed by harmonic terms. The meteorological constituents usually taken into account in the tidal analysis are

S<sub>a</sub>, S<sub>sa</sub>, and S<sub>1</sub> with periods corresponding respectively to the tropical year, the half tropical year, and the solar day. These constituents are represented also by terms in the development of the tide-producing force of the sun but they are considered of greater importance as meteorological tides. S<sub>sa</sub> occurs in the development of the principal solar force while S<sub>a</sub> and S<sub>1</sub> would appear in a development involving the 4th power of the solar parallax (par. 119). In the analysis of tide observations both S<sub>a</sub> and S<sub>sa</sub> are usually found to have an appreciable effect on the water level. Constituent S<sub>1</sub> is relatively of little importance in its effect on the height of the tide but has been more noticeable in the velocity of off-shore tidal currents, probably as a result of periodic land and sea breezes.

**138.** The shallow-water constituents result from the fact that when a wave runs into shallow water its trough is retarded more than its crest and the wave loses its simple harmonic form. The shallow-water constituents are classified as overtides and compound tides, the overtide having a speed that is an exact multiple of one of the elementary constituents and the compound tide a speed that equals the sum or difference of the speeds of two or more elementary constituents.

**139.** The overtides were so named because of their analogy to the overtones in musical sounds and they may be considered as the higher harmonics of the fundamental tides. The only overtides usually taken into account in tidal work are the harmonics of the principal lunar and solar semidiurnal constituents M<sub>2</sub> and S<sub>2</sub>, the lunar series being designated by the symbols M<sub>4</sub>, M<sub>6</sub>, and M<sub>8</sub>, and the solar series by S<sub>4</sub>, S<sub>6</sub>, and S<sub>8</sub>. The subscript indicates the number of periods in the constituent day. These overtides with their arguments and speeds are included in table 2a, the arguments and speeds being taken as exact multiples of those of the fundamental constituent. There are no theoretical expressions for the coefficients of the overtides but it is assumed that the amplitudes of the lunar series undergo variations due to changes in the longitude of the moon's node which are analogous to those in the fundamental tide. The node factors for M<sub>4</sub>, M<sub>6</sub>, and M<sub>8</sub>, respectively, are taken as the square, the cube, and the fourth power of the corresponding factor for M<sub>2</sub>. For the solar terms this factor is always zero.

**140.** Compound tides were suggested by Helmholtz's theory of sound waves. Innumerable combinations are possible but the principal elementary constituents involved are M<sub>2</sub>, S<sub>2</sub>, N<sub>2</sub>, K<sub>1</sub>, and O<sub>1</sub>. Table 2a includes the compound tides listed in International Hydrographic Bureau Special Publication No. 26, which is a compilation of the tidal harmonic constants for the world. The argument of a compound tide equals the sum or difference of the arguments of the elementary constituents of which it is compounded. The node factor is taken as the product of the node factors of the same constituents. Table 2a contains the arguments, speeds, and node factors of these tides.

**141.** Omitted from table 2a are a number of compound tides which have the same speeds as elementary constituents included in table 2. Thus, 2MS<sub>2</sub>, compounded by formula 2M<sub>2</sub>—S<sub>2</sub>, has the same speed as constituent μ<sub>2</sub> represented by term A<sub>45</sub> of formula (64). Considered as a compound tide there would be a small difference in the *u* of the argument and also in the node factor. Since there is no practical way of separating the elementary constituent from the compound

tide of the same speed, this has been treated solely as an elementary constituent. Constituent MSf represented by term  $A_5$  of formula (62) has the same speed as a compound tide of formula  $S_2 - M_2$ . This constituent is relatively unimportant and it makes little difference whether treated as an elementary or a compound tide. Following the previous practice in this office it is treated in the harmonic analysis as a compound tide with corresponding argument and node factor. When included in the computation of tidal forces, however, the argument and node factor indicated in table 2 should be used.

## ANALYSIS OF OBSERVATIONS

### HARMONIC CONSTANTS

**142.** In the preceding discussion it has been shown that under the equilibrium theory the height of a theoretical tide at any place can be expressed mathematically by the sum of a number of harmonic terms involving certain astronomical data and the location of the place. It has also been pointed out that for obvious reasons the actual tide of nature does not conform to the theoretical equilibrium tide. However, the tide of nature can be conceived as being composed of the sum of a number of harmonic constituents having the same periods as those found in the tide-producing force. Although the complexity of the tidal movement is too great to permit a theoretical computation based upon astronomical conditions only, it is possible through the analysis of observational data at any place to obtain certain constants which can be introduced into the theoretical formulas and thus adapt them for the computation of the tide for any desired time.

**143.** In the formulas obtained for the height of the equilibrium tide each constituent term consists of the product of a coefficient by the cosine of an argument. For corresponding formulas expressing the actual height of the tide at any place, the entire theoretical coefficient including the latitude factor and the common general coefficient is replaced by a coefficient determined from an analysis of observational data for the station. This tidal coefficient, which is known as the *amplitude* of the constituent, is assumed to be subject to the same variations arising from changes in the longitude of the moon's node as the coefficient of the corresponding term in the equilibrium tide. The amplitude pertaining to any particular year is usually designated by the symbol  $R$  while its mean value for an entire node period is represented by the symbol  $H$ . Amplitudes derived directly from an analysis of a limited series of observations must be multiplied by the reduction factor  $F$  (par. 78) to obtain the mean amplitudes of the harmonic constants. For the prediction of tides, the mean amplitudes must be multiplied by the node factor  $f$  (par. 77) to obtain the amplitudes pertaining to the year for which the predictions are to be made.

**144.** The phases of the constituents of the actual tide do not in general coincide with the phases of the corresponding constituents of the equilibrium tide but there may be lags varying from 0 to  $360^\circ$ . The interval between the high water phase of an equilibrium constituent and the following high water of the corresponding constituent in the actual tide is known as the *phase lag* or *epoch* of the constituent and is represented by the symbol  $\kappa$  (kappa) which is expressed in angular measure. The amplitudes and epochs together are called harmonic constants and are the quantities sought in the harmonic analysis of tides. Each locality has a separate set of harmonic constants which can be derived only from observational data but which remain the same over a long period of time provided there are no

physical changes in the region that might affect the tidal conditions.

**145.** If we let  $y_1$  equal the height of one of the tidal constituents as referred to mean sea level, it may be represented by the following formula:

$$y_1 = fH \cos (E - \kappa) = fH \cos (V + u - \kappa) \quad (236)$$

The combination symbol  $V + u$  is the equivalent of  $E$  and represents the argument or phase of the equilibrium constituent.

**146.** Formula (236) is illustrated graphically in figure 7 by a cosine curve with amplitude  $fH$ . The horizontal line represents mean sea level and the vertical line through  $T$  may be taken to indicate any instant of time under consideration. If the point  $M$  represents the time when the constituent argument equals zero, the interval from  $M$  to the following high water of the constituent will be the epoch  $\kappa$ . The interval from the preceding high water to  $M$  is measured by the explement of  $\kappa$  which may be expressed as  $-\kappa$ . The phase of the constituent argument at time  $T$  is reckoned from  $M$  and is expressed by the symbol  $(V + u)$ . The phase of the constit-




FIGURE 7.

uent itself at this time is reckoned from the preceding high water and therefore equals  $(V + u - \kappa)$ .

#### OBSERVATIONAL DATA

**147.** The most satisfactory observational data for the harmonic analysis are from the record of an automatic tide gage that traces a continuous curve from which the height of the tide may be scaled at any desired interval of time. This record is usually tabulated to give the height of the tide at each solar hour of the series in the kind of time normally used at the place. It is important, however, that the time should be accurate and that the same system be used for the entire series of observations regardless of the fact that daylight saving time may have been adopted temporarily for other purposes during a portion of the year. When the continuous record from an automatic gage is not available, hourly heights of the tide as observed by other methods may be used. The record should be complete with each hour of the series represented. If a part of the record has been lost, the hiatus may be filled by interpolated values; or, if the gap is very extensive, the record may be broken up into shorter series which do not include the defective portion.

**148.** If hourly heights have not been observed but a record of high and low waters is available, an approximate evaluation of the more important constituents may be obtained by a special treatment. The results, however, are not nearly as satisfactory as those obtained from the hourly heights.

**149.** Although the hourly interval for the tabulated heights of the tide has usually been adopted as most convenient and practicable for the purposes of the harmonic analysis, a greater or less interval might be used. A shorter interval would cause a considerable increase in the amount of work without materially increasing the accuracy of the results for the constituents usually sought. However, if an attempt were made to analyze for the short period seiches a closer interval would be necessary. An interval greater than one hour would lessen the work of the analysis but would not be sufficient for the satisfactory development of the overtides.

**150.** In selecting the length of series of observations for the purpose of the analysis, consideration has been given to the fact that the procedure is most effective in separating two constituents from each other when the length of series is an exact multiple of the synodic period of these constituents. By synodic period is meant the interval between two consecutive conjunctions of like phases. Thus, if the speeds of the two constituents in degrees per solar hour are represented by  $a$  and  $b$ , the synodic period will equal  $360^\circ / (a - b)$  hours. If there were only two constituents in the tide the best length of series could be easily fixed, but in the actual tide there are many constituents and the length of series most effective in one case may not be best adapted to another case. It is therefore necessary to adopt a length that is a compromise of the synodic periods involved, consideration being given to the relative importance of the different constituents.

**151.** Fortunately, the exact length of series is not of essential importance and for convenience all series may be taken to include an integral number of days. Theoretically, different lengths of series should be used in seeking different constituents, but practically it is more convenient to use the same length for all constituents, an exception being made in the case of a very short series. The longer the series of observations the less important is its exact length. Also the greater the number of synodic periods of any two constituents the more nearly complete will be their separation from each other. Constituents like  $S_2$  and  $K_2$  which have nearly equal speeds and a synodic period of about 6 months will require a series of not less than 6 months for a satisfactory separation. On the other hand, two constituents differing greatly in speed such as a diurnal and a semidiurnal constituent may have a synodic period that will not greatly exceed a day, and a moderately short series of observations will include a relatively large number of synodic periods. For this reason, when selecting the length of series no special consideration need be given to the effect of a diurnal and a semidiurnal constituent upon each other.

**152.** The following lengths of series have been selected as conforming approximately to multiples of synodic periods involving the more important constituents—14, 15, 29, 58, 87, 105, 134, 163, 192, 221, 250, 279, 297, 326, 355, and 369 days. The 369-day series is considered as a standard length to be used for the analysis whenever observations covering this period are available. This length conforms very closely with multiples of the synodic periods of practically all of the short-

period constituents and is well adapted for the elimination of seasonal meteorological effects. When observations at any station are available for a number of years, it is desirable to have separate analyses made for different years in order that the results may be compared and serve as a check on each other. Although not essential, there are certain conveniences in having each such series commence on January 1 of the year, regardless of the fact that series of consecutive years may overlap by several days because the length of series is a little longer than the calendar year.

153. If the available observations cover a period less than 369 days, the next longest series listed above which is fully covered by the observations will usually be taken, any extra days of observations being rejected. However, if the observations lack only a few hours of being equal to the next greater length, it may be advantageous to extrapolate additional hourly heights to complete the larger series. The 29-day series is usually considered as a minimum standard for short series of observations. This is a little shorter than the synodical month and a little longer than the nodical, tropical, and anomalistic months. It is the minimum length for a satisfactory development of the more important constituents.

154. For observations of less than 29 days, but more than 14 days, provisions are made for an analysis of a 14-day series for the diurnal constituents and a 15-day series for the semidiurnal constituents, the first conforming to the synodic period of constituents  $K_1$  and  $O_1$ , and the latter to the synodic period of  $M_2$  and  $S_2$ . Through special treatment involving a comparison with another station, it is possible to utilize even shorter series of observations. This treatment is rarely required in case of tide observations but is useful in connection with tidal currents where observations may be limited to only a few days.

#### SUMMATIONS FOR ANALYSIS

155. The first approximate separation of the constituents of the observed tide is accomplished by a system of summations, separate summations being made for all constituents with incommensurable periods. Designating the constituent sought by  $A$ , assume that the entire series of observations is divided into periods equal to the period of  $A$  and each period is subdivided into a convenient number of equal parts, the subdivisions of each period being numbered consecutively beginning with zero at the initial instant of each period. All subdivisions of like numbers will then include the same phase of constituent  $A$  but different phases for all other constituents with incommensurable speeds. The subdivisions will also include irregular variations arising from meteorological causes. By summing and averaging separately all heights corresponding to each of the numbered subdivisions over a sufficient length of time, the effects of constituents with incommensurable periods as well as the meteorological variations will be averaged out leaving intact constituent  $A$  with its overtides.

156. The principle just described for separating constituent  $A$  from the rest of the tide is applicable if the original periods into which the series of observations is divided are taken as some multiple of constituent  $A$  period. In general practice, that multiple of the constituent period which is most nearly equal to the solar day is taken as the unit. This is the constituent day and includes one or more

periods according to whether the constituent is diurnal, semidiurnal, etc. The constituent day is divided into 24 equal parts, the beginning of each part being numbered consecutively from 0 to 23 and these are known as constituent hours.

157. To carry out strictly the plan described above would require separate tabulations of the heights of the tide at different intervals for all constituents of incommensurable periods, a procedure involving an enormous amount of work. In actual practice the tabulated solar hourly heights are used for all of the summations, these heights being assigned to the nearest constituent hour. Corrections are afterwards applied to take account of any systematic error in this approximation.

158. There are two systems for the distribution and assignment of the solar hourly heights which differ slightly in detail. In the system ordinarily used and which is sometimes called the standard system, each solar hourly height is used once, and once only, by being assigned to its nearest constituent hour. By this system some constituent hours will be assigned two consecutive solar hourly heights or receive no assignment according to whether the constituent day is longer or shorter than the solar day. In the other system of distribution, each constituent hour receives one and only one solar hourly height necessitating the occasional rejection or double assignment of a solar hourly height. The difference in the results obtained from the two systems is practically negligible but the first system is generally used as it affords a quick method of checking the summations.

#### STENCILS

159. The distribution of the tabulated solar hourly heights of the tide for the purpose of the harmonic analysis is conveniently accomplished by a system of stencils (fig. 10) which were devised by L. P. Shidy of the Coast and Geodetic Survey early in 1885 (Report of U. S. Coast and Geodetic Survey, 1893, vol. I, p. 108). Although the original construction of the stencils involves considerable work, they are serviceable for many years and have resulted in a very great saving of labor. These stencils are cut from the same forms which are used for the tabulation of the hourly heights of the tide and 106 sheets are required for the summation of a 369-day series of observations for a single constituent. Separate sets are provided for different constituents. Constituents with commensurable periods are included in a single summation and no stencils are required for constituents,  $S_1$ ,  $S_2$ ,  $S_4$ , etc.

160. The use of the stencils makes a standardized form for the tabulation of the hourly heights essential. This form (fig. 9) is a sheet 8 by  $10\frac{1}{2}$  inches, with spaces arranged for the tabulation of the 24 hourly heights of each day in a vertical column, with 7 days of record on each page. The hours of the day are numbered consecutively from  $0^{\text{h}}$  at midnight to  $23^{\text{h}}$  at 11 p. m. When the tabulated heights are entered, each day is indicated by its calendar date and also by a serial number commencing with 1 as the first day of series. The days on the stencil sheets are numbered serially to correspond with the tabulation sheets and may be used for any series regardless of the calendar dates.

161. The openings in the stencils are numbered to indicate the constituent hours that correspond most closely with the times of the

height values showing through the openings when the stencil is applied to the tabulations. Openings applying to the same constituent hour are connected by a ruled line which clearly indicates to the eye the tabular heights which are to be summed together. For convenience in construction two stencil sheets are prepared for each page of tabulations, one sheet providing for the even constituent hours and the other sheet for the odd constituent hours.

**162.** The stencils are adapted for use with tabulations made in any kind of time provided the time used is uniform for the entire series of observations. For convenience the tabulations are usually made in the standard time of the place. The series to be analyzed, however, must commence with the zero hour of the day and this is also taken as the zero constituent hour for each constituent. Successive solar hours will fall either earlier or later than the corresponding constituent hour according to whether the constituent day is longer or shorter than the solar day.

**163.** For the construction of the stencils it is necessary to calculate the constituent hour that most nearly coincides with each solar hour of the series.

Let  $a$ =speed or rate of change in argument of constituent sought in degrees per solar hour.

$p$ =number of constituent periods in constituent day; 1 for diurnal tides, 2 for semidiurnal tides, etc.

$sh$ =number of solar hour reckoned from 0 at beginning of each solar day.

$shs$ =number of solar hour reckoned from 0 at beginning of series.

$dos$ =day of series counting from 1 as the first day.

$ch$ =number of constituent hour reckoned from 0 at beginning of each constituent day.

$chs$ =number of constituent hour reckoned from 0 at beginning of series.

Then

$$1 \text{ constituent period} = \frac{360}{a} \text{ solar hours.} \quad (237)$$

$$1 \text{ constituent day} = \frac{360p}{a} \text{ solar hours.} \quad (238)$$

$$1 \text{ constituent hour} = \frac{15p}{a} \text{ solar hours.} \quad (239)$$

$$1 \text{ solar hour} = \frac{a}{15p} \text{ constituent hours.} \quad (240)$$

Therefore,

$$(chs) = \frac{a}{15p} (shs) = \frac{a}{15p} [24\{(dos)-1\} + (sh)] \quad (241)$$

**164.** The above formula gives the constituent hour of the series ( $chs$ ) corresponding to any solar hour of the series ( $shs$ ). The observed heights of the tide being tabulated for the exact solar hours of the day, the ( $shs$ ) with which we are concerned will represent successive integers counting from 0 at the beginning of the series. The ( $chs$ ) as derived from the formula will generally be a mixed number. As

it is desired to obtain the integral constituent hour corresponding most nearly with each solar hour, the (*chs*) should be taken to the nearest integer by rejecting a fraction less than 0.5, or counting as an extra hour a fraction greater than 0.5, or adopting the usual rule for computations if the fraction is exactly 0.5. The constituent hour of the constituent day (*ch*) required for the construction of the stencils may be obtained by rejecting multiples of 24 from the (*chs*).

**165.** In the application of the above formula it will be found that the integral constituent hour will differ from the corresponding solar hour by a constant for a succession of solar hours, and then, with the difference changed by one, it will continue as a constant for another group of solar hours, etc. This fact is an aid in the preparation of a table of constituent hours corresponding to the solar hours of the series, as it renders it unnecessary to make an independent calculation for each hour. Instead of using the above formula for each value the time when the difference between the solar and constituent hours changes may be determined. The application of the differences to the solar hours will then give the desired constituent hours.

**166.** Formula (241) is true for any value of (*shs*), whether integral or fractional. It represents the constituent time of any instant in the series of observations in terms of the solar time of that same instant, both kinds of time being reckoned from the beginning of the series as the zero hour. The difference between the constituent and the solar time of any instant may therefore be expressed by the following formula:

$$\text{Difference} = \frac{a}{15p} (\text{shs}) - (\text{shs}) = \frac{a - 15p}{15p} (\text{shs}) \quad (242)$$

**167.** If the constituent day is shorter than the solar day, the speed *a* will be greater than  $15p$ , and the constituent hour as reckoned from the beginning of the series will be greater than the solar hour of the same instant. If the constituent day is longer than the solar day the constituent hour at any instant will be less than the solar hour of the same instant. At the beginning of the series the difference between the constituent and solar time will be zero, but the difference will increase uniformly with the time of the series. As long as the difference does not exceed 0.5 of an hour the integral constituent hours will be designated by the same ordinals as the integral solar hours with which they most nearly coincide. Differences between 0.5 and 1.5 will be represented by the integer 1, differences between 1.5 and 2.5 by the integer 2, etc. If we let *d* represent the integral difference, the time when the difference changes from  $(d-1)$  to *d*, will be the time when the difference derived from formula (242) equals  $(d-0.5)$ . Substituting this in the formula, we may obtain

$$(\text{shs}) = \frac{15p}{a - 15p} (d - 0.5) \quad (243)$$

in which (*shs*) represents the solar time when the integral difference between the constituent and solar time will change by one hour from  $(d-1)$  to *d*. By substituting successively the integers 1, 2, 3, etc., for *d* in the formula (243) the time of each change throughout the series may be obtained. The value of (*shs*) thus obtained will

generally be a mixed number; that is to say, the times of the changes will usually come between integral solar hours. The first integral solar hour after the change will be the one to which the new difference will apply if the usual system of distribution is to be adopted. In this case we are not concerned with the exact value of the fractional part of ( $shs$ ) but need note only the integral hours between which this value falls.

168. If, however, the second system of distribution should be desired, it should be noted whether the fractional part of ( $shs$ ) is greater or less than 0.5 hour. With a constituent day shorter than the solar day and the differences of formula (242) increasing positively, the application of the differences to the consecutive solar hours will result in the jumping or omission of a constituent hour at each change of difference. Under the second system of distribution each of the hours must be represented, and it will therefore be necessary in this case to apply two consecutive differences to the same solar hour to represent two consecutive constituent hours. The solar hour selected for this double use will be the one occurring nearest to the time of change of differences. If the fractional part of the ( $shs$ ) in (243) is less than 0.5 hour, the old and new differences will both be applied to the preceding integral solar hour; but if the fraction is greater than 0.5 hour the old and new differences will be applied to the integral solar hour following the change.

169. With a constituent day longer than the solar day and the differences of formula (242) increasing negatively, the application of the differences to the consecutive solar hours will result in two solar hours being assigned to the same constituent hour at each change of differences. Under the second system of distribution this must be avoided by the rejection of one of the solar hours. In this case the integral solar hour nearest the time of change will be rejected, since at the time of change the difference between the integral and the true difference is a maximum. Thus, if the fractional part of the ( $shs$ ), is less than 0.5 hour, the preceding solar hour will be rejected; but if the fraction is greater than 0.5 hour the next following solar hour will be rejected.

170. Table 31, computed from formula (243), gives the first solar hour of the group to which each difference applies when the usual system of distribution is adopted. Multiples of 24 have been rejected from the differences, since we are concerned only with the constituent hour of the constituent day rather than with the constituent hour of the series, and these differences may be applied directly to the solar hours of the day. For convenience equivalent positive and negative differences are given. By using the negative difference when it does not exceed the solar hour to which it is to be applied, and at other times using the positive difference, the necessity for adding or rejecting multiples of 24 hours from the results is avoided.

171. The tabulated solar hour is the integer hour that immediately follows the value for the ( $shs$ ) is formula (243). An asterisk (\*) indicates that the fractional part of the ( $shs$ ) exceeds 0.5, and that the tabular hour is therefore the one nearest the exact value of ( $shs$ ). If the second system for the distribution of the hourly heights is adopted, the solar hours marked with the asterisk will be used with both old and new difference to represent two constituent hours, or will be rejected altogether according to whether the constituent day

is shorter or longer than the solar day. If the tabular hour is unmarked, the same rule of double use or rejection will apply to the untabulated solar hour immediately preceding the tabular unmarked hour. For the ordinary stencils no attention need be given to the asterisks. By the formula constituents with commensurable periods will have the same tabular values, and no distinction is made in the construction of the stencils. Thus, stencils for constituent M serve not only for  $M_2$  but also for  $M_3$ ,  $M_4$ ,  $M_6$ , etc.

172. For the construction of a set of stencils for any constituent a preliminary set of the hourly height forms is prepared with days of series numbered consecutively beginning with 1 and each hourly height space numbered with its constituent hour as derived by the differences in table 31. The even and odd constituent hours are then transferred to separate sets of forms and the marked spaces cut out. In the Coast and Geodetic Survey this is done by a small machine with a punch operated by a hand lever. Spaces corresponding to the same constituent hour are connected by ruled lines which are numbered the same as the hours represented. Black ruling with red numbering is recommended, the red emphasizing the distinction between these numbers and the tabulated hourly heights which are to be summed.

173. When in use the stencils are placed one at a time on the sheets of tabulated heights, with days of series on stencils matching those on the tabulations, and all heights on the page corresponding to each constituent hour are then summed separately. For constituent S no stencils are necessary as the constituent hours in this case are identical with the solar hours. For constituents K, P, R, and T with speeds differing little from that of S, the lines joining the hourly spaces frequently become horizontal and the marginal sum previously obtained for constituent S becomes immediately available for the summation at hand. In these cases a hole in the margin of the stencil for the sum replaces the holes for the individual heights covered by the sum.

#### SECONDARY STENCILS

174. After the sums for certain principal constituents have been obtained by the stencils described in the preceding section, which for convenience will be called the primary stencils, the summations for other constituents may be abbreviated by the use of secondary stencils which are designed to regroup the hourly page sums already obtained for one constituent into new combinations conforming to the periods of other constituents. Certain irregularities are introduced by the process, but in a long series, such as 369 days, these are for the most part eliminated, and the resulting values for the harmonic constants compare favorably with those obtained by use of the primary stencils directly, the differences in the results obtained by the two methods being negligible. For short series the irregularities are less likely to be eliminated, and since the labor of summing for such a series is relatively small, the abbreviated form of summing is not recommended. As the length of series increases the saving in labor by the use of the secondary stencils increases, while the irregularities due to the short process tend to disappear. It is believed that the use of the secondary stencils will be found advantageous for all series more than 6 months in length.

175. In the primary summations there are obtained 24 sums for each page of tabulations, representing the 24 constituent hours of a constituent day. In general each sum will include 7 hourly heights, and the average interval between the first and last heights will be 6 constituent days. A few of the sums may, however, include a greater or less number of hourly heights within limits which may be a day greater or less than 6 constituent days.

176. Let the constituent for which summations have been made by use of the primary stencils be designated as  $A$  and the constituent which is to be obtained by use of the secondary stencils as  $B$ . For convenience let it be first assumed that the heights included in the sums for constituent  $A$  refer to the exact  $A$ -hours. This assumption is true for constituent  $S$  but only approximately true for the other constituents. It is now proposed to assign each hourly page sum obtained for constituent  $A$  to the integral  $B$ -hour with which it most nearly coincides. Constituent  $A$  and constituent  $B$ -hours separate at a uniform rate, and the proposed assignment will depend upon the relation of the hours on the middle day of each page of tabulations. The tabulated hourly heights on each full page of record run from zero (0) solar hour on the first day to the 23d solar hour on the seventh or last day of the page. The middle of the record on each such page is therefore at 11.5 solar hours on the fourth day, or 83.5 solar hours from the beginning of the page of record.

177. Let  $a$  and  $b$  represent the hourly speeds of the constituents  $A$  and  $B$ , respectively, and  $p$  and  $p_1$  their respective subscripts, and let  $n$  equal the number of the page of tabulation under consideration, beginning with number one as the first page.

The middle of page  $n$  will then be

$$[168(n-1) + 83.5] \text{ or } (168n - 84.5) \text{ solar hours} \quad (244)$$

from the beginning of the series.

Since one solar hour equals  $a/15p$  constituent  $A$ -hours (formula 240), the middle of page  $n$  will also correspond to

$$(168n - 84.5) \frac{a}{15p} \text{ constituent } A\text{-hours} \quad (245)$$

from the beginning of the series.

As there are 24 constituent hours in each constituent day, the middle constituent  $A$ -day of each page will commence 12 constituent  $A$ -hours earlier than the time represented by the middle of the page, or at

$$[(168n - 84.5) \frac{a}{15p} - 12] \text{ constituent } A\text{-hours} \quad (246)$$

from the beginning of the series.

178. The 24 integral constituent  $A$ -hours of the middle constituent day of the page will therefore be the integral constituent  $A$ -hours which immediately follow the time indicated by the last formula. The numerical value of this formula will usually be a mixed number. Let  $f$  equal the fractional part, and let  $m$  be an integer representing the number of any integral constituent hour according to its order in the middle constituent day of each page. For each page  $m$  will have

successive values from 1 to 24. The integral constituent *A*-hours falling within the middle constituent day of each page of tabulations will then be represented by the general formula.

$$[(168n - 84.5) \frac{a}{15p} - 12 - f + m] \text{ constituent } A\text{-hours} \quad (247)$$

from the beginning of the series.

**179.** The relation of the lengths of the constituent *A*- and constituent *B*-hours is given by the formula

$$1 \text{ constituent } A\text{-hour} = \frac{pb}{p_1 a} \text{ constituent } B\text{-hours} \quad (248)$$

The constituent *B*-hour corresponding to the integral constituent *A*-hour of formula (247) is therefore

$$[(168n - 84.5) \frac{a}{15p} - 12 - f + m] \frac{pb}{p_1 a} \text{ constituent } B\text{-hours} \quad (249)$$

from the beginning of the series.

The last formula will, in general, represent a mixed number. The integral constituent *B*-hour to which the sum for the constituent *A*-hour is to be assigned will be the nearest integral number represented by this formula. Let *g* be a fraction not greater than 0.5, which, applied either positively or negatively to the formula, will render it an integer.

**180.** The assignment of the hourly page sums for constituent *A*-hours to the constituent *B*-hours may now be represented as follows, multiples of 24 hours being rejected:

$$[(168n - 84.5) \frac{a}{15p} - 12 - f + m - \text{multiple of 24}] \text{ constituent } A\text{-hour} \quad (250)$$

sum to be assigned to

$$\left[ \left\{ (168n - 84.5) \frac{a}{15p} - 12 - f + m \right\} \frac{pb}{p_1 a} \pm g - \text{multiple of 24} \right] \text{ constituent } B\text{-hour.} \quad (251)$$

The difference between the constituent *A*-hour and the constituent *B*-hour to which the *A*-hour sum is to be assigned is

$$\left[ \left\{ (168n - 84.5) \frac{a}{15p} - 12 - f + m \right\} \left\{ \frac{pb}{p_1 a} - 1 \right\} \pm g - \text{multiple of 24} \right] \quad (252)$$

By means of the above formula table 33 has been prepared, giving the differences to be applied to the constituent *A*-hours of each page to obtain the constituent *B*-hours with which they most nearly coincide.

**181.** For the construction of secondary stencils the forms designated for the compilation of the stencil sums from the primary summations may be used. Because of the practical difficulties of constructing stencils with openings in adjacent line spaces it is desirable that the original compilation of the primary sums should be made so that each alternate line in the form for stencil sums is left vacant. As with the

primary stencils, it will generally be found convenient to use two stencils for each page of the compiled primary sums, although in some cases it may be found desirable to use more than two stencils in order to separate more clearly the groups to be summed. The actual construction of the secondary stencils is similar to that of the primary stencils. A preliminary set of forms is filled out with constituent *B*-hours as derived by differences from table 33 applied to the constituent *A*-hours. The odd and even constituent *B*-hours are then transferred to separate forms and the spaces indicated cut out. The openings corresponding to the same constituent *B*-hour are connected with ruled lines and numbered to accord with the constituent hour represented. The page numbering corresponding to the page numbering on the compiled primary sums and referring to the pages of the original tabulated hourly heights is to be entered in the column provided near the left margin of the stencil.

182. In using the stencils each sheet is to be applied to the page of compiled primary sums having the same page numbering in the left-hand column as is given on the stencil. The primary sums applying to the same constituent *B*-hour are added and the results brought together in a stencil sum form, where the totals and means are obtained. A table of divisors for obtaining the means may be readily derived as follows: In a set of stencil sum forms corresponding to those used for the compilation of constituent *A* primary sums the number of hourly heights included in each primary sum is entered in the space corresponding to that used for such primary sum. The secondary stencils for constituent *B* are then applied and the sums of the numbers obtained and compiled in the same manner as that in which the constituent *B* height sums are obtained. The divisors having been once obtained are applicable for all series of the same length.

183. In the analysis the means obtained by use of the secondary stencils may be treated as though obtained directly by the primary summations except that a special augmenting factor, to be discussed later, must be applied. The closeness of the agreement between the hourly means obtained by use of the secondary stencils and those obtained directly by use of primary stencils will depend to a large extent upon the relation of the speeds of constituents *A* and *B*. The smaller the difference in the speeds the closer will be the agreement.

184. To determine the extreme difference in the time of an individual hourly height and of the *B*-hour to which it is assigned by the secondary stencils, let an assumed case be first considered in which the tabulated heights coincide exactly with the integral *A*-hours, and that on the middle day of the page of tabulated hourly heights one of the integral *B*-hours coincides exactly with an *A*-hour. At the corresponding *A*-hour, one *A*-day later, the *B*-hour will have increased by  $24 \frac{pb}{pa}$  constituent *B*-hours. Rejecting a multiple of 24 hours, this becomes  $24\left(\frac{pb}{pa} - 1\right)$ , so that at the end of one *A*-day after the coincidence of integral hours of constituents *A* and *B* the constituent *A* hourly height will differ in time from the integral constituent *B*-hour to which it is to be assigned by  $24\left(\frac{pb}{pa} - 1\right)$  constituent *B*-hours.

At the end of the third *A*-day this difference becomes  $72\left(\frac{pb}{p_1a} - 1\right)$

constituent *B*-hours. The same difference with opposite sign will apply to the third constituent day before the middle day of the page. Now, taking account of the fact that the *B*-hour on the middle day of the page may differ by an amount as great 0.5 of a *B*-hour from the integral *A*-hour, and that the integral *A*-hour may differ as much as 0.5 of a constituent *A*, or  $0.5 pb/p_1a$  of a constituent *B* hour from the time of the actual observation of the solar hourly height, the extreme difference between the time of observation of an hourly height and the time represented by the *B*-hour with which this height is grouped by the secondary stencils may be represented by the formula

$$\pm \left[ 72\left(\frac{pb}{p_1a} - 1\right) + 0.5\left(\frac{pb}{p_1a} + 1\right) \right] \text{constituent } B\text{-hours.} \quad (253)$$

The differences may be either positive or negative, and in a long series it may reasonably be expected that the number of positive and negative values will be approximately equal.

**185.** The above formula for the extreme difference furnishes a criterion by which to judge, to some extent, the reliability of the method. Testing the following schedule of constituents for which it is proposed to use the secondary stencils, the extreme differences as indicated are obtained. The differences are expressed in constituent *B*-hours and also in constituent *B*-degrees. It will be noted that one constituent hour is equivalent to a change of  $15^\circ$  in the phase of a diurnal constituent,  $30^\circ$  in the phase of a semidiurnal constituent, etc.

Constituent <i>A</i> .....	J		S				
Constituent <i>B</i> .....	OO	2SM	K <sub>1</sub>	K <sub>2</sub>	R <sub>2</sub>	T <sub>2</sub>	P <sub>1</sub>
Difference in hours .....	3.58	1.36	1.20	1.20	1.10	1.10	1.20
Difference in degrees .....	54	41	18	36	33	33	18
Constituent <i>A</i> .....	L			2MK			
Constituent <i>B</i> .....	MS	$\lambda_2$	MK	MN	$\nu_2$	N <sub>2</sub>	
Difference in hours .....	1.09	1.18	1.43	1.24	1.26	1.45	
Difference in degrees .....	65	35	64	74	38	44	
Constituent <i>A</i> .....	O						
Constituent <i>B</i> .....	$\mu_2$	2N	$\rho_1$	Q	2Q		
Difference in hours .....	1.21	1.02	3.42	3.79	6.58		
Difference in degrees .....	36	31	51	57	99		

**186.** In the ordinary primary summation the extreme difference between the time of the observation of a solar hourly height and the integral constituent hour to which it is assigned is one-half of a constituent hour and, represented by constituent degrees, it is  $7.5^\circ$  for diurnal,  $15^\circ$  for semidiurnal,  $22.5^\circ$  for terdiurnal,  $30^\circ$  for quarter

diurnal,  $45^\circ$  for sixth-diurnal, and  $60^\circ$  for eighth-diurnal constituents. By the above schedule it will be noted that the extreme difference exceeds  $60^\circ$  in only a few cases. The largest difference is  $99^\circ$  for constituent  $2Q$  when based upon the primary summations for O. This is a small and unimportant constituent, and heretofore no analysis has been made for it, the value of its harmonic constants being inferred from those of constituent O. Although theoretically too small to justify a primary summation in general practice, the lesser work involved in the secondary summations may produce constants for this constituent which will be more satisfactory than the inferred constants.

#### FOURIER SERIES

187. A series involving only sines and cosines of whole multiples of a varying angle is generally known as the Fourier series. Such a series is of the form

$$h = H_0 + C_1 \cos \theta + C_2 \cos 2\theta + C_3 \cos 3\theta + \dots + S_1 \sin \theta + S_2 \sin 2\theta + S_3 \sin 3\theta + \dots \quad (254)$$

It can be shown that by taking a sufficient number of terms the Fourier series may be made to represent any periodic function of  $\theta$ .

This series may be written also in the following form:

$$h = H_0 + A_1 \cos(\theta + \alpha_1) + A_2 \cos(2\theta + \alpha_2) + A_3 \cos(3\theta + \alpha_3) + \dots \quad (255)$$

in which

$$A_m = [C_m^2 + S_m^2]^{\frac{1}{2}} \text{ and } \alpha_m = -\tan^{-1} \frac{S_m}{C_m}$$

$m$  being the subscript of any term.

188. From the summations for any constituent 24 hourly means are obtained, these means being the approximate heights of the constituent tide at given intervals of time. These mean constituent hourly heights, together with the intermediate heights, may be represented by the Fourier series, in which

$H_0$ =mean value of the function corresponding to the height of mean sea level above the adopted datum.

$\theta$ =an angle that changes uniformly with time and completes a cycle of  $360^\circ$  in one constituent day. The values of  $\theta$  corresponding to the 24 hourly means will be  $0^\circ, 15^\circ, 30^\circ, \dots, 330^\circ$ , and  $345^\circ$ .

Formula (254), or its equivalent (255), is the equation of a curve with the values of  $\theta$  as the abscissæ and the corresponding values of  $h$  as the ordinates. If the 24 constituent hourly means are plotted as ordinates corresponding to the values of  $0^\circ, 15^\circ, 30^\circ, \dots$  for  $\theta$ , it is possible to find values for  $H_0$ ,  $C_m$ , and  $S_m$ , which when substituted in (255) will give the equation of a curve that will pass exactly through each of the 24 points representing these means.

189. In order to make the following discussion more general, let it be assumed that the period of  $\theta$  has been divided into  $n$  equal parts, and that the ordinate or value of  $h$  pertaining to the beginning of each of those parts is known. Let  $u$  equal the interval between these ordinates, then

$$n u = 2\pi, \text{ or } 360^\circ \quad (256)$$

Let the given ordinates be  $h_0, h_1, h_2, \dots, h_{(n-1)}$  corresponding to the abscissæ  $o, u, 2u, \dots, (n-1)u$ , respectively.

It is now proposed to show that the curve represented by the following Fourier series will pass through the  $n$  points of which the ordinates are given:

$$\begin{aligned} h &= H_0 + C_1 \cos \theta + C_2 \cos 2\theta + \dots + C_k \cos k\theta \\ &\quad + S_1 \sin \theta + S_2 \sin 2\theta + \dots + S_l \sin l\theta \\ &= H_0 + \sum_{m=1}^{m=k} C_m \cos m\theta + \sum_{m=1}^{m=l} S_m \sin m\theta \end{aligned} \quad (257)$$

in which the limit  $k = \frac{n}{2}$  if  $n$  is an even number, or  $k = \frac{n-1}{2}$  if  $n$  is an odd number; and the limit  $l = \frac{n}{2} - 1$  if  $n$  is even, or  $\frac{n-1}{2}$  if  $n$  is odd.

**190.** By substituting successively the coordinates of the  $n$  given points in (257) we may obtain  $n$  equations of the form

$$h_a = H_0 + \sum_{m=1}^{m=k} C_m \cos mau + \sum_{m=1}^{m=l} S_m \sin mau \quad (258)$$

in which  $a$  represents successively the integers 0 to  $(n-1)$ .

By the solution of these  $n$  equations the values of  $n$  unknown quantities may be obtained, including  $H_0$  and the  $(n-1)$  values for  $C_m$  and  $S_m$ . It will be noted that the sum of the limits  $k$  and  $l$  of (257) or (258) equals  $(n-1)$  for both even and odd values of  $n$ .

**191.** The reason for these limits is as follows:

A continued series  $\Sigma C_m \cos m a u$  may be written

$$\begin{aligned} C_1 \cos a u + C_2 \cos 2 a u + \dots + C_n \cos n a u \\ + C_{(n+1)} \cos (n+1) a u + C_{(n+2)} \cos (n+2) a u + \dots + C_{2n} \cos 2 n a u \\ + C_{(2n+1)} \cos (2n+1) a u + C_{(2n+2)} \cos (2n+2) a u + \dots \\ + C_{3n} \cos 3 n a u \\ + \dots \end{aligned} \quad (259)$$

Since  $n u = 2\pi$  and  $a$  is an integer, the above may be written

$$\begin{aligned} &[C_1 + C_{(n+1)} + C_{(2n+1)} + \dots] \cos a u \\ &+[C_2 + C_{(n+2)} + C_{(2n+2)} + \dots] \cos 2 a u \\ &+ \dots \\ &+[C_{(n-1)} + C_{(2n-1)} + C_{(3n-1)} + \dots] \cos (n-1) a u \\ &+[C_n + C_{2n} + C_{3n} + \dots] \cos n a u \end{aligned} \quad (260)$$

Since  $\cos n a u = \cos 2a\pi = 1$ ;  $\cos (n-1) a u = \cos (2a\pi - a u) = \cos a u$ ;  $\cos (n-2) a u = \cos 2 a u$ ; etc., (260) may be written

$$\begin{aligned} &[C_n + C_{2n} + C_{3n} + \dots] \cos 0 \\ &+[C_1 + C_{(n+1)} + C_{(2n+1)} + \dots] \cos a u \\ &+[C_{(n-1)} + C_{(2n-1)} + C_{(3n-1)} + \dots] \cos a u \\ &+[C_2 + C_{(n+2)} + C_{(2n+2)} + \dots] \cos 2 a u \\ &+[C_{(n-2)} + C_{(2n-2)} + C_{(3n-2)} + \dots] \cos 2 a u \\ &+ \dots \\ &+[C_k + C_{(n+k)} + C_{(2n+k)} + \dots] \cos k a u \\ &+[C_{(n-k)} + C_{(2n-k)} + C_{(3n-k)} + \dots] \cos k a u \end{aligned} \quad (261)$$

The first term of the above is a constant which will be included with the  $H_0$  in the solution of (258). From an examination of (261) it is evident that the cosine terms will be completely represented when  $k = \frac{n}{2}$ , or  $\frac{n-1}{2}$ , according to whether  $n$  is even or odd.

Similarly, the continued series  $\Sigma S_m \sin m a u$  may be written

$$\begin{aligned}
 & [S_n + S_{2n} + S_{3n} + \dots] \sin 0 \\
 & + [S_1 + S_{(n+1)} + S_{(2n+1)} + \dots] \sin a u \\
 & - [S_{(n-1)} - S_{(2n-1)} - S_{(3n-1)} - \dots] \sin a u \\
 & + [S_2 + S_{(n+2)} + S_{(2n+2)} + \dots] \sin 2 a u \\
 & - [S_{(n-2)} - S_{(2n-2)} - S_{(3n-2)} - \dots] \sin 2 a u \\
 & + [S_l + S_{(n+l)} + S_{(2n+l)} + \dots] \sin l a u \\
 & - [S_{(n-l)} - S_{(2n-l)} - S_{(3n-l)} - \dots] \sin l a u
 \end{aligned} \quad (262)$$

The first term in the above equals zero. The remaining terms will take complete account of the series  $\sum S_m \sin m a u$ , if  $l = \frac{n}{2} - 1$  when  $n$  is even, or  $\frac{n-1}{2}$  when  $n$  is odd.

From the foregoing it is evident that the limit of  $m$  will not exceed  $\frac{n}{2}$ .

**192.** If we let  $u$  and  $\alpha$  represent any angles with fixed values,  $m$  and  $p$  any integers with fixed values, and  $a$  an integer having successive values from 0 to  $(n-1)$ , it may be shown that

$$\sum_{a=0}^{(n-1)} \sin(a m u + \alpha) = \frac{\sin \frac{1}{2} n m u}{\sin \frac{1}{2} m u} \sin [\frac{1}{2} (n-1) m u + \alpha] \quad (263)$$

$$\sum_{a=0}^{(n-1)} \cos(a m u + \alpha) = \frac{\sin \frac{1}{2} n m u}{\sin \frac{1}{2} m u} \cos [\frac{1}{2} (n-1) m u + \alpha] \quad (264)$$

$$\begin{aligned}
 \sum_{a=0}^{(n-1)} \sin a p u \sin a m u &= \frac{1}{2} \frac{\sin \frac{1}{2} n (p-m) u \cos \frac{1}{2} (n-1) (p-m) u}{\sin \frac{1}{2} (p-m) u} \\
 &\quad - \frac{1}{2} \frac{\sin \frac{1}{2} n (p+m) u \cos \frac{1}{2} (n-1) (p+m) u}{\sin \frac{1}{2} (p+m) u}
 \end{aligned} \quad (265)$$

$$\begin{aligned}
 \sum_{a=0}^{(n-1)} \cos a p u \cos a m u &= \frac{1}{2} \frac{\sin \frac{1}{2} n (p-m) u \cos \frac{1}{2} (n-1) (p-m) u}{\sin \frac{1}{2} (p-m) u} \\
 &\quad + \frac{1}{2} \frac{\sin \frac{1}{2} n (p+m) u \cos \frac{1}{2} (n-1) (p+m) u}{\sin \frac{1}{2} (p+m) u}
 \end{aligned} \quad (266)$$

$$\begin{aligned}
 \sum_{a=0}^{(n-1)} \sin a p u \cos a m u &= \frac{1}{2} \frac{\sin \frac{1}{2} n (p-m) u \sin \frac{1}{2} (n-1) (p-m) u}{\sin \frac{1}{2} (p-m) u} \\
 &\quad + \frac{1}{2} \frac{\sin \frac{1}{2} n (p+m) u \sin \frac{1}{2} (n-1) (p+m) u}{\sin \frac{1}{2} (p+m) u}
 \end{aligned} \quad (267)$$

**193.** If we let  $\alpha = 0$  and  $u = \frac{2\pi}{n}$ , or  $n u = 2\pi$ , then formulas (263) to (267) may be written as follows:

$$\sum_{a=0}^{(n-1)} \sin a m u = \frac{\sin m \pi \sin \left( m \pi - \frac{m}{n} \pi \right)}{\sin \frac{m}{n} \pi} \quad (268)$$

$$\sum_{a=0}^{(n-1)} \cos a m u = \frac{\sin m \pi \cos \left( m \pi - \frac{m}{n} \pi \right)}{\sin \frac{m}{n} \pi} \quad (269)$$

$$\sum_{a=0}^{(n-1)} \sin a p u \sin a m u = \frac{1}{2} \frac{\sin (p-m) \pi \cos \left[ (p-m) \pi - \frac{p-m}{n} \pi \right]}{\sin \frac{(p-m)}{n} \pi} - \frac{1}{2} \frac{\sin (p+m) \pi \cos \left[ (p+m) \pi - \frac{p+m}{n} \pi \right]}{\sin \frac{p+m}{n} \pi} \quad (270)$$

$$\sum_{a=0}^{(n-1)} \cos a p u \cos a m u = \frac{1}{2} \frac{\sin (p-m) \pi \cos \left[ (p-m) \pi - \frac{p-m}{n} \pi \right]}{\sin \frac{p-m}{n} \pi} + \frac{1}{2} \frac{\sin (p+m) \pi \cos \left[ (p+m) \pi - \frac{p+m}{n} \pi \right]}{\sin \frac{p+m}{n} \pi} \quad (271)$$

$$\sum_{a=0}^{(n-1)} \sin a p u \cos a m u = \frac{1}{2} \frac{\sin (p-m) \pi \sin \left[ (p-m) \pi - \frac{p-m}{n} \pi \right]}{\sin \frac{p-m}{n} \pi} + \frac{1}{2} \frac{\sin (p+m) \pi \sin \left[ (p+m) \pi - \frac{p+m}{n} \pi \right]}{\sin \frac{p+m}{n} \pi} \quad (272)$$

194. If  $p$  and  $m$  are unequal integers and neither exceeds  $\frac{n}{2}$ , the above (268) to (272) become equal to zero. Thus,

$$\left. \begin{array}{l} \sum_{a=0}^{(n-1)} \sin a m u = 0 \\ \sum_{a=0}^{(n-1)} \cos a m u = 0 \\ \sum_{a=0}^{(n-1)} \sin a p u \sin a m u = 0 \\ \sum_{a=0}^{(n-1)} \cos a p u \cos a m u = 0 \\ \sum_{a=0}^{(n-1)} \sin a p u \cos a m u = 0 \end{array} \right\} \quad (273)$$

195. If  $p$  and  $m$  are equal integers and do not exceed  $\frac{n}{2}$ , formulas (270), (271), and (272) will contain the indeterminate quantity  $\frac{\sin (p-m)\pi}{\sin \frac{p-m}{n}\pi} = \frac{0}{0}$ , and also when  $p$  and  $m$  each equal  $\frac{n}{2}$ , the indeterminate quantity  $\frac{\sin (p+m)\pi}{\sin \frac{(p+m)}{n}\pi} = \frac{0}{0}$ .

Evaluating these quantities we have

$$\left[ \frac{\sin(p-m)\pi}{\sin \frac{p-m}{n}\pi} \right]_{(p-m)=0} = \left[ \frac{\pi \cos(p-m)\pi}{\frac{\pi}{n} \cos \frac{p-m}{n}\pi} \right]_{(p-m)=0} = n \quad (274)$$

and

$$\left[ \frac{\sin(p+m)\pi}{\sin \frac{p+m}{n}\pi} \right]_{(p+m)=n} = \left[ \frac{\pi \cos(p+m)\pi}{\frac{\pi}{n} \cos \frac{p+m}{n}\pi} \right]_{(p+m)=n} = -n \quad (275)$$

In (275) it will be noted that when the integers  $p$  and  $m$  each equal  $\frac{n}{2}$ ,  $n$  must be an even number, and therefore  $\cos n\pi$  is positive, while  $\cos \pi$  is negative.

**196.** Assuming the condition that  $p$  and  $m$  are equal integers, each less than  $\frac{n}{2}$ , we have by substituting (274) in (270), (271), and (272),

$$\sum_{a=0}^{a=(n-1)} \sin a p u \sin a m u = \sum_{a=0}^{a=(n-1)} \sin^2 a m u = \frac{1}{2} n \quad (276)$$

$$\sum_{a=0}^{a=(n-1)} \cos a p u \cos a m u = \sum_{a=0}^{a=(n-1)} \cos^2 a m u = \frac{1}{2} n \quad (277)$$

$$\sum_{a=0}^{a=(n-1)} \sin a p u \cos a m u = \sum_{a=0}^{a=(n-1)} \sin a m u \cos a m u = 0 \quad (278)$$

**197.** Assuming the condition that  $p$  and  $m$  are each equal to  $\frac{n}{2}$  we have by substituting (274) and (275) in (270), (271), and (272),

$$\sum_{a=0}^{a=(n-1)} \sin^2 a m u = \frac{1}{2} n + \frac{1}{2} n \cos \pi = 0 \quad (279)$$

$$\sum_{a=0}^{a=(n-1)} \cos^2 a m u = \frac{1}{2} n - \frac{1}{2} n \cos \pi = n \quad (280)$$

$$\sum_{a=0}^{a=(n-1)} \sin a m u \cos a m u = 0 \quad (281)$$

**198.** Returning now to the solution of (258), by substituting the successive values of  $a$  from 0 to  $(n-1)$ , we have

$$\left. \begin{aligned} h_0 &= H_0 + C_1 \cos 0 + C_2 \cos 0 + \dots + C_k \cos 0 \\ &\quad + S_1 \sin 0 + S_2 \sin 0 + \dots + S_l \sin 0 \\ h_1 &= H_0 + C_1 \cos u + C_2 \cos 2u + \dots + C_k \cos ku \\ &\quad + S_1 \sin u + S_2 \sin 2u + \dots + S_l \sin lu \\ h_2 &= H_0 + C_1 \cos 2u + C_2 \cos 4u + \dots + C_k \cos 2ku \\ &\quad + S_1 \sin 2u + S_2 \sin 4u + \dots + S_l \sin 2lu \\ h_{(n-1)} &= H_0 + C_1 \cos (n-1)u + C_2 \cos 2(n-1)u + \dots \\ &\quad + C_k \cos (n-1)ku \\ &\quad + S_1 \sin (n-1)u + S_2 \sin 2(n-1)u + \dots \\ &\quad + S_l \sin (n-1)lu \end{aligned} \right\} \quad (282)$$

**199.** To obtain value of  $H_o$ , add above equations

$$\begin{aligned} & \sum_{a=0}^{a=(n-1)} h_a = n H_o \\ & + C_1 \sum_{a=0}^{a=(n-1)} \cos a u + C_2 \sum_{a=0}^{a=(n-1)} \cos 2 a u + \dots + C_k \sum_{a=0}^{a=(n-1)} \cos a k u \\ & + S_1 \sum_{a=0}^{a=(n-1)} \sin a u + S_2 \sum_{a=0}^{a=(n-1)} \sin 2 a u + \dots + S_l \sum_{a=0}^{a=(n-1)} \sin a l u \\ & = n H_o + \sum_{m=1}^{m=k} C_m \sum_{a=0}^{a=(n-1)} \cos a m u + \sum_{m=1}^{m=l} S_m \sum_{a=0}^{a=(n-1)} \sin a m u \end{aligned} \quad (283)$$

From (273),  $\sum_{a=0}^{a=(n-1)} \cos a m u$  and  $\sum_{a=0}^{a=(n-1)} \sin a m u$  each equals zero,

since neither  $k$  nor  $l$ , the maximum values of  $m$  exceeds  $\frac{n}{2}$

Therefore

$$\sum_{a=0}^{a=(n-1)} h_a = n H_o \quad (284)$$

and

$$H_o = \frac{1}{n} \sum_{a=0}^{a=(n-1)} h_a \quad (285)$$

**200.** To obtain the value of any coefficient  $C$ , such as  $C_p$ , multiply each equation of (282) by  $\cos a p u$ . Then

$$\begin{aligned} h_0 \cos 0 &= H_0 \cos 0 \\ &+ C_1 \cos 0 + C_2 \cos 0 + \dots + C_k \cos 0 \\ &+ S_1 \sin 0 + S_2 \sin 0 + \dots + S_l \sin 0 \\ h_1 \cos p u &= H_0 \cos p u \\ &+ C_1 \cos u \cos p u + C_2 \cos 2u \cos p u + \dots + C_k \cos k u \cos p u \\ &+ S_1 \sin u \cos p u + S_2 \sin 2u \cos p u + \dots + S_l \sin l u \cos p u \\ h_2 \cos 2p u &= H_0 \cos 2p u \\ &+ C_1 \cos 2u \cos 2p u + C_2 \cos 4u \cos 2p u + \dots \\ &+ C_k \cos 2k u \cos 2p u \\ &+ S_1 \sin 2u \cos 2p u + S_2 \sin 4u \cos 2p u + \dots \\ &+ S_l \sin 2l u \cos 2p u \\ \hline h_{(n-1)} \cos (n-1)p u &= H_0 \cos (n-1)p u \\ &+ C_1 \cos (n-1)u \cos (n-1)p u + C_2 \cos 2(n-1)u \cos (n-1)p u + \\ &+ C_k \cos (n-1)k u \cos (n-1)p u \\ &+ S_1 \sin (n-1)u \cos (n-1)p u + S_2 \sin 2(n-1)u \cos (n-1)p u + \dots \\ &+ S_l \sin (n-1)l u \cos (n-1)p u \end{aligned} \quad (286)$$

Summing the above equations

$$\begin{aligned} & \sum_{a=0}^{a=(n-1)} h_a \cos a p u = H_o \sum_{a=0}^{a=(n-1)} \cos a p u \\ & + C_1 \sum_{a=0}^{a=(n-1)} \cos a u \cos a p u + S_1 \sum_{a=0}^{a=(n-1)} \sin a u \cos a p u \end{aligned}$$

(Formula continued next page)

$$+ C_2 \sum_{a=0}^{a=(n-1)} \cos 2a u \cos a p u + S_2 \sum_{a=0}^{a=(n-1)} \sin 2a u \cos a p u$$

---


$$+ C_k \sum_{a=0}^{a=(n-1)} \cos a k u \cos a p u + S_l \sum_{a=0}^{a=(n-1)} \sin a l u \cos a p u$$

$$= H_0 \sum_{a=0}^{a=(n-1)} \cos a p u + \sum_{m=1}^{m=k} C_m \sum_{a=0}^{a=(n-1)} \cos a m u \cos a p u$$

$$+ \sum_{m=1}^{m=l} S_m \sum_{a=0}^{a=(n-1)} \sin a m u \cos a p u \quad (287)$$

**201.** Examining the limits of (287), it will be noted by a reference to page 63 that  $k$ , the maximum value of  $m$  for the  $C$  terms is  $\frac{n}{2}$  when  $n$  is even and  $\frac{n-1}{2}$  when  $n$  is odd; also, that  $l$  has a value of  $\frac{n}{2}-1$  when  $n$  is even and  $\frac{n-1}{2}$  when  $n$  is odd. The limits of  $p$ , which is a particular value of  $m$ , will, of course, be the same as those of  $m$ .

By (273) the quantity  $\sum_{a=0}^{a=(n-1)} \cos a p u$  becomes zero for all the values of  $p$ , and the quantity  $\sum_{a=0}^{a=(n-1)} \cos a m u \cos a p u$  becomes zero for all values of  $m$  and  $p$  except when  $p$  equals  $m$ . By (273), (278) and (281) the quantity  $\sum_{a=0}^{a=(n-1)} \sin a m u \cos a p u$  becomes zero for all values of  $m$  and  $p$ .

Formula (287) may therefore be reduced to the form

$$\sum_{a=0}^{a=(n-1)} h_a \cos a p u = C_p \sum_{a=0}^{a=(n-1)} \cos^2 a p u \quad (288)$$

For any value of  $p$  less than  $\frac{n}{2}$

$$\sum_{a=0}^{a=(n-1)} \cos^2 a p u = \frac{1}{2} n \quad (277)$$

but when  $p=\frac{n}{2}$ , this quantity becomes equal to  $n$  (280).

Therefore for all values of  $p$  less than  $\frac{n}{2}$

$$C_p = \frac{2}{n} \sum_{a=0}^{a=(n-1)} h_a \cos a p u \quad (289)$$

but when  $p$  is exactly  $\frac{n}{2}$

$$C_p = \frac{1}{n} \sum_{a=0}^{a=(n-1)} h_a \cos a p u \quad (290)$$

Since in tidal work  $p$  is always taken less than  $\frac{n}{2}$ , we are not especially concerned with the latter formula.

**202.** To obtain the value of any coefficient  $S_p$ , such as  $S_p$ , multiply each equation of (282) by  $\sin a_p u$ . Sum the resulting equations and obtain

$$\begin{aligned} \sum_{a=0}^{a=(n-1)} h_a \sin a_p u &= H_0 \sum_{a=0}^{a=(n-1)} \sin a_p u \\ &+ \sum_{m=1}^{m=k} C_m \sum_{a=0}^{a=(n-1)} \cos a_m u \sin a_p u \\ &+ \sum_{m=1}^{m=l} S_m \sum_{a=0}^{a=(n-1)} \sin a_m u \sin a_p u \end{aligned} \quad (291)$$

By (273), (278), and (281) the quantities  $\sum_{a=0}^{a=(n-1)} \sin a_p u$  and  $\sum_{a=0}^{a=(n-1)} \cos a_m u \sin a_p u$  are zero for all the values of  $m$  and  $p$ ; and  $\sum_{a=0}^{a=(n-1)} \sin a_m u \sin a_p u$  becomes zero for all the values of  $m$  and  $p$  except when  $m$  and  $p$  are equal. In this case the limit of  $l$  for  $m$  and  $p$  is less than  $\frac{n}{2}$  and by (276), the quantity  $\sum_{a=0}^{a=(n-1)} \sin^2 a_p u = \frac{1}{2} n$ . Therefore, formula (291) reduces to the form

$$\sum_{a=0}^{a=(n-1)} h_a \sin a_p u = \frac{1}{2} n S_p \quad (292)$$

and

$$S_p = \frac{2}{n} \sum_{a=0}^{a=(n-1)} h_a \sin a_p u \quad (293)$$

**203.** By substituting (285), (289), (290), and (293) in (257), the following equation of a curve, which will pass through the  $n$  given points, will be obtained

$$\begin{aligned} h &= \frac{1}{n} \sum_{a=0}^{a=(n-1)} h_a + \left[ \frac{2}{n} \sum_{a=0}^{a=(n-1)} h_a \cos a u \right] \cos \theta \\ &+ \left[ \frac{2}{n} \sum_{a=0}^{a=(n-1)} h_a \sin a u \right] \sin \theta \\ &+ \left[ \frac{2}{n} \sum_{a=0}^{a=(n-1)} h_a \cos 2 a u \right] \cos 2 \theta \\ &+ \left[ \frac{2}{n} \sum_{a=0}^{a=(n-1)} h_a \sin 2 a u \right] \sin 2 \theta \\ &\dots \\ &+ \left[ \frac{2}{n} \sum_{a=0}^{a=(n-1)} h_a \cos k a u \right] \cos k \theta \\ &+ \left[ \frac{2}{n} \sum_{a=0}^{a=(n-1)} h_a \sin l a u \right] \sin l \theta \end{aligned} \quad (294)$$

\*If  $n$  is even and  $k = \frac{n}{2}$ , this fraction is  $\frac{1}{n}$  instead of  $\frac{2}{n}$ .

**204.** Although by taking a sufficient number of terms the Fourier series may thus be made to represent a curve which will be exactly satisfied by the  $n$  given ordinates, this is, in general, neither necessary nor desirable in tidal work, since it is known that the mean ordinates obtained from the summations of the hourly heights of the tide include many irregularities due to the imperfect elimination of the meteorological effects and also residual effects of constituents having periods incommensurable with that of the constituent sought. It is desirable to include only the terms of the series which represent the true periodic elements of the constituent. With series of observations of sufficient length, the coefficient of the other terms, if sought, will be found to approximate to zero.

**205.** The short-period constituents as derived from the equilibrium theory are, in general, either diurnal or semidiurnal. If the period of  $\theta$  in formula (257) is taken to correspond to the constituent day, the diurnal constituents will be represented by the terms with coefficient  $C_1$  and  $S_1$ , and the semidiurnal constituents by the terms with coefficients  $C_2$  and  $S_2$ . For the long-period constituents, the period of  $\theta$  may be taken to correspond to the constituent month or to the constituent year, in which case the coefficients  $C_1$  and  $S_1$  will refer to the monthly or annual constituents and the coefficients  $C_2$  and  $S_2$  to the semimonthly or semiannual constituents. For most of the constituents the coefficients  $C_1$ ,  $S_1$ ,  $C_2$ , and  $S_2$  will be the only ones required, but for the tides depending upon the fourth power of the moon's parallax and for the overtides and the compound tides, other coefficients will be required. Terms beyond those with coefficients  $C_8$  and  $S_8$ , for the overtides of the principal lunar constituent are not generally used in tidal work.

**206.** When it is known that certain periodic elements exist in a constituent tide and that the mean ordinates obtained from observations include accidental errors that are not periodic, it may be readily shown by the method known as the least square adjustment, using the observational equations represented by (258), that the most probable values of the constant  $H_o$  and the coefficients  $C_p$  and  $S_p$  are the same as those given by formulas (285), (289), and (293), respectively.

**207.** Since in tidal work the value of  $H_o$ , which is the elevation of mean sea level above the datum of observations, is generally determined directly from the original tabulation of hourly heights, formula (285) is unnecessary except for checking purposes. Formulas (289) and (293) are used for obtaining the most probable values of the coefficients  $C_p$  and  $S_p$  from the hourly means obtained from the summations.

**208.** When 24 hourly means are used  $n=24$  and  $u=15^\circ$ , and the formulas may be written

$$C_p = \frac{1}{12} \sum_{a=0}^{a=23} h_a \cos 15 a p \quad (295)$$

$$S_p = \frac{1}{12} \sum_{a=0}^{a=23} h_a \sin 15 a p \quad (296)$$

in which the angles are expressed in degrees.

If only 12 means are used, the formulas become

$$C_p = \frac{1}{6} \sum_{a=0}^{a=11} h_a \cos 30 a p \quad (297)$$

$$S_p = \frac{1}{6} \sum_{a=0}^{a=11} h_a \sin 30^\circ a p \quad (298)$$

**209.** The upper part of Form 194 (fig. 16) is designed for the computation of the coefficients  $C_p$  and  $S_p$  in accordance with formulas (295) and (296) to take account of the 24 constituent hourly means.

It is now desired to express each constituent in the form

$$y = A \cos(p\theta + \alpha) \quad (299)$$

or using a more specialized notation by

$$y = A \cos(p\theta - \xi) \quad (300)$$

By trigonometry

$$\begin{aligned} A \cos(p\theta - \xi) &= A \cos \xi \cos p\theta + A \sin \xi \sin p\theta \\ &= C_p \cos p\theta + S_p \sin p\theta \end{aligned} \quad (301)$$

$$\text{in which } C_p = A \cos \xi \quad \text{and} \quad S_p = A \sin \xi \quad (302)$$

Therefore,

$$\tan \xi = \frac{S_p}{C_p} \quad (303)$$

and

$$A = \frac{C_p}{\cos \xi} = \frac{S_p}{\sin \xi} = \sqrt{C_p^2 + S_p^2} \quad (304)$$

Substituting in formulas (303) and (304) the values of  $C_p$  and  $S_p$  from formulas (295) and (296), the corresponding values for  $A$  and  $\xi$  may be obtained. Substituted in formula (300), these furnish an approximate representation of one of the tidal constituents, but a further processing is necessary in order to obtain the mean amplitude and epoch of the constituent.

#### AUGMENTING FACTORS

**210.** In the usual summations with the primary stencils for all the short period constituents, except constituent  $S$ , the hourly ordinates which are summed in any single group are scattered more or less uniformly over a period from one-half of a constituent hour before to one-half of a constituent hour after the exact constituent hour which the group represents. Because of this the resulting mean will differ a little from the true mean ordinate that would be obtained if all the ordinates included were read on the exact constituent hour, as with constituent  $S$ , and the amplitude obtained will be less than the true amplitude of the constituent. The factor necessary to take account of this fact is called the augmenting factor.

**211.** Let any constituent be represented by the curve

$$y = A \cos(at + \alpha) \quad (305)$$

in which

$A$ =the true amplitude of the constituent

$a$ =the speed of the constituent (degrees per solar hours)

$t$ =variable time (expressed in solar hours)

$\alpha$ =any constant.

The mean value of  $y$  for a group of consecutive ordinates from  $\tau/2$  hours before to  $\tau/2$  hours after any given time  $t$ ,  $\tau$  being the number of solar hours covered by the group, is

$$\begin{aligned} \frac{A}{\tau} \int_{t-\tau/2}^{t+\tau/2} \cos(at + \alpha) dt &= \left[ \frac{180}{\pi} \frac{A}{a\tau} \sin(at + \alpha) \right]_{t-\tau/2}^{t+\tau/2} \\ &= \frac{180}{\pi} \frac{A}{a\tau} \left[ \sin\left(at + \alpha + \frac{a\tau}{2}\right) - \sin\left(at + \alpha - \frac{a\tau}{2}\right) \right] \\ &= \frac{360}{\pi} \frac{A}{a\tau} \cos(at + \alpha) \sin \frac{a\tau}{2} = \frac{360}{\pi a\tau} \sin \frac{a\tau}{2} A \cos(at + \alpha) \end{aligned} \quad (306)$$

**212.** Since the true value of  $y$  at any time  $t$ , is equal to  $A \cos(at + \alpha)$  by (305), it is evident that the relation of this true value to the mean value (306) for the group  $\tau$  hours in length is

$$\frac{A \cos(at + \alpha)}{\frac{360}{\pi a\tau} \sin \frac{a\tau}{2} A \cos(at + \alpha)} = \frac{\pi a\tau}{360 \sin \frac{a\tau}{2}} \quad (307)$$

The quantity  $\frac{\pi a\tau}{360 \sin \frac{a\tau}{2}}$  is the augmenting factor which is to be

applied to the mean ordinate to obtain the true ordinate. In the use of this factor it is assumed that all the consecutive ordinates within the time  $\tau/2$  hours before to  $\tau/2$  hours after the given time have been used in obtaining the mean. This assumption is, of course, only approximately realized in the summation for any constituent, but the longer the series of observations the more nearly to the truth it approaches.

**213.** According to the usual summations with the primary stencils, the hourly heights included in a single group may be distributed over an interval from one-half of a constituent hour before to one-half of a constituent hour after the hour to be represented. In this case  $\tau$  equals one constituent hour, or  $\frac{15p}{a}$  solar hours.

Substituting this in (307), the

$$\text{augmenting factor} = \frac{\pi p}{24 \sin \frac{15p}{2}} \quad (308)$$

which is the formula generally adopted for the short-period constituents and is the one used in the calculation of the augmenting factors in Form 194. For the long-period constituents special factors are necessary which will be explained later.

**214.** If the second system of distribution of the hourly heights as described on page 53 is adopted,  $\tau$  equals one solar hour and formula (307) becomes

$$\text{augmenting factor} = \frac{\pi a}{360 \sin \frac{a}{2}} \quad (309)$$

It will be noted that formula (308) depends upon the value of  $p$  and therefore will be the same for all short period constituents (S excepted) with like subscripts. Formula (309) depends upon the speed  $a$  of the constituent and will therefore be different for each constituent.

**215.** When the secondary stencils are used, the grouping of the ordinates is less simple than that provided by the primary stencils only. Let it be assumed that the series is of sufficient length so that the distribution of the ordinates is more or less uniform in accordance with the system adopted.

Suppose the original primary summations have been made for constituent  $A$  with speed  $a$  and that the secondary stencils have been used for constituent  $B$  with speed  $b$ . Then let  $p$  and  $p'$  represent the subscripts of constituents  $A$  and  $B$ , respectively.

The equation for constituent  $B$  may be written

$$y = B \cos(bt + \beta) \quad (310)$$

**216.** In the primary summation for constituent  $A$ , the group of ordinates included in a single sum covers a period of one constituent  $A$  hour or  $\frac{15p}{a}$  solar hours. Expressed in time  $t$ , midway of this interval and representing the exact integral constituent  $A$  hour to which the group applied, the average value of the  $B$  ordinates included in such a group may be written

$$\begin{aligned} & \frac{a}{15p} B \int_{t-\frac{15p}{a}}^{t+\frac{15p}{a}} \cos(bt + \beta) dt \\ &= \frac{180}{\pi} \frac{a}{15pb} B \left[ \sin\left(bt + \beta + \frac{15pb}{2a}\right) - \sin\left(bt + \beta - \frac{15pb}{2a}\right) \right] \\ &= \left( \frac{24}{\pi} \frac{a}{pb} \sin \frac{15pb}{2a} \right) B \cos(bt + \beta) \\ &= F_1 B \cos(bt + \beta) \end{aligned} \quad (311)$$

In which  $F_1$ , for brevity, is substituted for the coefficient  $\frac{24}{\pi} \frac{a}{pb} \sin \frac{15pb}{2a}$  and gives the relation of the average  $B$  ordinate included in the  $A$  grouping to the true  $B$  ordinate for the time  $t$  represented by that group. The reciprocal of this coefficient will be that part of the augmenting factor necessary to take account of this primary grouping. If the primary summing has been for the constituent S, this coefficient may be taken as unity since the original S sums refer to the exact S hour.

**217.** When the secondary stencils are applied to the constituent  $A$  group sums, the groups applying to an exact constituent  $A$  hour at any time  $t$  and represented by that time, will be distributed over an interval of a constituent  $B$  hour, or  $\frac{15p'}{b}$  solar hours.

For an integral constituent  $B$  hour at any time  $t$  within the middle day represented by a seven-day page of original tabulations the limits of this interval will be  $\left(t - \frac{15p'}{2b}\right)$  and  $\left(t + \frac{15p'}{2b}\right)$ . For the same page

of tabulations, letting  $t$  represent the same time in the middle day, the limits of the group interval for the day following the middle one, are  $\left(t + \frac{360p}{a} - \frac{15p'}{2b}\right)$  and  $\left(t + \frac{360p}{a} + \frac{15p'}{2b}\right)$ . If we let  $n = -3, -2, -1, 0, +1, +2, +3$ , respectively, for the seven successive days represented by a single page of original tabulations, the limits of the group interval for any day of the page may be represented by

$$\left(t + \frac{360pn}{a} - \frac{15p'}{2b}\right) \text{ and } \left(t + \frac{360pn}{a} + \frac{15p'}{2b}\right)$$

**218.** Formula (311) gives the mean value of the  $B$  ordinate for grouping of the  $A$  summations. The mean value of (311) obtained by combining the groups falling in any particular day of page of tabulations in the limits indicated above is

$$\begin{aligned} & \frac{b}{15p'} F_1 B \int_{t + \frac{360pn}{a} - \frac{15p'}{2b}}^{t + \frac{360pn}{a} + \frac{15p'}{2b}} \cos(bt + \beta) dt \\ &= \frac{180}{\pi} \frac{1}{15p'} F_1 B \left[ \sin\left(bt + \beta + \frac{360bpn}{a} + \frac{15p'}{2}\right) \right. \\ &\quad \left. - \sin\left(bt + \beta + \frac{360bpn}{a} - \frac{15p'}{2}\right) \right] \\ &= \left( \frac{24}{\pi} \frac{1}{p'} \sin \frac{15p'}{2} \right) F_1 B \cos\left(bt + \beta + \frac{360pn}{a}\right) \\ &= F_1 F_2 B \cos\left(bt + \beta + \frac{360bpn}{a}\right) \end{aligned} \quad (312)$$

if we put  $F_2 = \frac{24}{\pi} \frac{1}{p'} \sin \frac{15p'}{2}$  for brevity.

**219.** Formula (312) represents the mean value of the  $B$  ordinate for a particular day of the page record. The average value for the 7 days may be written

$$\begin{aligned} & \frac{1}{7} F_1 F_2 B \sum_{n=-3}^{n=+3} \cos\left(t + \beta + \frac{360bpn}{a}\right) \\ &= \frac{1}{7} F_1 F_2 B \left[ \cos(bt + \beta) \cos\left(-3 \frac{360bp}{a}\right) - \sin(bt + \beta) \sin\left(-3 \frac{360bp}{a}\right) \right. \\ &\quad + \cos(bt + \beta) \cos\left(-2 \frac{360bp}{a}\right) - \sin(bt + \beta) \sin\left(-2 \frac{360bp}{a}\right) \\ &\quad + \cos(bt + \beta) \cos\left(-1 \frac{360bp}{a}\right) - \sin(bt + \beta) \sin\left(-1 \frac{360bp}{a}\right) \\ &\quad + \cos(bt + \beta) \cos 0 - \sin(bt + \beta) \sin 0 \\ &\quad \left. + \cos(bt + \beta) \cos\left(\frac{360bp}{a}\right) - \sin(bt + \beta) \sin\left(\frac{360bp}{a}\right) \right] \end{aligned}$$

(Formula continued next page)

$$\begin{aligned}
& + \cos(bt+\beta) \cos\left(2 \frac{360bp}{a}\right) - \sin(bt+\beta) \sin\left(2 \frac{360bp}{a}\right) \\
& + \cos(bt+\beta) \cos\left(3 \frac{360bp}{a}\right) - \sin(bt+\beta) \sin\left(3 \frac{360bp}{a}\right) \Big] \\
& = \frac{1}{7} F_1 F_2 B \left[ 1 + 2 \cos \frac{360bp}{a} + 2 \cos 2 \frac{360bp}{a} + 2 \cos 3 \frac{360bp}{a} \right] \cos(bt+\beta) \\
& = \frac{1}{7} F_1 F_2 B \left[ 2 \frac{\sin 2 \frac{360bp}{a} \cos \frac{3}{2} \frac{360bp}{a}}{\sin \frac{360bp}{a}} - 1 \right] \cos(bt+\beta) \\
& = \frac{1}{7} F_1 F_2 B \left[ \frac{\sin \frac{1260bp}{a}}{\sin \frac{180bp}{a}} \right] \cos(bt+\beta). \tag{313}
\end{aligned}$$

**220.** Replacing the equivalents of  $F_1$  and  $F_2$  in (313), the average value of the  $B$  ordinate as obtained by the secondary summations may be written

$$\left[ \frac{24a}{\pi pb} \sin \frac{15bp}{2a} \right] \left[ \frac{24}{\pi p'} \sin \frac{15p'}{2} \right] \left[ \frac{\sin \frac{1260bp}{a}}{7 \sin \frac{180bp}{a}} \right] B \cos(bt+\beta) \tag{314}$$

Since the true ordinate of constituent  $B$  at any time  $t$  is equal to  $B \cos(bt+\beta)$ , the reciprocal of the bracketed coefficient will be the augmenting factor necessary to reduce the  $B$  ordinate as obtained from the summations to their true values.

This augmenting factor may be written

$$\left[ \frac{\pi bp}{24a \sin \frac{15bp}{2a}} \right] \left[ \frac{\pi p'}{24 \sin \frac{15p'}{2}} \right] \left[ \frac{7 \sin \frac{180bp}{a}}{\sin \frac{1260bp}{a}} \right] \tag{315}$$

The first factor of the above is to be omitted if the primary summations are for constituent S. It will be noted that the middle factor is the same as the augmenting factor that would be used if constituent  $B$  had been subjected to the primary summations.

#### PHASE LAG OR EPOCH

**221.** The phase lag or epoch of a tidal constituent, which is represented by the Greek kappa ( $\kappa$ ), is the difference between the phase of the observed constituent and the phase of its argument at the same time. This difference remains approximately constant for any constituent in a particular locality. The phase of a constituent argument for any time may be obtained from the argument formula in table 2 by making suitable substitutions for the astronomical elements. The argument itself is represented by the general symbol  $(V+u)$  or  $E$  and

its phase or value pertaining to an initial instant of time, such as the beginning of a series of observations, is expressed by  $(V_o + u)$ . Referring to formula (300), since  $\theta$  is reckoned from the beginning of the series, the angular quantity  $(-\xi)$  is the corresponding phase of the observed constituent at this time. The phase lag may therefore be expressed by the following general formula:

$$\kappa = V_o + u - (-\xi) = V_o + u + \xi \quad (316)$$

**222.** Since the argument formulas of all short-period constituents contain some multiple of the hour angle ( $T$ ) of the mean sun, the arguments themselves will have different values in different longitudes at the same instant of time. If  $p$  equals the coefficient of  $T$  or the subscript of the constituent and  $L$  equals the longitude of the place in degrees reckoned west from Greenwich,  $L$  being considered as negative for east longitude, the relation between the local and Greenwich argument for any constituent may be expressed as follows:

$$\text{local } (V + u) = \text{Greenwich } (V + u) - pL \quad (317)$$

**223.** Also, since the absolute time of the beginning of a day or the beginning of a year depends upon the time meridian used in the locality, the initial instant taken for the beginning of a series of observations may differ in different localities even though expressed in the same clock time of the same calendar day. If we let  $S$  equal the longitude of the time meridian in degrees, positive for west and negative for east, the same meridian expressed in hours becomes  $S/15$ . Letting  $a$  equal the speed or hourly rate of change in the constituent argument, the difference in argument due to the difference in the absolute beginning of the series becomes  $aS/15$ , and the relation between the local and Greenwich argument due to this difference may be expressed as follows:

$$\text{local } (V_o + u) = \text{Greenwich } (V_o + u) - pL + aS/15 \quad (318)$$

In the above formula the local and Greenwich  $(V_o + u)$  pertain to the same clock time but not the same absolute time unless both clocks are set for the meridian of Greenwich.

**224.** Values of  $(V_o + u)$  for the meridian of Greenwich at the beginning of each calendar year 1850 to 2000 are given in table 15 for all constituents represented in the Coast and Geodetic Survey tide-predicting machine. Tables 16 to 18 provide differences for referring the arguments to other days and hours of the year. In the preparation of table 15 that portion of the argument included in the  $u$  was treated as a constant with a value pertaining to the middle of the calendar year. If the Greenwich  $(V_o + u)$  with its corrections is substituted for the local  $(V_o + u)$  in formula (316), we obtain

$$\kappa = \text{Greenwich } (V_o + u) - pL + aS/15 + \xi \quad (319)$$

**225.** The phase lag designated by  $\kappa$  is sometimes called the local epoch to distinguish it from certain modified forms which may be used for special purposes. In the preparation of the harmonic constants for predictions it is convenient to combine the longitude and time meridian corrections with the local epoch to form a modified epoch

designated by  $k'$  or by the small  $g$ . The relation of the modified epoch to the local epoch may then be expressed by the following formula:

$$k' \text{ or } g = \kappa + pL - aS/15 = \text{Greenwich } (V_o + u) + \xi \quad (320)$$

**226.** The phases of the same tidal constituent in different parts of the world are not directly comparable through their local epochs since these involve the longitude of the locality. For such a comparison it is desirable to have a Greenwich epoch that is independent of both longitude and time meridian. Such an epoch may be designated by the capital  $G$  and its relation to the corresponding local epoch expressed as follows:

$$\text{Greenwich epoch } (G) = \kappa + pL = \text{Greenwich } (V_o + u) + aS/15 + \xi \quad (321)$$

**227.** The angle  $\kappa$  may be graphically represented by figures 7 and 8. In figure 7, we have a simple representation of a single con-




FIGURE 8.

stituent. In this figure changes in the phase or angle are measured along the horizontal line, positive change toward the right and negative change toward the left. The full vertical line indicates the beginning of the series, at which time the angle  $p\theta$ , or  $at$ , equals 0. At the left of this vertical line, the symbol of a moon ( $M$ ) indicates the zero value of the equilibrium argument that precedes the beginning of the series. For the principal lunar or solar constituent, this will be simultaneous with a transit of the mean moon (modified by longitude of moon's node) or of the mean sun, and for other short-period constituents with the transit of a fictitious star representing such constituent (p. 23). At the point represented by this moon, the angle  $(V+u)$  has a value of zero. This angle increases to the right, and at the beginning of the series has a value represented by  $(V_o+u)$ , which may be readily computed for the beginning of any series. This interval from  $M$  to the time of occurrence of the first following constituent high water is the epoch  $\kappa$ . This represents the lag or difference between the actual constituent high water at any

place and the theoretical time as determined by the equilibrium theory. The distance from the beginning of the series to the following high water is the  $\zeta$  of formula (300), which is determined directly from the analysis of the observations. From the figure it is evident that the  $\kappa$  is the sum of  $(V_o + u)$  and  $\zeta$ , and also that it is independent of the time of the beginning of the series.

**228.** Figure 8 gives a more detailed representation of the epoch of a constituent. In this figure the horizontal line represents changes in time. Distances along this line will be proportional to the changes in the angle of any single constituent, but since each constituent has a different speed equal distances along this line will not represent equal angles for different constituents. The time between the events may be converted into an equivalent constituent angle by multiplying by the speed of the constituent. The figure is to some extent self-explanatory. The word "transit" signifies the transit of the fictitious moon representing any constituent and also the time when the equilibrium argument of that constituent has a zero value. For all short-period constituents the time of such zero value will depend upon the longitude of the place of observation as well as upon absolute time. For long-period constituents the zero values are independent of the longitude of the place of observation, and the "transits" over the several meridians may be considered as occurring simultaneously, which is equivalent to taking the coefficient  $p$  equal to zero. The figure illustrates the relation between the Greenwich  $(V_o + u)$  calculated for the meridian of Greenwich and referring to standard Greenwich time and local  $(V_o + u)$  referring to the meridian of observation and the actual time of the beginning of the observations.

#### INFERENCE OF CONSTANTS

**229.** Under the conditions assumed for the equilibrium theory the amplitudes of the constituents could be computed directly by means of the coefficient formulas without the necessity of securing tidal observations, and the phases would correspond with the equilibrium arguments of the constituents. Under the conditions that actually exist it has been found from observations that the amplitudes of the constituents of a similar type at any place, although differing greatly from their theoretical values, have a relation that, in general, agrees fairly closely with the relations of their theoretical coefficients. It has also been ascertained from the results obtained from observations that the difference in the epochs or lags of the constituents have a relation conforming, in general, with the relation of the differences in their speeds. This last relation is based upon an assumption that the ages of the inequalities due to the disturbing influence of other constituents of a similar type are equal when expressed in time.

**230.** If the mean amplitudes, epochs, and speeds of several constituents  $A$ ,  $B$ ,  $C$ , are represented by  $H(A)$ ,  $H(B)$ ,  $H(C)$ ,  $\kappa(A)$ ,  $\kappa(B)$ ,  $\kappa(C)$ , and  $a$ ,  $b$ ,  $c$ , respectively, the above relations may be expressed by the following formulas:

$$H(B) = \frac{\text{mean coefficient of } B}{\text{mean coefficient of } A} H(A) \quad (322)$$

$$\kappa(C) - \kappa(A) = \frac{c-a}{b-a} [\kappa(B) - \kappa(A)] \quad (323)$$

or,

$$\kappa(C) = \kappa(A) + \frac{c-a}{b-a} [\kappa(B) - \kappa(A)] \quad (324)$$

By formula (322) the amplitude of a constituent ( $B$ ) may be inferred from the known amplitude of a constituent ( $A$ ), and by formula (324) the epoch of a constituent ( $C$ ) may be inferred from the known epochs of constituents ( $A$ ) and ( $B$ ).

**231.** These formulas have, however, certain limitations. They are not applicable to shallow water and meteorological constituents, nor are they adapted to the determination of a diurnal constituent from a semidiurnal constituent or of a semidiurnal constituent from a diurnal constituent. The results obtained by the application of the formulas to tides of similar type may be considered only as rough approximations to the truth. They may, however, be preferable to the values obtained for certain constituents when the series of observations is short.

**232.** By substituting the mean values of the coefficients and the speeds from table 2 the following special formulas may be derived from the general formulas (322) and (324)

### Diurnal constituents

$$H(J_1) = 0.079 H(O_1); \quad \kappa(J_1) = \kappa(K_1) + 0.496 [\kappa(K_1) - \kappa(O_1)] \quad (325)$$

$$H(M_1) = 0.071 H(O_1); \quad \kappa(M_1) = \kappa(K_1) - 0.500 [\kappa(K_1) - \kappa(O_1)] \quad (326)$$

$$H(OO) = 0.043 H(O_1); \quad \kappa(OO) = \kappa(K_1) + 1.000 [\kappa(K_1) - \kappa(O_1)] \quad (327)$$

$$H(P_1) = 0.331 H(K_1); \quad \kappa(P_1) = \kappa(K_1) - 0.075 [\kappa(K_1) - \kappa(O_1)] \quad (328)$$

$$H(Q_1) = 0.194 H(O_1); \quad \kappa(Q_1) = \kappa(K_1) - 1.496 [\kappa(K_1) - \kappa(O_1)] \quad (329)$$

$$H(2Q) = 0.026 H(O_1); \quad \kappa(2Q) = \kappa(K_1) - 1.992 [\kappa(K_1) - \kappa(O_1)] \quad (330)$$

$$H(\rho_1) = 0.038 H(O_1); \quad \kappa(\rho_1) = \kappa(K_1) - 1.429 [\kappa(K_1) - \kappa(O_1)] \quad (331)$$

### Semidiurnal constituents

$$H(K_2) = 0.272 H(S_2); \quad \kappa(K_2) = \kappa(S_2) + 0.081 [\kappa(S_2) - \kappa(M_2)] \quad (332)$$

$$H(L_2) = 0.028 H(M_2); \quad \kappa(L_2) = \kappa(S_2) - 0.464 [\kappa(S_2) - \kappa(M_2)] \quad (333)$$

$$= 0.143 H(N_2); \quad \kappa(M_2) = \kappa(S_2) + 1.000 [\kappa(S_2) - \kappa(N_2)] \quad (334)$$

$$H(N_2) = 0.194 H(M_2); \quad \kappa(N_2) = \kappa(S_2) - 1.536 [\kappa(S_2) - \kappa(M_2)] \quad (335)$$

$$H(2N) = 0.026 H(M_2); \quad \kappa(2N) = \kappa(S_2) - 2.072 [\kappa(S_2) - \kappa(M_2)] \quad (336)$$

$$= 0.133 H(N_2); \quad \kappa(M_2) = \kappa(S_2) - 2.000 [\kappa(S_2) - \kappa(N_2)] \quad (337)$$

$$H(R_2) = 0.008 H(S_2); \quad \kappa(R_2) = \kappa(S_2) + 0.040 [\kappa(S_2) - \kappa(M_2)] \quad (338)$$

$$H(T_2) = 0.059 H(S_2); \quad \kappa(T_2) = \kappa(S_2) - 0.040 [\kappa(S_2) - \kappa(M_2)] \quad (339)$$

$$H(\lambda_2) = 0.007 H(M_2); \quad \kappa(\lambda_2) = \kappa(S_2) - 0.536 [\kappa(S_2) - \kappa(M_2)] \quad (340)$$

$$H(\mu_2) = 0.024 H(M_2); \quad \kappa(\mu_2) = \kappa(S_2) - 2.000 [\kappa(S_2) - \kappa(M_2)] \quad (341)$$

$$H(\nu_2) = 0.038 H(M_2); \quad \kappa(\nu_2) = \kappa(S_2) - 1.464 [\kappa(S_2) - \kappa(M_2)] \quad (342)$$

$$= 0.194 H(N_2); \quad \kappa(M_2) = \kappa(S_2) - 0.866 [\kappa(S_2) - \kappa(N_2)] \quad (343)$$

**233.** In order to test the reliability of the results obtained by inference as above, 60 stations representing various types of tide in different parts of the world where the harmonic constants had been determined from observations were selected and a comparison was made between the values for certain constants as obtained by inference and by observations. The tests were applied to the diurnal constituents

$M_1$ ,  $P_1$ , and  $Q_1$ , and to the semidiurnal constituents  $K_2$ ,  $L_2$ , and  $\nu_2$ , and formulas (326), (328), (329), (332), (333), and (342) were used for the purpose. The following results were obtained for the differences between values as obtained from inference and from observations. The average gross difference is the average difference without regard to the signs of the individual items, and the average net difference takes into account these signs so that a positive difference may offset a negative difference in the mean. The last two lines in the table show the percentage of cases in which the differences were less than 0.05 and 0.10 foot, respectively, for the amplitudes, and less than  $10^\circ$  and  $20^\circ$ , respectively, for the epochs.

	$M_1$ ampli- tude	$M_1$ epoch	$P_1$ ampli- tude	$P_1$ epoch	$Q_1$ ampli- tude	$Q_1$ epoch
Maximum difference.....	Ft. 0.05	Deg. 149	Ft. 0.27	Deg. .03	Ft. 0.05	Deg. 105
Average gross difference.....	.02	31	.01	8	.01	14
Average net difference.....	.01	1	.01	3	.00	0
Differences less than 0.05 foot or $10^\circ$ .....	% 93	% 37	% 85	% 76	% 96	% 58
Differences less than 0.10 foot or $20^\circ$ .....	100	57	92	92	100	82

	$K_2$ ampli- tude	$K_2$ epoch	$L_2$ ampli- tude	$L_2$ epoch	$\nu_2$ ampli- tude	$\nu_2$ epoch
Maximum difference.....	Ft. 0.28	Deg. 51	Ft. 1.09	Deg. 104	Ft. 0.28	Deg. 53
Average gross difference.....	.02	9	.09	25	.04	14
Average net difference.....	.00	5	.08	4	.02	4
Differences less than 0.05 foot or $10^\circ$ .....	% 87	% 65	% 58	% 20	% 71	% 48
Differences less than 0.10 foot or $20^\circ$ .....	97	93	78	44	88	83

By using formulas (334) and (343) for  $L_2$  and  $\nu_2$  the results are slightly improved, the average net differences for the amplitude and epoch of  $L_2$  becoming 0.07 foot and  $3^\circ$ , respectively, the difference for the epoch of  $\nu_2$  becoming  $2^\circ$ , while the average net difference for the amplitude of  $\nu_2$  remains unchanged.

**234.** Although there is a fairly good agreement indicated by the average differences, it is evident that the inferred constants, especially the epochs, cannot be depended upon for a high degree of refinement. It may be stated, however, that for constituents with very small amplitudes the epochs determined from actual observations may be equally unreliable. This becomes evident when results from different years of observations are compared. Fortunately, the large discrepancies in epochs are found only in constituents of small amplitude and are therefore of little practical importance.

**235.** Constituent  $\mu_2$  as determined by inference is relatively unimportant. However, this constituent has the same period as the compound tide  $2MS_2$  and when obtained directly from the analysis of observations frequently differs considerably from the inferred  $\mu_2$  both in amplitude and epoch. The inferred values for this constituent cannot therefore be considered as very satisfactory.

**236.** Prior to the elimination process described in the next section, certain preliminary corrections are applied to the amplitudes and

epochs of constituents  $S_2$  and  $K_1$  because of the disturbing effects of  $K_2$  and  $T_2$  on the former and  $P_1$  on the latter. In a short series of observations these effects may be considerable because of the small differences in the periods of the constituents involved.

**237.** Let

$$y_1 = A \cos (at + \alpha) \quad (344)$$

and

$$y_2 = B \cos (bt + \beta) \quad (345)$$

represent two constituents, the first being the principal or predominating constituent and the latter a secondary constituent whose effect is to modify the amplitude and epoch of the principal constituent. The resultant tide will then be represented by

$$y = y_1 + y_2 = A \cos (at + \alpha) + B \cos (bt + \beta) \quad (346)$$

Values of  $t$  which will render (344) a maximum must satisfy the derived equation

$$Aa \sin (at + \alpha) = 0 \quad (347)$$

and the values of  $t$  which will render (346) a maximum must satisfy the equation

$$Aa \sin (at + \alpha) + Bb \sin (bt + \beta) = 0 \quad (348)$$

For a maximum of (344)

$$t = \frac{2n\pi - \alpha}{a} \quad (349)$$

in which  $n$  is any integer.

**238.** Let  $\frac{\theta}{a}$  = the acceleration in the principal constituent  $A$  due to the disturbing constituent  $B$ . Then for a maximum of (346)

$$t = \frac{2n\pi - \alpha - \theta}{a} \quad (350)$$

This value of  $t$  must satisfy equation (348), therefore we have

$$\begin{aligned} & Aa \sin (2n\pi - \theta) + Bb \sin \left[ \frac{b}{a}(2n\pi - \theta - \alpha) + \beta \right] \\ &= -Aa \sin \theta + Bb \sin \left[ \frac{b-a}{a}(2n\pi - \theta - \alpha) + \beta - \alpha - \theta \right] = 0 \end{aligned} \quad (351)$$

At the time of this maximum, when

$$t = \frac{2n\pi - \alpha - \theta}{a},$$

the phase of constituent  $A$  will equal

$$(2n\pi - \alpha - \theta) + \alpha$$

and the phase of constituent  $B$  will equal

$$\frac{b}{a}(2n\pi - \alpha - \theta) + \beta$$

Let  $\phi$  = phase of constituent  $B$  - phase of constituent  $A$  at this time. Then

$$\phi = \frac{b-a}{a}(2n\pi - \alpha - \theta) + \beta - \alpha \quad (352)$$

Substituting the above in (351)

$$\begin{aligned} & -Aa \sin \theta + Bb \sin (\phi - \theta) \\ & = -Aa \sin \theta + Bb \sin \phi \cos \theta - Bb \cos \phi \sin \theta \\ & = -(Aa + Bb \cos \phi) \sin \theta + Bb \sin \phi \cos \theta = 0 \end{aligned} \quad (353)$$

Then

$$\tan \theta = \frac{Bb \sin \phi}{Aa + Bb \cos \phi} \quad (354)$$

**239.** For the resultant amplitude at the time of this maximum substitute the values of  $t$  from (350), in (346), and we have

$$\begin{aligned} y &= A \cos (2n \pi - \theta) + B \cos \left[ \frac{b}{a} (2n \pi - \theta - \alpha) + \beta \right] \\ &= A \cos \theta + B \cos \left[ \frac{b-a}{a} (2n \pi - \theta - \alpha) + \beta - \alpha - \theta \right] \\ &= A \cos \theta + B \cos (\phi - \theta) \\ &= A \cos \theta + B \cos \phi \cos \theta + B \sin \phi \sin \theta \\ &= (A + B \cos \phi) \cos \theta + B \sin \phi \sin \theta \\ &= \sqrt{A^2 + B^2 + 2AB \cos \phi} \cos \left( \theta - \tan^{-1} \frac{B \sin \phi}{A + B \cos \phi} \right) \end{aligned} \quad (355)$$

**240.** From (354)

$$\theta = \tan^{-1} \frac{B \sin \phi}{A \frac{a}{b} + B \cos \phi} = \tan^{-1} \frac{\sin \phi}{\frac{Aa}{Bb} + \cos \phi} \quad (356)$$

In the special cases under consideration the ratio  $\frac{a}{b}$  is near unity, and the difference between  $\theta$  and  $\tan^{-1} \frac{B \sin \phi}{A + B \cos \phi}$  is therefore very small, so that the cosine may be taken as unity.

The resultant amplitude may therefore be expressed by

$$\sqrt{A^2 + B^2 + 2AB \cos \phi} = A \sqrt{1 + \frac{B^2}{A^2} + 2 \frac{B}{A} \cos \phi} \quad (357)$$

The true amplitude of the constituent sought being  $A$ , the resultant amplitude must be divided by the factor

$$\sqrt{1 + \frac{B^2}{A^2} + 2 \frac{B}{A} \cos \phi} \quad (358)$$

in order to correct for the influence of the disturbing constituent.

**241.** The corrections for acceleration and amplitude as indicated by formulas (356) and (358) may to advantage be applied to the constants for constituent  $K_1$  for an approximate elimination of the effects of constituent  $P_1$  and to the constants for  $S_2$  for an approximate elimination of the effects of constituents  $K_2$  and  $T_2$ . By taking the relations of the theoretical coefficients for the ratios  $\frac{B}{A}$  and the differences in the equilibrium arguments as the approximate equivalents of the phase differences represented by  $\phi$ , tables may be prepared giving the acceleration and resultant amplitudes with the arguments referring to certain solar elements.

Thus, from table 2, the following values may be obtained.

	$\frac{B}{A}$	$\frac{Aa}{Bb}$	$\phi$
Effect of $P_1$ on $K_1$ .....	0.33086	3.03904	$-2h + \nu' + 180^\circ$ .
Effect of $K_2$ on $S_2$ .....	0.27213	3.66469	$2h - 2\nu''$ .
Effect of $T_2$ on $S_2$ .....	0.05881	17.02813	$-h + p_1$ .

Substituting the above in (356) and (358) we have  
 Effect of  $P_1$  on  $K_1$

$$\text{Acceleration} = \tan^{-1} \frac{\sin (2h - \nu')}{3.0390 - \cos (2h - \nu')} \quad (359)$$

$$\text{Resultant amplitude} = 0.813 \sqrt{1.6767 - \cos (2h - \nu')} \quad (360)$$

Effect of  $K_2$  on  $S_2$

$$\text{Acceleration} = \tan^{-1} \frac{\sin (2h - 2\nu'')}{3.6647 + \cos (2h - 2\nu'')} \quad (361)$$

$$\text{Resultant amplitude} = 0.738 \sqrt{1.9734 + \cos (2h - 2\nu'')} \quad (362)$$

Effect of  $T_2$  on  $S_2$

$$\text{Acceleration} = \tan^{-1} \frac{-\sin (h - p_1)}{17.0281 + \cos (h - p_1)} \quad (363)$$

$$\text{Resultant amplitude} = 0.343 \sqrt{8.5318 + \cos (h - p_1)} \quad (364)$$

**242.** The above formulas give the accelerations and resulting amplitudes for any individual high water. For the correction of the constants derived from a series covering many high waters it is necessary to take averages covering the period of observations. Tables 21 to 26 give such average values for different lengths of series, the argument in each case referring to the beginning of the series.

In the preceding formulas the mean values of the coefficients were taken to obtain the ratios  $\frac{A}{B}$ . To take account of the longitude of the moon's node, the node factor should be introduced. If the mean coefficients are indicated by the subscript  $o$ , formulas (356) and (358) may be written

$$\text{Acceleration} = \tan^{-1} \frac{\sin \phi}{\frac{f(A)A_o a}{f(B)B_o b} + \cos \phi} \quad (365)$$

$$\text{Resultant amplitude} = \sqrt{1 + \left( \frac{f(B)B_o}{f(A)A_o} \right)^2 + 2 \frac{f(B)B_o}{f(A)A_o} \cos \phi} \quad (366)$$

**243.** In the cases under consideration the ratio  $\frac{f(A)}{f(B)}$  will not differ greatly from unity, the ratio  $\frac{A_o a}{B_o b}$  will be rather large compared with  $\cos \phi$ , which can never exceed unity, and the acceleration itself is relatively small. Because of these conditions the following may be taken as the approximate equivalent of (365):

$$\text{Acceleration} = \frac{f(B)}{f(A)} \tan^{-1} \frac{\sin \phi}{\frac{A_o a}{B_o b} + \cos \phi} \quad (367)$$

Also because  $\frac{B_o}{A_o}$  in these cases is small compared with unity, the following may be taken as the approximate equivalent of (366):

$$\text{Resulting amplitude} = 1 + \frac{f(B)}{f(A)} \left[ \sqrt{1 + \left( \frac{B_o}{A_o} \right)^2} + 2 \frac{B_o}{A_o} \cos \phi - 1 \right] \quad (368)$$

To allow for the effects of the longitude of the moon's node, the tabular value of the acceleration should, therefore, be multiplied by the ratio  $\frac{f(B)}{f(A)}$  and the amount by which the resultant amplitude differs from unity by the same factor. In the particular cases under consideration the factor  $f$ , for constituents  $P_1$ ,  $S_2$ , and  $T_2$ , is unity for each. Therefore, for the effect of  $P_1$  on  $K_1$ , the ratio  $\frac{f(B)}{f(A)} = \frac{1}{f(K_1)} = F(K_1)$ , and for the effect of  $K_2$  upon  $S_2$ , this ratio is  $f(K_2)$ . For the effect of  $T_2$  upon  $S_2$  the ratio is unity.

#### ELIMINATION

**244.** Because of the limited length of a series of observations analyzed the amplitudes and epochs of the constituents as obtained by the processes already described are only approximately freed from the effects of each other. The separation of two constituents from each other might be satisfactorily accomplished by having the length of series equal to a multiple of the synodic period of the two constituents. To completely effect the separation of all the constituents from each other by the same process would require a series of such a length that it would contain an exact multiple of the period of each constituent. The length of such a series would be too great to be given practical consideration. In general, it is therefore desirable to apply certain corrections to the constants as directly obtained from the analysis in order to eliminate the residual effects of the constituent upon each other.

**245.** Let  $A$  be the designation of a constituent for which the true constants are sought and let  $B$  be the general designation for each of the other constituents in the tide, the effects of which are to be eliminated from constituent  $A$ .

Let the original tide curve which has been analyzed be represented by the formula

$$y = A \cos(at + \alpha) + \sum B \cos(bt + \beta) \quad (369)$$

in which

$y$ =the height of the tide above mean sea level at any time  $t$ .  
 $t$ =time reckoned in mean solar hours from the beginning of the series as the origin.

$A=R(A)$ =true amplitude of the constituent  $A$  for the time covered by series of observations.

$B=R(B)$ =true amplitude of constituent  $B$  for the time covered by series of observations.

$\alpha=-\zeta(A)$ =true initial phase of constituent  $A$  at beginning of series.

$\beta = -\zeta(B)$  = true initial phase of constituent  $B$  at beginning of series.

$a$  = speed of constituent  $A$ .

$b$  = speed of constituent  $B$ .

246. Formula (369) may be written

$$\begin{aligned} y &= A \cos \alpha \cos at + \Sigma B \cos \{(b-a)t + \beta\} \cos at \\ &\quad - A \sin \alpha \sin at - \Sigma B \sin \{(b-a)t + \beta\} \sin at \\ &= [A \cos \alpha + \Sigma B \cos \{(b-a)t + \beta\}] \cos at \\ &\quad - [A \sin \alpha + \Sigma B \sin \{(b-a)t + \beta\}] \sin at \end{aligned} \quad (370)$$

The mean values of the coefficients of  $\cos at$  and  $\sin at$  of formula (370) correspond to the coefficients  $C_p$  and  $S_p$  of formulas (295) and (296) which are obtained from the summations for constituent  $A$ .

247. Let  $A'$  and  $\alpha'$  = the uneliminated amplitude and initial phase, respectively, of constituent  $A$ , as obtained directly from the analysis.

The equation of the uneliminated constituent  $A$  tide may be written

$$y = A' \cos (at + \alpha') = A' \cos \alpha' \cos at - A' \sin \alpha' \sin at \quad (371)$$

Comparing (370) and (371), it will be found that

$$A' \cos \alpha' = \text{mean value of } [A \cos \alpha + \Sigma B \cos \{(b-a)t + \beta\}] \quad (372)$$

$$A' \sin \alpha' = \text{mean value of } [A \sin \alpha + \Sigma B \sin \{(b-a)t + \beta\}] \quad (373)$$

248. Let  $\tau$  = length of series in mean solar hours. Then the mean value of

$B \cos \{(b-a)t + \beta\}$  within the limits  $t=0$  and  $t=\tau$ , is

$$\begin{aligned} \frac{1}{\tau} \int_0^\tau B \cos \{(b-a)t + \beta\} dt &= \frac{180}{\pi} \frac{B}{(b-a)\tau} [\sin \{(b-a)\tau + \beta\} - \sin \beta] \\ &= \frac{180}{\pi} \frac{\sin \frac{1}{2}(b-a)\tau}{\frac{1}{2}(b-a)\tau} B \cos \{\frac{1}{2}(b-a)\tau + \beta\} \end{aligned} \quad (374)$$

The mean value of  $B \sin \{(b-a)t + \beta\}$  within the same limits is

$$\begin{aligned} \frac{1}{\tau} \int_0^\tau B \sin \{(b-a)t + \beta\} dt &= -\frac{180}{\pi} \frac{B}{(b-a)\tau} [\cos \{(b-a)\tau + \beta\} - \cos \beta] \\ &= \frac{180}{\pi} \frac{\sin \frac{1}{2}(b-a)\tau}{\frac{1}{2}(b-a)\tau} B \sin \{\frac{1}{2}(b-a)\tau + \beta\} \end{aligned} \quad (375)$$

Substituting (374) and (375) in (372) and (373), and for brevity letting

$$F_b = \frac{180}{\pi} \frac{\sin \frac{1}{2}(b-a)\tau}{\frac{1}{2}(b-a)\tau} B \quad (376)$$

we have

$$A' \cos \alpha' = A \cos \alpha + \Sigma F_b \cos \{\frac{1}{2}(b-a)\tau + \beta\} \quad (377)$$

$$A' \sin \alpha' = A \sin \alpha + \Sigma F_b \sin \{\frac{1}{2}(b-a)\tau + \beta\} \quad (378)$$

Transposing,

$$A \cos \alpha = A' \cos \alpha' - \Sigma F_b \cos \{\frac{1}{2}(b-a)\tau + \beta\} \quad (379)$$

$$A \sin \alpha = A' \sin \alpha' - \Sigma F_b \sin \{\frac{1}{2}(b-a)\tau + \beta\} \quad (380)$$

Multiplying (379) and (380) by  $\sin \alpha'$  and  $\cos \alpha'$ , respectively,

$$A \sin \alpha' \cos \alpha = A' \sin \alpha' \cos \alpha' - \sum F_b \cos \left\{ \frac{1}{2}(b-a)\tau + \beta \right\} \sin \alpha' \quad (381)$$

$$A \cos \alpha' \sin \alpha = A' \sin \alpha' \cos \alpha' - \sum F_b \sin \left\{ \frac{1}{2}(b-a)\tau + \beta \right\} \cos \alpha' \quad (382)$$

Subtracting (382) from (381)

$$A \sin (\alpha' - \alpha) = \sum F_b \sin \left\{ \frac{1}{2}(b-a)\tau + \beta - \alpha' \right\} \quad (383)$$

Multiplying (379) and (380) by  $\cos \alpha'$  and  $\sin \alpha'$ , respectively,

$$A \cos \alpha' \cos \alpha = A' \cos^2 \alpha' - \sum F_b \cos \left\{ \frac{1}{2}(b-a)\tau + \beta \right\} \cos \alpha' \quad (384)$$

$$A \sin \alpha' \sin \alpha = A' \sin^2 \alpha' - \sum F_b \sin \left\{ \frac{1}{2}(b-a)\tau + \beta \right\} \sin \alpha' \quad (385)$$

Taking the sum of (384) and (385)

$$A \cos (\alpha' - \alpha) = A' - \sum F_b \cos \left\{ \frac{1}{2}(b-a)\tau + \beta - \alpha' \right\} \quad (386)$$

Dividing (383) by (386)

$$\tan (\alpha' - \alpha) = \frac{\sum F_b \sin \left\{ \frac{1}{2}(b-a)\tau + \beta - \alpha' \right\}}{A' - \sum F_b \cos \left\{ \frac{1}{2}(b-a)\tau + \beta - \alpha' \right\}} \quad (387)$$

From (386)

$$A = \frac{A' - \sum F_b \cos \left\{ \frac{1}{2}(b-a)\tau + \beta - \alpha' \right\}}{\cos (\alpha' - \alpha)} \quad (388)$$

**249.** Substituting the value  $F_b$  from (376) and the equivalents  $R'(A)$ ,  $R(A)$ ,  $R(B)$ ,  $-\xi'(A) - \xi(A)$ , and  $-\xi(B)$  for  $A'$ ,  $A$ ,  $B$ ,  $\alpha'$ ,  $\alpha$ , and  $\beta$ , respectively, we have by (387) and (388)

$$\tan [\xi(A) - \xi'(A)] =$$

$$\frac{\sum \frac{180}{\pi} \frac{\sin \frac{1}{2}(b-a)\tau}{\frac{1}{2}(b-a)\tau} R(B) \sin \left\{ \frac{1}{2}(b-a)\tau - \xi(B) + \xi'(A) \right\}}{R'(A) - \sum \frac{180}{\pi} \frac{\sin \frac{1}{2}(b-a)\tau}{\frac{1}{2}(b-a)\tau} R(B) \cos \left\{ \frac{1}{2}(b-a)\tau - \xi(B) + \xi'(A) \right\}} \quad (389)$$

$$R'(A) - \sum \frac{180}{\pi} \frac{\sin \frac{1}{2}(b-a)\tau}{\frac{1}{2}(b-a)\tau} R(B) \cos \left\{ \frac{1}{2}(b-a)\tau - \xi(B) + \xi'(A) \right\} \\ R(A) = \frac{\cos [\xi(A) - \xi'(A)]}{\sin [\xi(A) - \xi'(A)]} \quad (390)$$

**250.** Formula (389) gives an expression for obtaining the difference to be applied to the uneliminated  $\xi'(A)$  in order to obtain the true  $\xi(A)$ , and formula (390) gives an expression for obtaining the true amplitude  $R(A)$ . These formulas cannot, however, be rigorously applied, because the true values of  $R(B)$  and  $\xi(B)$  of the disturbing constituents are, in general, not known, but very satisfactory results may be obtained by using the approximate values of  $R(B)$  and  $\xi(B)$  derived from the analysis or by inference.

By a series of successive approximations, using each time in the formulas the newly eliminated values for the disturbing constituents, any desired degree of refinement may be obtained, but the first approximation is usually sufficient and all that is justified because of the greater irregularities existing from other causes.

**251.** Form 245 (fig. 19) provides for the computations necessary in applying formulas (389) and (390). In these formulas the factors represented by  $\frac{180}{\pi} \frac{\sin \frac{1}{2}(b-a)\tau}{\frac{1}{2}(b-a)\tau}$  and the angles represented by

$\frac{1}{2}(b-a)\tau$  will depend upon the length of series; but for any given length of series they will be constant for all times and places. Table 29 has been computed to give these quantities for different lengths of series. The factor as directly obtained may be either positive or negative, but for convenience the tabular values are all given as positive, and when the factor as directly obtained is negative the angle has been modified by  $\pm 180^\circ$  in order to compensate for the change of sign in the factor and permit the tabular values to be used directly in formulas (389) and (390).

252. An examination of formulas (389) and (390) will show that the disturbing effect of one constituent upon another will depend largely upon the magnitude of the fraction  $\frac{\sin \frac{1}{2}(b-a)\tau}{\frac{1}{2}(b-a)\tau}$ . Assuming that  $b$  is not equal to  $a$ , this fraction and the disturbing effect it represents will approach zero as the length of series  $\tau$  approaches in value  $\frac{360^\circ}{(b-a)}$ , or any multiple thereof, or, in other words, as  $\tau$  approaches in length any multiple of the synodic period of constituents  $A$  and  $B$ . Also, since the numerator of the fraction can never exceed unity, while the denominator may be increased indefinitely, the value of the fraction will, in general, be diminished by increasing the length of series and will approach zero as  $\tau$  approaches infinity. The greater the difference  $(b-a)$  between the speeds of the two constituents the less will be their disturbing effects upon each other. For this reason the effects upon each other of the diurnal and semidiurnal constituents are usually considered as negligible.

253. The quantities  $R(B)$  and  $\xi(B)$  of formulas (389) and (390) refer to the true amplitudes and epochs of the disturbing constituents. These true values being in general unknown when the elimination process is to be applied, it is desirable that there should be used in the formulas the closest approximation to such values as are obtainable. If the series of observations covers a period of a year or more, the amplitudes and epochs as directly obtained from the analysis may be considered sufficiently close approximations for use in the formulas. For short series of observations, however, the values as directly obtained for the amplitudes and epochs of some of the constituents may be so far from the true values that they are entirely unserviceable for use in the formulas. In such cases inferred values for the disturbing constituents should be used.

#### LONG-PERIOD CONSTITUENTS

254. The preceding discussions have been especially applicable to the reduction of the short-period constituents—those having a period of a constituent day or less. They are the constituents that determine the daily or semidaily rise and fall of the tide. Consideration will now be given to the long-period tides which affect the mean level of the water from day to day, but which have practically little or no effect upon the times of the high and low waters. There are five such long-period constituents that are usually treated in works on harmonic analysis—the lunar fortnightly Mf, the lunisolar synodic fortnightly MSf, the lunar monthly Mm, the solar semiannual Ssa, and the solar annual Sa. The first three are usually too small to be of practical importance, but the last two, depending largely upon

meteorological conditions, often have an appreciable effect upon the mean daily level of the water.

**255.** To obtain the long-period constituents, methods similar to those adopted for the short-period constituents with certain modifications may be used. For the fortnightly and monthly constituents the constituent month may be divided into 24 equal parts, analogous to the 24 constituent hours of the day. Similarly, for the semiannual and annual constituents the constituent year may be divided into 24 equal parts, although it will often be found more convenient to divide the year into 12 parts to correspond approximately with the 12 calendar months.

**256.** Instead of distributing the individual hourly heights, as for the short-period constituents, a considerable amount of labor can be saved by using the daily sums of these heights. The mean of each sum is to be considered as applying to the middle instant of the period from 0 hour to 23d hour; that is, at the 11.5 hour of the day. If the constituent month or year is divided into 24 equal parts, the instants separating the groups may be numbered consecutively, like the hours, from 0 to 23, with the 0 instant of the first group taken at the exact beginning of the series. A table may now be prepared (table 34) which will show to which division each daily sum, or mean, of the series must be assigned.

**257. Letting**

$a$ =the hourly speed of any constituent, in degrees.

$p=1$  when applied to a monthly or an annual constituent, and

$p=2$  when applied to a fortnightly or a semiannual constituent.

$d$ =day of series.

$s$ =solar hour of day.

Then

$$1 \text{ constituent period} = \frac{360}{a} \text{ solar hours} \quad (391)$$

and

$$1 \text{ constituent month} = \frac{360p}{a} \text{ solar hours} \quad (392)$$

also

$$1 \text{ constituent year} = \frac{360p}{a} \text{ solar hours} \quad (393)$$

Dividing the constituent month or year into 24 equal parts, the length of

$$1 \text{ constituent division} = \frac{15p}{a} \text{ solar hours} \quad (394)$$

Therefore, to express the time of any solar hour in units of the constituent divisions to which the solar hourly heights are to be assigned, the solar hour should be multiplied by the factor  $a/15p$ .

Thus,

$$\begin{aligned} \text{Constituent division} &= \frac{a}{15p} \text{ (solar hour of series)} \\ &= \frac{a}{15p} [24(d-1) + s] \\ &= \frac{a}{15p} [24(d-1) + 11.5] \end{aligned} \quad (395)$$

since in using the daily sums, the solar hour of the day to which each such sum applies will always be 11.5 hour.

By substituting the speeds of the constituents from table 2 the following numerical values are obtained for the coefficient  $\frac{a}{15p}$ :

$$\text{Mf, } 0.036,601,10; \text{ MSf, } 0.033,863,19; \text{ Mm, } 0.036,291,65; \\ \text{Sa and Ssa, } 0.002,737,91.$$

By using the appropriate coefficient and substituting successively the numerals corresponding to the day of series ( $d$ ), the corresponding value of the constituent division to which each daily sum is to be assigned may be readily obtained. The value of such division as obtained directly from the formula will usually be a mixed number. For table 34 the nearest integral number, less any multiple of 24, is used.

**258.** The distribution of the daily sums for the analysis of the long-period constituents may be conveniently accomplished by copying such sums in Form 142 (fig. 12), taking the constituent divisions as the equivalents of the constituent hours and using table 34 to determine the division or hour to which each sum should be assigned. The total sum and mean for each division may then be readily obtained. These means can then be treated as the hourly means of the short-period tides according to the processes outlined in Form 194 (fig. 16) with such modifications as will now be described.

**259.** In using the daily means as ordinates of a long-period constituent consideration must be given to the residual effects of any of the short-period constituents upon such means and steps taken to clear the means of these effects when necessary. Constituent  $S_2$  with a period commensurate with the solar day, may be considered as being completely eliminated from each daily mean. Constituents  $K_1$  and  $K_2$  are very nearly eliminated because the  $K$  day is very nearly equal to the solar day. Other short-period constituents may affect the daily means to a greater or less extent, depending largely upon their amplitudes. Of these the principal ones are constituents  $M_2$ ,  $N_2$ , and  $O_1$ . In the distribution and grouping of the daily means for the analysis of the several long-period constituents the disturbing effects of the short-period constituents just enumerated, excepting the effects of  $M_2$  upon MSf, will be greatly reduced, and in a series covering several years may be practically eliminated. Because the period of MSf is the same as the synodic period of  $M_2$  and  $S_2$  there will always remain a residual effect of the constituent  $M_2$  in the constituent MSf sums of the daily means, no matter how long the series, which must be removed by a special process.

**260.** Let the equation of one of the short-period constituents be

$$y = A \cos(at + \alpha) \quad (396)$$

Letting  $d$ =day of series, the values of  $t$  for the hours 0 to 23 of  $d$  day will be

$$24(d-1), 24(d-1)+1, 24(d-1)+2, \dots, 24(d-1)+23.$$

Substituting these values for  $t$  in (396) and designating the corresponding values of the ordinate  $y$  as  $y_0, y_1, y_2, \dots, y_{23}$  the following are obtained:

$$\left. \begin{aligned} y_0 &= A \cos [24(d-1)a + \alpha] \\ y_1 &= A \cos [24(d-1)a + \alpha + a] \\ y_2 &= A \cos [24(d-1)a + \alpha + 2a] \\ &\dots \\ y_{23} &= A \cos [24(d-1)a + \alpha + 23a] \end{aligned} \right\} \quad (397)$$

Representing the mean of these 24 ordinates for  $d$  day by  $y_d$ , we have

$$\begin{aligned} y_d &= \frac{1}{24} A \cos \{24(d-1)a + \alpha\} [1 + \cos a + \cos 2a + \dots + \cos 23a] \\ &\quad - \frac{1}{24} A \sin \{24(d-1)a + \alpha\} [\sin a + \sin 2a + \dots + \sin 23a] \\ &= \frac{1}{24} A \frac{\sin 12a}{\sin \frac{1}{2}a} \left[ \cos \{24(d-1)a + \alpha\} \cos \frac{23}{2} a \right. \\ &\quad \left. - \sin \{24(d-1)a + \alpha\} \sin \frac{23}{2} a \right] \\ &= \frac{1}{24} A \frac{\sin 12a}{\sin \frac{1}{2}a} \cos \{24(d-1)a + \alpha + 11.5a\} \end{aligned} \quad (398)$$

**261.** Formula (398), representing the average value of the constituent  $A$  ordinates contained in the daily mean for  $d$  day, is the correction or clearance that must be subtracted from the mean for that day in order to eliminate the effects of  $A$ . It will be noted that if we let  $A$  represent any of the solar constituents,  $S_1$ ,  $S_2$ ,  $S_3$ ,  $S_4$ , etc., the factor  $\sin 12a$ , and consequently the entire formula, becomes zero for all values of  $d$ . By formula (398) clearances for each of the disturbing short-period constituents for each day of series may be computed and these clearances then applied individually to the daily means, or, if first multiplied by the factor 24, to the daily sums.

**262.** The labor involved in making independent calculations for the clearance of the effect of each short-period constituent for each day of series would be considerable, but this may be avoided to a large extent by means of a tide-computing machine.

If we let  $t$  = time reckoned in mean solar hours from the beginning of the series, then for any value of  $y_d$ , which must apply to the 11.5 hour of  $d$  day,

$$t = 24(d-1) + 11.5$$

and

$$at = 24(d-1)a + 11.5a \quad (399)$$

If the above equivalent is substituted in (398) and  $y_d$  replaced by  $y_a$ , we have

$$y_a = \frac{1}{24} A \frac{\sin 12a}{\sin \frac{1}{2}a} \cos (at + \alpha) \quad (400)$$

which represents a continuous function of  $t$ ; and for any value of  $t$  corresponding to the 11.5 hour of  $d$  day the corresponding value of  $y_a$  will be  $y_d$ . This formula is the same as that for the short-period

constituent  $A$ , except that it includes the factor  $\frac{1}{24} \frac{\sin 12a}{\sin \frac{1}{2}a}$  in the coefficient. The speed  $a$  is a known constant and the values of  $A$  and  $\alpha$  are presumed to have already been determined from the harmonic analysis of the short-period constituents. Similarly, the disturbing effects of other short-period constituents may be represented by

$$\begin{aligned} y_b &= \frac{1}{24} B \frac{\sin 12b}{\sin \frac{1}{2}b} \cos(bt + \beta) \\ y_c &= \frac{1}{24} C \frac{\sin 12c}{\sin \frac{1}{2}c} \cos(ct + \gamma) \\ &\text{etc.} \end{aligned} \quad (401)$$

The combined disturbing effect of all the short-period constituents may, therefore, be represented by the equation

$$\begin{aligned} y = y_a + y_b + \text{etc.} &= \frac{1}{24} A \frac{\sin 12a}{\sin \frac{1}{2}a} \cos(at + \alpha) \\ &+ \frac{1}{24} B \frac{\sin 12b}{\sin \frac{1}{2}b} \cos(bt + \beta) + \text{etc.} \end{aligned} \quad (402)$$

**263.** This formula is adapted to use on the tide-computing machine. With the constituent cranks set in accordance with the coefficients and initial epochs of the above formula, the machine will indicate the values of  $y$  corresponding to successive values of  $t$ . The values of  $y$  desired for the clearances are those which correspond to  $t$  at the 11.5 hour on each day. Thus, the clearance for each successive day of series may be read directly from the dials of the machine. In practice, it may be found more convenient to use the daily sums rather than the daily means for the analysis. In this case the coefficients of the terms of (402) should be multiplied by the factor 24 before being used in the tide-computing machine.

**264.** Assuming that all the daily sums are used in the analysis, the augmenting factor represented by formula (308) which is used for the short-period constituent is also applicable to the long-period constituents, with  $p$  representing the number of constituent periods in a constituent month or year. Thus, for Mm and Sa,  $p$  equals 1, and for Mf, MSf, and Ssa,  $p$  equals 2. For the long-period constituents a further correction or augmenting factor is necessary, because the mean or sum of the 24 hourly heights of the day is used to represent the single ordinate at the 11.5 hour of the day.

**265.** If we let formula (396) be the equation of the long-period constituent sought, formula (400) will give the mean value of the 24 ordinates of the day which, in the grouping for the analysis, is taken as representing the 11.5 hour of the day or the  $t_d$  hour of the series. Since the true constituent ordinate for this hour should be  $A \cos(at_d + \alpha)$ , it is evident that an augmenting factor of  $24 \frac{\sin \frac{1}{2}a}{\sin 12a}$  must be applied to the mean ordinates as derived from the sum of the 24 hourly heights of the day in order to reduce the means to the 11.5 hour of each day.

**266.** The complete augmenting factor for the long-period constituents, the year or month being represented by 24 means, will be obtained by combining the above factor with that given in formula (308). Thus

$$\text{augmenting factor} = \frac{\pi p}{24 \sin \frac{15p}{2}} \times \frac{24 \sin \frac{1}{2}a}{\sin 12a} \quad (403)$$

If the year or month is represented by only 12 means as when monthly means are used in evaluating  $S_a$  and  $S_{sa}$ , the formula becomes

$$\text{augmenting factor} = \frac{\pi p}{12 \sin 15p} \times \frac{24 \sin \frac{1}{2}a}{\sin 12a} \quad (404)$$

Values obtained from these formulas are given in table 20.

**267.** The following method of reducing the long-period tides, which conforms to the system outlined by Sir George H. Darwin, differs to some extent from that just described. In this discussion it is assumed that a series of 365 days is used. Let the entire tide due to the five long-period constituents already named be represented by the equation

$$y = A \cos(at + \alpha) + B \cos(bt + \beta) + C \cos(ct + \gamma) + D \cos(dt + \delta) + E \cos(et + \epsilon) \quad (405)$$

**268.** For convenience in this discussion let  $t$  be reckoned from the 11.5th solar hour of the first day of series instead of the midnight beginning that day. Every value of  $t$  to which the daily means refer will then be either 0 or a multiple of 24.

Let  $A'$ ,  $B'$ ,  $C'$ ,  $D'$ , and  $E'$ , equal

$$\begin{aligned} &A \cos \alpha, B \cos \beta, C \cos \gamma, D \cos \delta, \text{ and } E \cos \epsilon, \text{ respectively, and} \\ &A'', B'', C'', D'', \text{ and } E'', \text{ equal} \\ &-A \sin \alpha, -B \sin \beta, -C \sin \gamma, -D \sin \delta, \text{ and } -E \sin \epsilon, \text{ respectively.} \end{aligned} \quad (406)$$

Then formula (405) may be written

$$y = A' \cos at + B' \cos bt + C' \cos ct + D' \cos dt + E' \cos et + A'' \sin at + B'' \sin bt + C'' \sin ct + D'' \sin dt + E'' \sin et \quad (407)$$

In the above equation there are 10 unknown quantities,  $A'$ ,  $A''$ ,  $B'$ ,  $B''$ , etc., for which values are sought in order to obtain from them the amplitudes and epochs of the five long-period constituents. The most probable values of these quantities may be found by the least square adjustment.

**269.** Let  $y_1, y_2, \dots, y_{365}$  represent the daily means for a 365 day series, as obtained from observations. If we let  $n$  be any day of the series, the value of  $t$  to which that mean applies will be  $24(n-1)$ . By substituting in formula (407) the successive values of  $y$  and the values of  $t$  to which they correspond, 365 observational equations are formed as follows:

$$\left. \begin{aligned} y_1 &= A' \cos 0 + B' \cos 0 + \dots \\ &\quad + A'' \sin 0 + B'' \sin 0 + \dots \\ y_2 &= A' \cos 24a + B' \cos 24b + \dots \\ &\quad + A'' \sin 24a + B'' \sin 24b + \dots \\ y_{365} &= A' \cos 24 \times 364a + B' \cos 24 \times 364b + \dots \\ &\quad + A'' \sin 24 \times 364a + B'' \sin 24 \times 364b + \dots \end{aligned} \right\} \quad (408)$$

**270.** A normal equation is now formed for each unknown quantity by multiplying each observational equation by the coefficient of the unknown quantity in that equation and adding the results. Thus, for the unknown quantity  $A'$ , we have

$$\left. \begin{aligned} y_1 \cos 0 &= A' \cos^2 0 + B' \cos 0 \cos 0 + \dots \\ &\quad + A'' \sin 0 \cos 0 + B'' \sin 0 \cos 0 + \dots \\ y_2 \cos 24a &= A' \cos^2 24a + B' \cos 24b \cos 24a + \dots \\ &\quad + A'' \sin 24a \cos 24a + B'' \sin 24b \cos 24a + \dots \\ y_{365} \cos (24 \times 364a) &= A' \cos^2 (24 \times 364a) \\ &\quad + B' \cos (24 \times 364b) \cos (24 \times 364a) + \dots \\ &\quad + A'' \sin (24 \times 364a) \cos (24 \times 364a) \\ &\quad + B'' \sin (24 \times 364b) \cos (24 \times 364a) + \dots \end{aligned} \right\} \quad (409)$$

Summing

$$\sum_{n=1}^{n=365} y_n \cos 24(n-1)a = A' \sum_{n=1}^{n=365} \cos^2 24(n-1)a + A'' \sum_{n=1}^{n=365} \sin 24(n-1)a \cos 24(n-1)a + B' \sum_{n=1}^{n=365} \cos 24(n-1)b \cos 24(n-1)a + B'' \sum_{n=1}^{n=365} \sin 24(n-1)b \cos 24(n-1)a + C' \sum_{n=1}^{n=365} \cos 24(n-1)c \cos 24(n-1)a + C'' \sum_{n=1}^{n=365} \sin 24(n-1)c \cos 24(n-1)a + D' \sum_{n=1}^{n=365} \cos 24(n-1)d \cos 24(n-1)a + D'' \sum_{n=1}^{n=365} \sin 24(n-1)d \cos 24(n-1)a + E' \sum_{n=1}^{n=365} \cos 24(n-1)e \cos 24(n-1)a + E'' \sum_{n=1}^{n=365} \sin 24(n-1)e \cos 24(n-1)a \quad (410)$$

which is the normal equation for the unknown quantity  $A'$ .

**271.** In a similar manner we have for the normal equation for the quantity  $A''$

$$\begin{aligned}
 & \Sigma y_n \sin 24(n-1)a \\
 & = A' \Sigma \cos 24(n-1)a \sin 24(n-1)a + A'' \Sigma \sin^2 24(n-1)a \\
 & + B' \Sigma \cos 24(n-1)b \sin 24(n-1)a + B'' \Sigma \sin 24(n-1)b \sin 24(n-1)a \\
 & + C' \Sigma \cos 24(n-1)c \sin 24(n-1)a + C'' \Sigma \sin 24(n-1)c \sin 24(n-1)a \\
 & + D' \Sigma \cos 24(n-1)d \sin 24(n-1)a + D'' \Sigma \sin 24(n-1)d \sin 24(n-1)a \\
 & + E' \Sigma \cos 24(n-1)e \sin 24(n-1)a + E'' \Sigma \sin 24(n-1)e \sin 24(n-1)a
 \end{aligned} \tag{411}$$

the limits of  $n$  being the same as before.

Normal equations of forms similar to (410) and (411) are easily obtained for the other unknown quantities.

**272.** By changing the notation of formulas (265) to (267) the following relations may be derived:

$$\begin{aligned}
 \sum_{n=1}^{365} \cos^2 24(n-1)a &= \frac{1}{2}n + \frac{1}{2} \frac{\sin 24na \cos 24(n-1)a}{\sin 24a} \\
 &= 182\frac{1}{2} + \frac{1}{2} \frac{\sin 8760a \cos 8736a}{\sin 24a}
 \end{aligned} \tag{412}$$

$$\begin{aligned}
 \sum_{n=1}^{365} \sin^2 24(n-1)a &= \frac{1}{2}n - \frac{1}{2} \frac{\sin 24na \cos 24(n-1)a}{\sin 24a} \\
 &= 182\frac{1}{2} - \frac{1}{2} \frac{\sin 8760a \cos 8736a}{\sin 24a}
 \end{aligned} \tag{413}$$

$$\begin{aligned}
 \sum_{n=1}^{365} \cos 24(n-1)b \cos 24(n-1)a \\
 &= \frac{1}{2} \frac{\sin 12n(b-a) \cos 12(n-1)(b-a)}{\sin 12(b-a)} \\
 &\quad + \frac{1}{2} \frac{\sin 12n(b+a) \cos 12(n-1)(b+a)}{\sin 12(b+a)} \\
 &= \frac{1}{2} \frac{\sin 4380(b-a) \cos 4368(b-a)}{\sin 12(b-a)} \\
 &\quad + \frac{1}{2} \frac{\sin 4380(b+a) \cos 4368(b+a)}{\sin 12(b+a)}
 \end{aligned} \tag{414}$$

$$\begin{aligned}
 \sum_{n=1}^{365} \sin 24(n-1)b \sin 24(n-1)a \\
 &= \frac{1}{2} \frac{\sin 12n(b-a) \cos 12(n-1)(b-a)}{\sin 12(b-a)} \\
 &\quad - \frac{1}{2} \frac{\sin 12n(b+a) \cos 12(n-1)(b+a)}{\sin 12(b+a)} \\
 &= \frac{1}{2} \frac{\sin 4380(b-a) \cos 4368(b-a)}{\sin 12(b-a)} \\
 &\quad - \frac{1}{2} \frac{\sin 4380(b+a) \cos 4368(b+a)}{\sin 12(b+a)}
 \end{aligned} \tag{415}$$

$$\begin{aligned}
 & \sum_{n=1}^{n=365} \sin 24(n-1)b \cos 24(n-1)a \\
 &= \frac{1}{2} \frac{\sin 12n(b-a) \sin 12(n-1)(b-a)}{\sin 12(b-a)} \\
 &\quad + \frac{1}{2} \frac{\sin 12n(b+a) \sin 12(n-1)(b+a)}{\sin 12(b+a)} \\
 &= \frac{1}{2} \frac{\sin 4380(b-a) \sin 4368(b-a)}{\sin 12(b-a)} \\
 &\quad + \frac{1}{2} \frac{\sin 4380(b+a) \sin 4368(b+a)}{\sin 12(b+a)} \tag{416}
 \end{aligned}$$

**273.** By substituting in (412) to (416) the numerical values of  $a$ ,  $b$ , etc., from table 2, the corresponding equivalents for these expressions are obtained. These, in turn, may be substituted in (410), (411), and similar equations for the other unknown quantities to obtain the 10 normal equations given below. In preparing these equations the symbols  $a$ ,  $b$ ,  $c$ ,  $d$ , and  $e$  are taken, respectively, as the speeds of constituents Mm, Mf, MSf, Sa, and Ssa.

$$\left. \begin{aligned}
 & \sum_{n=1}^{n=365} y_n \cos 24(n-1)a \\
 &= 183.05A' + 0.72B' + 0.76C' + 4.88D' + 4.96E' \\
 &\quad + 2.14A'' + 4.29B'' + 5.04C'' - 0.34D'' - 0.70E'' \\
 & \sum_{n=1}^{n=365} y_n \sin 24(n-1)a \\
 &= 2.14A' - 4.15B' - 4.90C' + 3.80D' + 3.88E' \\
 &\quad + 181.95A'' + 1.01B'' + 1.06C'' + 0.34D'' + 0.68E'' \\
 & \sum_{n=1}^{n=365} y_n \cos 24(n-1)b \\
 &= 0.72A' + 183.17B' + 0.56C' - 1.50D' - 1.51E' \\
 &\quad - 4.15A'' + 0.88B'' + 0.92C'' - 0.09D'' - 0.18E'' \\
 & \sum_{n=1}^{n=365} y_n \sin 24(n-1)b \\
 &= 4.29A' + 0.88B' + 0.92C' + 3.05D' + 3.06E' \\
 &\quad + 1.01A'' + 181.83B'' - 0.80C'' - 0.08D'' - 0.17E'' \\
 & \sum_{n=1}^{n=365} y_n \cos 24(n-1)c \\
 &= 0.76A' + 0.56B' + 183.19C' - 1.68D' - 1.70E' \\
 &\quad - 4.90A'' + 0.92B'' + 0.97C'' - 0.11D'' - 0.21E'' \\
 & \sum_{n=1}^{n=365} y_n \sin 24(n-1)c \\
 &= 5.04A' + 0.92B' + 0.97C' + 3.24D' + 3.25E' \\
 &\quad + 1.06A'' - 0.80B'' + 181.81C'' - 0.10D'' - 0.20E'' \\
 & \sum_{n=1}^{n=365} y_n \cos 24(n-1)d \\
 &= 4.88A' - 1.50B' - 1.68C' + 182.38D' - 0.24E' \\
 &\quad + 3.80A'' + 3.05B'' + 3.24C'' + 0.00D'' + 0.01E'' \\
 & \sum_{n=1}^{n=365} y_n \sin 24(n-1)d \\
 &= -0.34A' - 0.09B' - 0.11C' + 0.00D' + 0.00E' \\
 &\quad + 0.34A'' - 0.08B'' - 0.10C'' + 182.62D'' + 0.00E'' \tag{417a}
 \end{aligned} \right\}$$

$$\left. \begin{aligned}
 & \sum_{n=1}^{n=365} y_n \cos 24(n-1)e \\
 &= -0.76A' + 0.56B' + 183.19C' - 1.68D' - 1.70E' \\
 &\quad - 4.90A'' + 0.92B'' + 0.97C'' - 0.11D'' - 0.21E'' \\
 & \sum_{n=1}^{n=365} y_n \sin 24(n-1)e \\
 &= -5.04A' - 0.92B' - 0.97C' - 3.24D' - 3.25E' \\
 &\quad - 1.06A'' + 0.80B'' - 181.81C'' + 0.10D'' + 0.20E'' \tag{417b}
 \end{aligned} \right\}$$

$$\left. \begin{aligned}
 & \sum_{n=1}^{n=365} y_n \cos 24(n-1)d \\
 &= 4.88A' - 1.50B' - 1.68C' + 182.38D' - 0.24E' \\
 &\quad + 3.80A'' + 3.05B'' + 3.24C'' + 0.00D'' + 0.01E'' \\
 & \sum_{n=1}^{n=365} y_n \sin 24(n-1)d \\
 &= -0.34A' - 0.09B' - 0.11C' + 0.00D' + 0.00E' \\
 &\quad + 0.34A'' - 0.08B'' - 0.10C'' + 182.62D'' + 0.00E'' \tag{417c}
 \end{aligned} \right\}$$

$$\left. \begin{aligned}
 & \sum_{n=1}^{n=365} y_n \cos 24(n-1)e \\
 &= -0.76A' + 0.56B' + 183.19C' - 1.68D' - 1.70E' \\
 &\quad - 4.90A'' + 0.92B'' + 0.97C'' - 0.11D'' - 0.21E'' \\
 & \sum_{n=1}^{n=365} y_n \sin 24(n-1)e \\
 &= -5.04A' - 0.92B' - 0.97C' - 3.24D' - 3.25E' \\
 &\quad - 1.06A'' + 0.80B'' - 181.81C'' + 0.10D'' + 0.20E'' \tag{417d}
 \end{aligned} \right\}$$

$$\left. \begin{aligned} \sum_{n=1}^{n=365} y_n \cos 24(n-1)\alpha \\ = 4.96A' - 1.51B' - 1.70C' - 0.24D' + 182.38E' \\ + 3.88A'' + 3.06B'' + 3.25C'' + 0.00D'' + 0.00E'' \\ \sum_{n=1}^{n=365} y_n \sin 24(n-1)\alpha \\ = -0.70A' - 0.18B' - 0.21C' + 0.01D' + 0.00E' \\ + 0.68A'' - 0.17B'' - 0.20C'' + 0.00D'' + 182.62E'' \end{aligned} \right\} \quad (417e)$$

**274.** The numerical value of the first member of each of the above normal equations is obtained from the observations by taking the sum of the product of each daily mean by the cosine or sine of the angle indicated. The solution of the equations give the values of  $A'$ ,  $A''$ ,  $B'$ ,  $B''$ , etc., from which the corresponding values of quantities  $A$  and  $\alpha$ ,  $B$  and  $\beta$ , etc., of formula (405) are readily obtained, since

$$A = \sqrt{(A')^2 + (A'')^2} \text{ and } \alpha = \tan^{-1} \frac{-A''}{A'}.$$

In calculating the corrected epoch, it must be kept in mind that the  $t$  in this reduction is referred to the 11.5 hour of the first day of series instead of the preceding midnight.

**275.** Before solving equations (417), if the daily means have not already been cleared of the effects of the short-period constituents, it will be necessary to apply corrections to the first member of each of these equations in order to make the clearances.

The disturbance in a single daily mean due to the presence of a short-period constituent is represented by equation (398). Introducing the subscript  $s$  to distinguish the symbols pertaining to the short-period constituents, the disturbance in the daily mean of the  $n^{\text{th}}$  day of series due to the presence of the short-period constituent  $A_s$  may be written

$$[y_s]_n = \frac{1}{24} A_s \frac{\sin 12a_s}{\sin \frac{1}{2}a_s} \cos \{24(n-1)a_s + 11.5a_s + \alpha_s\} \quad (418)$$

The disturbances in the products of the daily means by

$$\cos 24(n-1)\alpha \text{ and } \sin 24(n-1)\alpha$$

may therefore be written

$$[y_s]_n \cos 24(n-1)\alpha$$

$$= \frac{1}{24} A_s \frac{\sin 12a_s}{\sin \frac{1}{2}a_s} \frac{1}{2} [\cos \{24(n-1)(a_s + a) + 11.5a_s + \alpha_s\} \\ + \cos \{24(n-1)(a_s - a) + 11.5a_s + \alpha_s\}] \quad (419)$$

and

$$\begin{aligned} [y_s]_n \sin 24(n-1)\alpha \\ = \frac{1}{24} A_s \frac{\sin 12a_s}{\sin \frac{1}{2}a_s} \frac{1}{2} [\sin \{24(n-1)(a_s + a) + 11.5a_s + \alpha_s\} \\ - \sin \{24(n-1)(a_s - a) + 11.5a_s + \alpha_s\}] \end{aligned} \quad (420)$$

**276.** Then, referring to formulas (263) and (264)

$$\sum_{n=1}^{365} [y_s]_n \cos 24(n-1)a = \frac{1}{48} A_s \frac{\sin 12a_s}{\sin \frac{1}{2}a_s} \left[ \frac{\sin 12 \times 365(a_s + a)}{\sin 12(a_s + a)} \cos \{12 \times 364(a_s + a) + 11.5a_s + \alpha_s\} + \frac{\sin 12 \times 365(a_s - a)}{\sin 12(a_s - a)} \cos \{12 \times 364(a_s - a) + 11.5a_s + \alpha_s\} \right] \quad (421)$$

and

$$\sum_{n=1}^{365} [y_s]_n \sin 24(n-1)a = \frac{1}{48} A_s \frac{\sin 12a_s}{\sin \frac{1}{2}a_s} \left[ \frac{\sin 12 \times 365(a_s + a)}{\sin 12(a_s + a)} \sin \{12 \times 364(a_s + a) + 11.5a_s + \alpha_s\} - \frac{\sin 12 \times 365(a_s - a)}{\sin 12(a_s - a)} \sin \{12 \times 364(a_s - a) + 11.5a_s + \alpha_s\} \right] \quad (422)$$

Now let

$$A'_s = A_s \cos \alpha_s \quad (423)$$

and

$$A''_s = -A_s \sin \alpha_s$$

then (421) and (422) may be reduced as follows:

$$\begin{aligned} & \sum_{n=1}^{365} [y_s]_n \cos 24(n-1)a \\ &= \frac{1}{48} \frac{\sin 12a_s}{\sin \frac{1}{2}a_s} \left[ \frac{\sin 12 \times 365(a_s + a)}{\sin 12(a_s + a)} \cos \{12 \times 364(a_s + a) + 11.5a_s\} \right. \\ & \quad \left. + \frac{\sin 12 \times 365(a_s - a)}{\sin 12(a_s - a)} \cos \{12 \times 364(a_s - a) + 11.5a_s\} \right] A'_s \\ &+ \frac{1}{48} \frac{\sin 12a_s}{\sin \frac{1}{2}a_s} \left[ \frac{\sin 12 \times 365(a_s + a)}{\sin 12(a_s + a)} \sin \{12 \times 364(a_s + a) + 11.5a_s\} \right. \\ & \quad \left. + \frac{\sin 12 \times 365(a_s - a)}{\sin 12(a_s - a)} \sin \{12 \times 364(a_s - a) + 11.5a_s\} \right] A''_s \end{aligned} \quad (424)$$

and

$$\begin{aligned} & \sum_{n=1}^{365} [y_s]_n \sin 24(n-1)a \\ &= \frac{1}{48} \frac{\sin 12a_s}{\sin \frac{1}{2}a_s} \left[ \frac{\sin 12 \times 365(a_s + a)}{\sin 12(a_s + a)} \sin \{12 \times 364(a_s + a) + 11.5a_s\} \right. \\ & \quad \left. - \frac{\sin 12 \times 365(a_s - a)}{\sin 12(a_s - a)} \sin \{12 \times 364(a_s - a) + 11.5a_s\} \right] A'_s \\ &- \frac{1}{48} \frac{\sin 12a_s}{\sin \frac{1}{2}a_s} \left[ \frac{\sin 12 \times 365(a_s + a)}{\sin 12(a_s + a)} \cos \{12 \times 364(a_s + a) + 11.5a_s\} \right. \\ & \quad \left. - \frac{\sin 12 \times 365(a_s - a)}{\sin 12(a_s - a)} \cos \{12 \times 364(a_s - a) + 11.5a_s\} \right] A''_s \end{aligned} \quad (425)$$

**277.** Formulas (424) and (425) represent the clearances for any long-period constituent  $A$  due to any short-period constituent  $A_s$ . The first must be subtracted from terms corresponding to  $\Sigma y_n \cos 24(n-1)a$  and the latter from terms corresponding to  $\Sigma y_n \sin 24(n-1)a$  of formula (417) before solving the latter.

**278.** In (424) and (425) the coefficients of  $A'_s$  and  $A''_s$ , which for brevity we may designate as  $C'$ ,  $C''$ ,  $S'$ , and  $S''$ , respectively, contain only values that are constant for all series and may therefore be computed once for all. Separate sets of such coefficients must, however, be computed for the effect of each short-period constituent upon each long-period constituent. In the usual reductions in which the effects of 3 short-period constituents upon 5 long-period constituents are considered, 15 sets of 4 coefficients each, or 60 coefficients in all, are required.

The coefficients are given in the following table: \*

	Long-period constituents				
	Mm	Mf	MSf	Sa	Ssa
M <sub>2</sub> (C')	-0.0556	+0.0030	+5.739	-0.1041	-0.1046
(C'')	-0.1704	-0.0377	-2.923	-0.0752	-0.0755
(S')	-0.1708	+0.0417	-2.840	-0.0018	-0.0035
(S'')	+0.0441	+0.0105	-5.727	+0.0048	+0.0096
N <sub>2</sub> (C')	-0.0588	+0.0368	+0.0294	-0.0176	-0.0176
(C'')	-0.0776	-0.2236	-0.1938	+0.0025	+0.0025
(S')	-0.0206	-0.1526	-0.1221	+0.0002	+0.0004
(S'')	+0.1138	-0.0854	-0.0808	+0.0001	+0.0002
O <sub>1</sub> (C')	-0.0648	+0.0166	+0.0157	-0.1924	-0.1934
(C'')	-0.3476	-0.0778	-0.0816	-0.1826	-0.1831
(S')	-0.3452	+0.0841	+0.0875	-0.0046	-0.0093
(S'')	+0.0405	+0.0338	+0.0331	+0.0090	+0.0180

In the above table the sign is so taken that the values are to be applied to the sums directly as indicated.

**279.** After the clearances have been applied and the normal equations (417) solved and the resulting amplitude and epoch obtained for each of the long-period constituents, the reductions will be completed in accordance with the processes already outlined, but it must be kept in mind that in this reduction the initial value of  $t$  is taken to correspond to 11:30 a. m. on the first day of series. In obtaining the numerical values of such quantities as  $\Sigma y_n \cos 24(n-1)a$  and  $\Sigma y_n \sin 24(n-1)a$ , in order to avoid the labor of separate multiplications for each day, the following abbreviations have been proposed by the British authorities. The values of  $\cos 24(n-1)a$  and of  $\sin 24(n-1)a$  are divided into 11 groups according as they fall nearest 0, 0.1, 0.2, 0.3, 0.4, 0.5, 0.6, 0.7, 0.8, 0.9, or 1.0. The daily values are then distributed into 11 corresponding groups, so that all values in one group will be multiplied by 0, another group by 0.1, etc. The  $\cos 24(n-1)a$  and  $\sin 24(n-1)a$  include negative as well as positive values. The former are taken into account by changing the sign of the daily mean to which the negative values apply.

**280.** As a part of the routine reductions of the tidal records from the principal tide stations it is the practice of the office to obtain the mean sea level for each calendar month. It is therefore desirable to

\*From Scientific Papers by Sir George H. Darwin, Vol. I, p. 64.

have a method of using these means directly in the analysis for the annual and semiannual constituents, thus avoiding any special summation for the purpose. The period of the annual constituent is approximately the length of the Julian year, that is, 365.25 days. If this period is divided into 12 equal groups and the mean of the hourly heights for each group taken, these means represent the approximate height of the combined annual and semiannual constituents for the middle of each group, and the middle of the first group will be the initial point from which the zeta ( $\zeta$ ) as obtained by the usual process is referred. As each group represents  $30^\circ$  of motion for the annual constituent, or  $60^\circ$  for the semiannual constituent, to refer this  $\zeta$  to the actual beginning of the series of observations it will be necessary to apply a correction of  $15^\circ$  for the annual constituent or  $30^\circ$  for the semiannual constituent.

**281.** In obtaining the monthly means by calendar months the year is divided only approximately into 12 equal groups. The following table shows the difference between the middle of each group representing a calendar month and the middle of the corresponding group obtained by dividing the Julian year into 12 equal parts. It is to be noted that the hourly heights included in a monthly sum extend from 0 hour on the first day of the month to the 23d hour on the last day. The middle of the group as reckoned from the beginning of the month will therefore be 13.98 days, 14.48 days, 14.98 days, or 15.48 days, respectively, according to whether the month has 28, 29, 30, or 31 days.

Month	Middle of group reckoned from beginning of year			Differences	
	Julian year	Common year	Leap year	Common year	Leap year
January	15.22	15.48	15.48	+0.26	+0.26
February	45.66	44.98	45.48	-0.68	-0.18
March	76.09	74.48	75.48	-1.61	-0.61
April	106.53	104.98	105.98	-1.55	-0.55
May	136.97	135.48	136.48	-1.49	-0.49
June	167.41	165.98	166.98	-1.43	-0.43
July	197.84	196.48	197.48	-1.36	-0.36
August	228.28	227.48	228.48	-0.80	+0.20
September	258.72	257.98	258.98	-0.74	+0.26
October	289.16	288.48	289.48	-0.68	+0.32
November	319.59	318.98	319.98	-0.61	+0.39
December	350.03	349.48	350.48	-0.55	+0.45
Sums				-11.24	-0.74
Means				-0.94	-0.06
Speed of Sa constituent per day = 0.9850°.				°	°
Mean differences reduced to degrees of Sa.				-0.93	-0.06
Correction to $\zeta$ of Sa.				14.07	14.94
Correction to $\zeta$ of Ssa.				28.14	29.88

**282.** From the above table it is evident that in the summation for the monthly means for a calendar year the middle of each group of a common year is on an average  $0.93^\circ$  earlier than the middle of the corresponding group when the Julian year is equally subdivided and the middle of each group of a leap year is on an average  $0.06^\circ$  earlier. Subtracting these values from  $15^\circ$ , the interval between the beginning of the observations and the middle of the first group of an equal subdivision, we have  $14.07^\circ$  and  $14.94^\circ$ , for common and leap years, respectively, as a correction to be applied to the  $\zeta$  of Sa as

directly obtained, in order to refer the  $\zeta$  to the 0 hour of the 1st day of January. For Ssa the corrections will be twice as great.

283. If the year commences on the first day of any month other than January, the corrections will differ a little from the above. Calculated in a manner similar to that above, the following table gives the correction to be applied to the  $\zeta$  to refer to the first day of any month at which the series commences. The correction to the  $\zeta$  of Ssa will be twice the tabular value for Sa.

Observations commence—	Correction to $\zeta$ of Sa to refer to begin- ning of month		Observations commence—	Correction to $\zeta$ of Sa to refer to begin- ning of month	
	Common year	Leap year		Common year	Leap year
Jan. 1	◦	◦	July 1	◦	◦
Feb. 1	14.07	14.94	Aug. 1	15.56	15.93
Mar. 1	13.50	14.45	Sept. 1	14.98	15.43
Apr. 1	15.89	15.93	Oct. 1	14.41	14.94
May 1	15.31	15.43	Nov. 1	14.82	15.43
June 1	15.72	15.93	Dec. 1	14.24	14.94
	15.15	15.43		14.65	15.43

284. If the monthly means extend over many calendar years, it may be convenient to combine them for a single analysis. In this case the ( $V_0 + u$ ) for January 1 may be taken as the average of the values for the beginning of each year included in the observations, and the correction to the  $\zeta$  to refer to the beginning of the year will be a mean of the values given above for common and leap years, weighted in accordance with the number of each kind of year included. If only a few years of observations are available, it is better to analyze each year separately in order that the results may serve as a check on each other.

285. The augmenting factors to be used for constituents Sa and Ssa when derived from the monthly sea level values are based upon formula (404) in paragraph 266 and are as follows:

Sa 1.0115, logarithm 0.00497.

Ssa 1.0472, logarithm 0.02003.

#### ANALYSIS OF HIGH AND LOW WATERS

286. The automatic tide gage, which furnishes a continuous record of the rise and fall of the tide, now being in general use, it is seldom necessary to rely only upon the high and low waters for an analysis. It may happen, however, that a record of high and low water observations is available for a more or less isolated locality where it has been impractical to secure continuous records. Such records, if they include all the high and low waters for a month or more may be utilized in determining approximate values of the principal harmonic constants, but the results are not as satisfactory as those obtained from an analysis of the hourly heights.

287. An elaborate mode of analysis of the high and low waters is contained in volume 1 of Scientific Papers, by Sir George H. Darwin. Other methods are given by Dr. R. A. Harris in his Manual of Tides. The process outlined below follows to some extent one of the methods of Doctor Harris, extending his treatment for the K and O to other constituents.

**288.** The lengths of series may be taken the same as the lengths used as the analysis of the hourly heights (see par. 152). It is sometimes convenient to divide a series, whatever its length, into periods of 29 days each. This permits a uniform method of procedure, and a comparison of the results from different series affords a check on the reliability of the work.

**289.** The first process in this analysis consists in making the usual high and low water reductions, including the computation of the lunitidal intervals. Form 138 provides for this reduction. The times and heights of the high and low waters, together with the times of the moon's transits, are tabulated. For convenience the standard time of the place of observations may be used for the times of the high and low waters, and the Greenwich mean civil time of the moon's transits over the meridian of Greenwich may be used for the moon's transits. The interval between each transit and the following high and low water is then found, and the mean of all the high water intervals and the mean of all the low water intervals are then obtained separately. The true mean intervals between the time of the moon's transit over the local meridian and the time of the following high and low waters being desired, the means as directly obtained must be corrected to allow for any difference in the kind of time used for the transit of the moon and the time of the tides and also for the difference in time between the transit of the moon over the local meridian and the transit over the meridian to which the tabular values refer.

**290.** If the tide is of the semidiurnal type, the approximate amplitude and epoch for  $M_2$  may be obtained directly from this high and low water reduction. On account of the presence of the other constituents the mean range from the high and low waters will always be a little larger than twice the amplitude of  $M_2$ . If the data are available for some other station in the general locality, the ratio of the  $M_2$  amplitude to the mean range of tide at that station may be used in finding the  $M_2$  amplitude from the mean range of tide at the station for which the results are sought. If this ratio cannot be obtained for any station in the general locality, the empirical ratio of 0.47 may be used with fairly satisfactory results. After the amplitude of  $M_2$  has been thus obtained, it should be corrected for the longitude of the moon's node by factor  $F$  from table 12.

**291.** The epoch of  $M_2$  may be obtained from the corrected high and low water lunitidal intervals  $HWI$ ,  $LWI$  by the following formula:

$$M^{\circ}_2 = \frac{1}{2}(HWI + LWI) \times 28.984 + 90^{\circ} \quad (426)$$

In the above formula  $HWI$  must be greater than  $LWI$ , 12.42 hours being added, if necessary, to the  $HWI$  as directly obtained from the high and low water reductions.

**292.** The difference between the average duration of rise and fall of the tide at any place, where the tide is of the semidiurnal type, depends largely upon the constituent  $M_4$ . It is possible to obtain from the high and low waters a constituent with the speed of  $M_4$  which, when used in the harmonic prediction of the tides, will cause the mean duration of rise and fall to be the same as that at the station. The effect of  $M_4$  upon the mean duration of rise will depend chiefly upon the relation of its amplitude and epoch to the amplitude and epoch of the principal constituent  $M_2$ . By assuming an  $M_4$  with epoch

such as to make the constituent symmetrically situated in regard to the maxima and minima of  $M_2$ , the amplitude necessary to account for the mean duration of rise of the tide may be readily calculated.

**293.** Let  $DR$ =duration of rise of tide in hours as obtained from the lunital intervals,

$$a = \text{Hourly speed of } M_2 = 28.^{\circ}984.$$

$$M_2 = \text{Amplitude of } M_2.$$

$$M_2^{\circ} = \text{Epoch of } M_2.$$

$$M_4 = \text{Amplitude of } M_4.$$

$$M_4^{\circ} = \text{Epoch of } M_4.$$

Then, for  $M_4$  to be symmetrically situated with respect to the maxima and minima of  $M_2$

$$M_4^{\circ} = 2 M_2^{\circ} \pm 90^{\circ} \quad (427)$$

in which the upper or lower sign is to be used according to whether  $a(DR)$  is greater or less, respectively, than  $180^{\circ}$ . Multiples of  $360^{\circ}$  may be added or rejected to obtain the result as a positive angle less than  $360^{\circ}$ .

The equations of the constituents  $M_2$  and  $M_4$  may be written

$$y_1 = M_2 \cos (at + \alpha) \quad (428)$$

$$y_2 = M_4 \cos (2at + \beta) \quad (429)$$

and the resultant curve

$$y = M_2 \cos (at + \alpha) + M_4 \cos (2at + \beta) \quad (430)$$

**294.** Values of  $t$  which will render (428) a maximum must satisfy the derived equation

$$aM_2 \sin (at + \alpha) = 0 \quad (431)$$

and for a maximum of (430)  $t$  must satisfy the derived equation

$$aM_2 \sin (at + \alpha) + 2aM_4 \sin (2at + \beta) = 0 \quad (432)$$

For a maximum of (428)

$$t = \frac{2n\pi - \alpha}{a} \quad (433)$$

in which  $n$  is any integer.

**295.** Let  $\frac{\theta}{a}$  = the acceleration in the high waters of  $M_2$  due to the presence of  $M_4$ . With the  $M_4$  wave symmetrically situated with respect to the  $M_2$  wave,  $\frac{\theta}{a}$  will also equal the retardation in the low water of  $M_2$ , due to the presence of  $M_4$ , and  $\frac{2\theta}{a}$  will equal the total amount by which the duration of rise of the tide has been diminished by  $M_4$ . If the duration of rise has been increased,  $\theta$  will be negative. Then, for a maximum of (430)

$$t = \frac{2n\pi - \alpha - \theta}{a} \quad (434)$$

and this value of  $t$  must satisfy equation (432).

**296.** Substituting in (432), we have

$$\begin{aligned} aM_2 \sin (2n\pi - \theta) + 2aM_4 \sin (4n\pi - 2\alpha + \beta - 2\theta) = \\ -aM_2 \sin \theta - 2aM_4 \sin (2\theta + 2\alpha - \beta) = 0 \end{aligned} \quad (435)$$

But

$$2\alpha - \beta = -2M_2^\circ + M_4^\circ \quad (436)$$

From (427)

$$-2M_2^\circ + M_4^\circ = \pm 90^\circ$$

according to whether the duration of rise is greater or less than  $\frac{180^\circ}{a}$ , or whether  $\theta$  is negative or positive.

Then

$$2\alpha - \beta = \mp 90^\circ \quad (437)$$

according to whether  $\theta$  is positive or negative.

Substituting this in (435)

$$-aM_2 \sin \theta \pm 2aM_4 \cos 2\theta = 0 \quad (438)$$

and

$$\frac{M_4}{M_2} = \pm \frac{1}{2} \frac{\sin \theta}{\cos 2\theta} \quad (439)$$

the upper or lower sign being used according to whether  $\theta$  is positive or negative. As under the assumed conditions  $\theta$  must come within the limits  $\pm 45^\circ$ , the ratio of  $\frac{M_4}{M_2}$  as derived from (439) will always be positive.

**297.** The duration of rise of tide due solely to the constituent  $M_2$  is  $\frac{180^\circ}{a}$ .

The duration of rise as modified by the presence of the assumed  $M_4$  is

$$DR = \frac{180^\circ}{a} - \frac{2\theta}{a} \quad (440)$$

Therefore

$$\theta = \frac{1}{2}(180^\circ - aDR) \quad (441)$$

Substituting the above in (439) we have

$$\frac{M_4}{M_2} = \pm \frac{1}{2} \frac{\sin (90^\circ - \frac{1}{2}aDR)}{\cos (180^\circ - aDR)} = \mp \frac{1}{2} \frac{\cos \frac{1}{2}aDR}{\cos aDR} \quad (442)$$

and

$$M_4 = \mp \frac{1}{2} \frac{\cos \frac{1}{2}aDR}{\cos aDR} M_2 \quad (443)$$

$M_4$  must be positive, and the sign of the above coefficient will depend upon whether  $aDR$  is less or greater than  $180^\circ$ .

**298.** The approximate constants for  $S_2$ ,  $N_2$ ,  $K_1$ , and  $O_1$  may be obtained from the observed high and low waters as follows: Add to each low-water height the mean range of tide. Copy the high and modified low water heights into the form for hourly heights (form 362), always putting the values upon the nearest solar hour. Sum for the desired constituents, using the same stencils as are used for the regular

analysis of the hourly heights. Account should be taken of the number of items entering into each sum and the mean for each constituent hour obtained. The 24 hourly means for each constituent are then to be analyzed in the usual manner.

**299.** The results obtained by this process are, of course, not as dependable as those obtained from a continuous record of hourly heights. The approximate results first obtained can, however, be improved by the following treatment if a tide-computing machine is available. Using the approximate constants as determined above for the principal constituents and inferred values for smaller constituents, set the machine for the beginning of the period of observations and find the predicted heights corresponding to the observed times of the high and low waters. Tabulate the differences between the observed and predicted heights for these times, using the hourly height form and entering the values according to the nearest solar hour. These differences are then to be summed and analyzed the same as the original observed heights. In this analysis of the residuals the constituent  $M_2$  should be included. The results from the analysis of the residuals are then combined with the constants used for the setting of the predicting machine.

**300.** In making the combinations the following formulas may be used:

Let  $A'$  and  $\kappa'$  represent the first approximate values of the constants of any constituent.

$A''$  and  $\kappa''$ , the constants as obtained from the residuals.

$A$  and  $\kappa$ , the resultant constants sought.

Then

$$A = \sqrt{(A' \cos \kappa' + A'' \cos \kappa'')^2 + (A' \sin \kappa' + A'' \sin \kappa'')^2} \quad (444)$$

and

$$\kappa = \tan^{-1} \frac{A' \sin \kappa' + A'' \sin \kappa''}{A' \cos \kappa' + A'' \cos \kappa''} \quad (445)$$

#### FORMS USED FOR ANALYSIS OF TIDES

**301.** Forms used by the Coast and Geodetic Survey for the harmonic analysis of tide observations are shown in figures 9 to 19. A series of tide observations at Morro, California, covering the period February 13 to July 25, 1919, is taken as an example to illustrate the detail of the work.

**302.** *Form 362, Hourly heights (fig. 9).*—The hourly heights of the tide are first tabulated in form 362. Although the zero of the tide staff is usually taken as the height datum, any other fixed plane will serve this purpose. For practical convenience it is desirable that the datum be low enough to avoid negative tabulations but not so low as to cause the readings to be inconveniently large for summing.

**303.** The hours refer to mean solar time, which may be either local or standard, astronomical or civil, but standard civil time will generally be the most convenient to use. The series must commence with the zero (0) hour of the adopted time, and all vacancies in the record should be filled by interpolated values in order that each hour of the series may be represented by a tabulated height. It is the general practice to use brackets with interpolated values to distinguish them from the observed heights. The record for successive days of the series must be entered in successive columns of the form, and these

columns are to be numbered consecutively, beginning with one (1) for the first day of the series.

**304.** The series analyzed should be one of the lengths indicated in paragraph 152. Series of observations very nearly equal to one of these standard lengths may be completed by the use of extrapolated hourly heights. If the observations cover a period of several years, the analysis for each year may be made separately, a comparison of the results affording an excellent check on the work.

**305.** The hourly heights on each page of form 362 are first summed horizontally and vertically. The total of the vertical sums must equal the total of the horizontal sums, and this page sum is entered in the lower right-hand corner of the page.

Form 362 DEPARTMENT OF COMMERCE U. S. COAST AND GEODETIC SURVEY										TIDES: HOURLY HEIGHTS	
Station: Morro, California. Chief of Party: E. B. Lathan. Time Meridian: 120° W. Year: 1919. Lat. 35° 22' N. Long. 120° 51' W. Tide Gauge No. 107 Scale 1:9 Reduced to Staff.										11-712	
Month and Day.	mo.	d. Feb. 13	d. 14	d. 15	d. 16	d. 17	d. 18	d. 19	d. 20	Horizontal Sum.	
Day of Series.	1	2	3	4	5	6	7	8	9	10	
Hour.	Feet.	Feet.	Feet.	Feet.	Feet.	Feet.	Feet.	Feet.	Feet.	Feet.	
0	3.9	4.2	4.6	4.5	4.4	4.7	4.6	4.6	4.6	30.9	
1	3.4	3.8	4.2	4.2	4.2	4.9	4.8	4.8	4.8	29.5	
2	3.0	3.3	3.5	3.7	3.8	4.6	4.9	4.9	4.9	26.8	
3	2.8	3.0	3.0	3.1	3.3	4.1	4.5	4.5	4.5	23.8	
4	3.0	2.8	2.6	2.5	2.7	3.5	3.8	3.8	3.8	20.9	
5	3.6	3.1	2.5	2.2	2.2	3.0	3.2	3.2	3.2	19.8	
6	4.4	3.6	2.8	2.2	1.9	2.6	2.7	2.7	2.7	20.2	
7	5.1	4.5	3.5	2.6	2.0	2.5	2.3	2.3	2.3	22.5	
8	5.7	5.3	4.3	3.3	2.4	2.7	2.2	2.2	2.2	25.9	
9	6.0	6.0	4.9	4.1	3.1	3.1	2.4	2.4	2.4	29.6	
10	5.6	6.2	5.4	4.6	3.9	3.6	2.8	2.8	2.8	32.1	
11	4.8	5.8	5.5	4.9	4.3	4.1	3.2	3.2	3.2	32.6	
Noon.	3.9	5.1	5.1	4.8	4.4	4.5	3.6	3.6	3.6	31.4	
13	3.4	4.3	4.4	4.3	4.2	4.5	3.8	3.8	3.8	28.9	
14	2.6	3.4	3.5	3.6	3.7	4.3	3.8	3.8	3.8	24.9	
15	1.9	2.6	2.8	2.9	3.1	3.8	3.6	3.6	3.6	20.7	
16	1.2	2.0	2.2	2.2	2.6	3.2	3.2	3.2	3.2	16.5	
17	1.0	1.6	1.7	1.6	2.1	2.7	2.8	2.8	2.8	13.5	
18	1.3	1.6	1.5	1.3	1.9	2.4	2.5	2.5	2.5	12.5	
19	2.3	2.2	1.8	1.4	1.9	2.3	2.3	2.3	2.3	14.2	
20	3.2	3.1	2.6	2.0	2.3	2.5	2.4	2.4	2.4	18.1	
21	4.0	3.9	3.4	2.8	3.0	3.0	2.9	2.9	2.9	23.0	
22	4.3	4.5	4.1	3.6	3.8	3.6	3.7	3.7	3.7	27.6	
23	4.5	4.7	4.5	4.1	4.4	4.2	4.2	4.2	4.2	30.6	
Sum.	84.9	90.6	84.4	76.5	75.6	84.4	80.2	80.2	80.2	876.6	
Sum for 29 days, 1 to 29 of										Divisor=696; mean for 29 days=	

FIGURE 9.

**306.** *Stencils* (figs. 10 and 11).—The first figure is a copy of the M stencil for the even hours of the first 7 days of the series, and the second figure illustrates the application of the same. This stencil being laid over the page of hourly heights shown in figure 9, the heights applying to each of the even constituent hours for this page show through the openings in the stencil, where they appear connected by diagonal lines, thus indicating each group to be summed.

**307.** For each constituent summation, excepting for S, there are provided two stencils for each page of tabulated hourly heights, one for the even constituent hours and the other for the odd constituent hours.

TIDES: HOURLY HEIGHTS											
Form 868 DEPARTMENT OF COMMERCE U. S. COAST AND GEODETIC SURVEY											
Station: <u>Stencil for component M.</u> Year: _____											
Chief of Party: _____ Lat. _____ Long. _____											
Time Meridian: _____ Tide Gauge No. _____ Scale 1: _____ Reduced to Staff. _____											
11-792											
Month and Day.	mo.	d.	d.	d.	d.	d.	d.	d.	d.	d.	Hor- izon- tal Sum.
Day of Series	1	2	3	4	5	6	7				
Hour.	Feet.				Feet.						
9	0		22			5		20			.
1											.
2	2							22			.
3											.
4	4							0			.
5											.
6	6							2			.
7											.
8	8							4			.
9											.
10	10							6			.
11											.
Noon	12							8			.
13											.
14	14										.
15								10			.
16											.
17	16							12			.
18											.
19	18							14			.
20											.
21	20							16			.
22	0										.
23	22			20				18			.
Sum.											
Sum for 29 days, 1 to 29 of _____ = Divisor = 696; mean for 29 days = _____											

FIGURE 10.

The stencils are numbered with the days of series to which they apply, and special care must be taken to see that the days of series on each stencil correspond with the days of series on the page of tabulations with which it is used. For constituent S no stencils are necessary, as the constituent hours correspond to the solar hours of the tabulations and the horizontal sums from form 362 may be taken directly as the constituent hour sums.

**308. Form 142, Stencil sums (figs. 12 and 13).**—The sums for each constituent hour are entered in form 142, one line of the form being used for each page of the original tabulations. The total of the hour

TIDES: HOURLY HEIGHTS										
Form 362 DEPARTMENT OF COMMERCE U. S. COAST AND GEODESIC SURVEY										
Station: <u>Stencil for component M.</u>	Year: _____									
Chief of Party: _____	Lat. _____ Long. _____									
Time Meridian: _____ Tide Gauge No. _____ Scale 1: _____ Reduced to Staff. _____										
Month and Day.	mo.	d.	Horizontal Sum.							
Day of Series	1	2	3	4	5	6	7			
Hour.	Feet.	Feet.	Feet.	Feet.	Feet.	Feet.	Feet.	Feet.		
0	3.9	0	4.6	2.2	4.5	.	4.7	2.0	.	
1	.	3.8	.	.	4.2	.	4.6	2.2	.	
2	3.0	2	3.5	.	.	.	4.6	2.2	.	
3	.	3.0	.	3.1	3.3	.	4.5	.	.	
4	3.0	4	2.6	.	.	.	3.5	0	.	
5	.	3.1	.	2.2	2.2	.	3.2	.	.	
6	4.4	6	2.8	.	.	2.6	2	.	.	
7	.	4.5	.	2.6	2.0	.	2.3	.	.	
8	5.7	8	4.3	.	2.4	2.7	4	.	.	
9	.	6.0	.	4.1	.	.	2.4	.	.	
10	5.6	10	5.4	.	3.9	3.6	6	.	.	
11	.	5.8	.	4.9	.	.	3.2	.	.	
Noon.	3.9	12	5.1	.	4.4	4.5	8	.	.	
13	.	4.3	.	4.3	.	.	3.8	.	.	
14	2.6	14	3.5	.	3.7	.	.	.	.	
15	1.9	2.6	.	2.9	.	3.8	10	3.6	.	
16	.	.	2.2	.	2.6	.	.	.	.	
17	1.0	16	1.6	.	1.6	.	2.7	12	2.8	.
18	.	.	1.5	.	1.9	.	.	.	.	
19	2.3	18	2.2	.	1.4	.	2.3	14	.	
20	.	.	2.6	.	2.3	.	2.4	.	.	
21	4.0	20	.	2.8	.	3.0	16	.	.	
22	.	4.5	4.1	.	3.8	.	3.7	.	.	
23	4.5	22	.	4.1	2.0	.	4.2	18	.	
Sum.	.	.	.	.	.	.	.	.	.	
Sum for 29 days, 1 to 29 of										
Divisor=696; mean for 29 days=										

FIGURE 11.

sums in each line of the form must equal the corresponding page sum of the hourly heights in form 362, this serving as a check on the summation. After the summing of all the pages of the series has been completed for any constituent the totals for each constituent hour are obtained, the divisors from table 32 entered, and the constituent hourly means computed (fig. 13). These means should be carefully checked before proceeding with the analysis. Large errors can usually be detected by plotting the means.

**309. Form 244, Computation of  $(V_0 + u)$**  (fig. 14).—This form provides for the computation of the equilibrium arguments for the beginning of the series of observations, the computation being in accordance with formulas given in table 2. For the most part the form is self-explanatory. The values of the mean longitude of the

FORM 244  
DEPARTMENT OF COMMERCE  
U. S. COAST AND GEODETIC SURVEY

TIDES: STENCIL SUMS.

Station: Morro, California. Lat.:  $35^{\circ} 22' N.$

Component: "M". Length of series: 163. Series begins: 1919-Feb-13-0. Long:  $120^{\circ} 51' W.$

Kind of time used:  $120^{\circ} W.$  Computed by Fred A. Kummell, Dec. 9, 1920.

11-47

Page.	0	1	2	3	4	5	6	7	8	9	10	11	
1	24.3	20.6	17.9	16.9	21.0	23.0	28.0	31.9	39.2	34.8	31.9	27.4	
2	21.8	17.5	14.4	13.6	11.2	12.3	14.6	20.5	21.7	23.9	24.7	24.3	
3	19.7	16.9	11.0	9.6	12.2	17.7	28.1	24.9	27.4	27.6	29.8	21.1	
4	26.4	18.0	17.3	17.3	22.7	22.6	26.0	34.3	36.5	41.2	33.3	28.5	
5	21.5	21.4	17.9	16.2	16.3	19.9	24.9	29.9	37.7	34.8	33.2	29.6	
6	20.3	16.8	15.8	12.1	12.6	15.1	21.0	21.4	23.6	24.8	25.0	28.1	
7	23.1	16.1	13.3	13.1	15.6	23.7	28.6	33.9	30.1	34.9	27.6	23.6	
8	25.5	23.0	21.4	21.6	20.8	23.5	27.2	29.7	43.5	36.4	32.6	27.5	
9	20.9	18.5	15.2	12.1	11.3	13.8	18.1	26.3	26.8	28.6	28.0	29.6	
10	16.9	13.2	10.2	8.7	11.5	15.5	18.3	21.5	24.4	25.3	28.7	24.3	
11	18.6	15.0	12.5	12.5	17.7	17.2	23.2	29.5	41.1	36.7	34.4	27.8	24.0
12	24.6	25.5	20.4	20.7	21.0	24.6	32.1	31.7	32.7	36.5	31.6	25.6	
13	25.7	17.3	13.2	10.0	11.9	12.5	16.2	20.3	24.3	30.2	30.3	24.4	
14	16.7	12.6	8.3	8.7	9.5	14.3	19.0	23.4	30.2	27.4	27.2	26.2	
15	19.0	16.1	16.3	15.7	20.1	26.5	37.7	37.7	40.2	39.3	35.6	29.6	
16	29.6	22.8	21.6	22.5	25.7	31.7	31.9	34.9	35.8	38.1	28.3	27.3	
17	22.9	18.6	14.5	11.1	10.5	12.3	15.3	19.0	25.4	24.5	24.9	23.6	
18	15.4	10.0	6.2	3.2	4.9	10.2	16.0	24.2	24.5	25.1	25.4	27.6	
19	16.7	15.4	13.1	15.3	19.8	29.4	31.8	35.8	38.3	37.9	38.7	28.3	
20	27.6	21.0	19.8	20.4	28.1	29.9	31.5	36.1	36.4	39.9	28.9	22.8	
<b>Sum:</b> 437.1 356.3 300.3 <b>286.5</b> 323.8 401.7 495.8 578.5 635.4 645.6 593.5 523.4													
Page.	12	13	14	15	16	17	18	19	20	21	22	23	
1	22.7	21.1	17.5	13.5	14.5	17.5	17.9	26.2	26.2	27.7	26.9	28.0	576.6
2	25.8	17.6	17.4	18.9	21.5	28.8	32.1	35.1	36.5	35.9	35.8	26.6	552.5
3	17.5	17.5	14.7	15.4	21.1	23.5	29.2	33.3	35.3	39.5	30.0	24.8	547.8
4	23.2	20.8	12.9	9.0	7.1	7.9	14.0	20.8	22.2	24.9	25.8	25.3	638.0
5	27.5	20.2	16.9	15.5	16.4	23.8	25.3	30.3	27.9	29.0	34.0	25.4	897.5
6	23.8	23.0	20.0	24.2	25.1	25.2	27.5	28.8	39.4	38.2	28.7	24.4	562.8
7	19.5	15.5	17.8	14.6	15.5	20.2	29.8	31.4	33.7	33.1	30.6	29.0	574.3
8	22.4	19.4	12.3	8.8	7.4	10.7	14.9	19.5	24.1	26.6	32.0	27.3	558.1
9	22.1	18.1	15.4	14.0	17.1	19.1	24.3	29.3	38.0	29.0	28.0	24.6	528.2
10	23.2	22.6	26.3	22.0	24.4	25.9	28.7	33.8	28.1	30.7	26.4	24.5	536.1
11	19.4	18.2	11.8	12.5	13.1	16.8	25.3	26.6	29.1	29.0	30.4	22.9	550.8
12	21.5	13.3	8.4	4.6	4.4	8.7	15.0	21.0	25.5	31.9	28.1	27.0	536.3
13	21.7	18.8	16.5	17.5	17.6	21.6	26.9	36.1	35.5	36.4	30.2	27.0	542.2
14	28.6	23.6	27.4	26.4	26.0	28.8	36.8	32.5	31.5	28.2	28.0	25.0	566.3
15	20.3	16.4	12.9	10.5	14.6	21.5	22.0	25.7	27.8	31.1	25.4	22.6	584.6
16	20.9	14.9	10.9	6.1	4.8	9.9	15.9	23.2	32.1	30.8	30.6	28.9	580.2
17	21.3	22.2	17.1	16.8	17.9	26.7	27.6	36.0	33.5	36.5	35.4	32.1	545.7
18	22.2	21.1	21.5	23.6	31.2	31.8	30.4	31.7	31.3	28.2	27.6	21.1	514.4
19	19.6	15.5	12.5	11.3	10.0	12.2	19.9	26.1	28.6	24.1	22.5	19.7	542.5
20	20.4	15.2	7.8	3.6	3.0	7.2	17.1	22.2	27.4	29.8	34.3	27.7	558.1
<b>Sum:</b> 443.6 375.0 318.0 286.9 310.7 386.8 491.6 569.6 613.7 620.6 590.7 513.9 11093.0													

FIGURE 12.

moon ( $s$ ), of the lunar perigee ( $p$ ), of the sun ( $h$ ), of the solar perigee ( $p_1$ ), and of the moon's ascending node ( $N$ ), may be obtained from table 4 for the beginning of any year between 1800 and 2000. The values for any year beyond these limits may be readily obtained by taking into account the rate of change in these elements as given in table 1. The corrections necessary in order to refer the elements to any desired month, day, and hour are given in table 5. As the tables refer to Greenwich mean civil time, the argument used in entering them should refer also to this kind of time, and in the lines for the beginning and middle of the series at the head of the form space is therefore provided for entering the equivalent Greenwich hour. Any change in the day may be avoided by using a negative Greenwich hour when necessary. For example, 1922, January 1, 0 hour, in the standard time of the meridian  $15^{\circ}$  east of Greenwich, may be written as 1922, January 1,  $-1$  hour in Greenwich time, instead of 1921, December 31, 23 hour, as would otherwise be necessary. If a negative argument is used in table 5, the corresponding tabular value must be taken with its sign reversed. For the middle of the series the nearest integral hour is sufficient.

**310.** The values of  $I$ ,  $\nu$ ,  $\xi$ ,  $\nu'$ , and  $2\nu''$  are obtained for the middle of the series from table 6, using  $N$  as the argument. If  $N$  is between  $180^{\circ}$  and  $360^{\circ}$ , each of the last four quantities will be negative, but  $I$

Form 148  
DEPARTMENT OF COMMERCE  
U. S. COAST AND GEODETIC SURVEY

## TIDES: STENCIL SUMS.

Station: Morro, California. Lat.:  $35^{\circ} 22' N.$   
Component:  $T^M$ . Length of series: 163. Series begins: 1919 - Feb. 13-0. Long.:  $120^{\circ} 51' W.$   
Dep. Kind of time used: 120 $^{\circ}$  W. Computed by Fred A. Kummell, Dec. 9, 1920.  
11-247 Date:

Page	0	1	2	3	4	5	6	7	8	9	10	11
21	25.9	18.1	14.8	14.5	10.6	11.1	17.3	23.8	23.1	24.4	24.1	22.6
22	16.8	14.8	7.7	5.7	6.6	11.1	19.5	23.2	26.5	27.6	30.8	24.9
23	17.8	15.7	15.1	20.1	21.6	30.7	33.3	37.3	39.0	42.8	33.9	28.4
24	7.2	6.8	6.2	6.1	6.5	8.0	9.7	10.9	18.3	12.1	11.0	9.4
Sums-21-24	67.7	55.4	43.8	46.4	45.5	60.9	79.8	95.2	106.9	106.9	99.8	85.3
" 1-20	437.1	356.3	300.3	286.5	323.8	401.7	495.8	578.5	635.4	645.6	593.5	523.4
Sums.-	504.8	411.7	344.1	332.9	369.1	462.6	575.6	673.7	742.3	752.5	693.3	608.7
Divisors.-	164	163	162	165	164	163	163	164	165	165	163	162
Means.-	3.08	2.53	2.12	2.02	2.25	2.84	3.53	4.13	4.53	4.56	4.25	3.76

Page	12	13	14	15	16	17	18	19	20	21	22	23	
21	23.3	18.2	17.0	17.3	23.3	24.0	29.7	32.9	35.9	42.1	34.7	31.1	558.8
22	22.5	20.7	20.2	26.0	26.9	31.7	36.2	34.0	40.0	31.3	26.2	20.5	551.4
23	23.1	16.3	15.5	11.6	11.9	13.7	19.6	25.1	26.6	26.2	24.0	24.5	573.8
24	3.3	4.7	3.0	1.7	0.9	0.9	1.7	3.4	5.5	7.0	7.7	7.8	159.8
Sums 21-24	72.2	59.9	55.7	56.6	63.0	70.3	86.2	95.4	108.1	106.6	92.6	83.9	1843.8
" 1-20	443.6	375.0	318.0	286.9	310.7	366.8	481.6	569.6	613.7	620.6	590.7	513.9	11093.0
Sums.-	515.8	434.9	373.7	345.5	373.7	459.1	567.8	665.0	721.7	727.2	683.3	597.8	12936.8
Divisors.-	162	163	163	163	162	162	163	163	162	162	163	163	
Means.-	3.18	2.67	2.29	2.12	2.31	2.83	3.48	4.08	4.45	4.49	4.19	3.67	

FIGURE 13.

is always positive. Although table 6 is computed for the epoch, January 1, 1900, it is applicable without material error for any series of observations.

311. The values of  $u$  of  $L_2$  and  $u$  of  $M_1$ , may be obtained from table 13 for any date between 1900 and 2000, inclusive, using the value of  $N$  for interpolation. If the series falls beyond the limits of this table, the following formulas may be used:

$$u \text{ of } L_2 = 2\xi - 2\nu - R \text{ (par. 129)} \quad (446)$$

$$u \text{ of } M_1 = \xi - \nu + Q \text{ (par. 123)} \quad (447)$$

The values of  $\xi$  and  $\nu$  may be taken from form 244, the values of  $R$  and  $Q$  from tables 8 and 10, respectively, using the arguments  $I$  and  $P$  for the middle of the series.

Form 244.  
DEPARTMENT OF COMMERCE,  
U. S. COAST AND GEODETIC SURVEY.

### TIDES: Computation of $V_e + u$ .

Station	Morro, California			Lat.	35° 22' N.	Long.	120° 51' W.
	Sec.	Min.	Sec.	( $^{\circ}$ Greenwich hr.)			
Beginning of series	1919	Feb.	13	0	(8)	Length of series	163 a
					( $^{\circ}$ Greenwich hr.)	Time mer.	120.00 W. = S
Middle of series	1919	May	5	12	(20)		

Compute all values to two decimal places.

Table in Harmonic Analysis and Prediction of the Tide.

		For the beginning of series.			For the middle of series.		
		(1) = $s$	(2) = $p$	(3) = $h$	(4) = $p_t$	(5) = $p$	(6) = $N$
(1) = $I$ (Table 6)	21.76						
(8) = $v$ (Table 6)	-12.68	+ (17) = 138.25	-27.41	279.60	281.55	+ 27.41	251.71
(9) = $e$ (Table 6)	-11.71	+ (8) = 12.68	-7.45	30.56	0.00	13.37	-6.35
(10) = $u'$ (Table 6)	-8.53	+ 150.93				0.45	-0.21
(11) = $2e'$ (Table 6)	-16.63					0.09	-0.04
Sum =	119.02	(G) = 32.24	(1) = 322.32	(1) = 281.55	(1) = 41.32	(G) = 245.11	
(7) = $J_1$		$M_C$		$R_2$			
(8) = $v + u$		$M_2$		$S_1$			
(9) = $e + u$		$M_3$		$S_2$			
(10) = $u + 2(M_1)$		$M_4$		$S_3$			
(11) = $2e + 2(M_2)$		$M_5$		$S_4$			
(12) = $F = (2) - (9)$	53.03	$K_1$	$N_2$	$T_2$			
(13) = $E$ (Table 13)	170.6	$N_3$	$S_1$	$(M_N)_1$			
(14) = $2(e + 90')$	123.6	$N_4$	$S_2$	$(M_N)_2$			
(15) = $2(e + 90')$	51.47	$N_5$	$S_3$	$(M_N)_3$			
(16) = $2(e + 90')$	86.78	$N_6$	$S_4$	$(M_N)_4$			
(17) = $(1) - (2)$	138.25	$L_2$	$O_1$	$T_2$			
(18) = $(1) - (2) + (17) = (13)$	282.94	$O_2$	$T_2$	$(MS)_4$			
(19) = $(2) - (16)$	321.47	$O_3$	$T_2$	$(MS)_4$			
(20) = $(2) - (16) + (17) = (13)$	282.94	$O_4$	$T_2$	$(MS)_4$			
(21) = $(1) + (2)$	151.26	$O_5$	$T_2$	$(MS)_4$			
(22) = $(20) - (21)$	131.68	$O_6$	$T_2$	$(MS)_4$			
		$M_1$	$(O_1)$	$M_2$			
(23) = $(16) - (1)$	202.45	$M_2$	$M_1$	$(2SM)_2$			
(24) = $(16) - (1) + (17) = (13)$	123.6	$M_3$	$M_1$	$(2SM)_2$			
(25) = $(9) - (5)$	0.97	$M_4$	$M_1$	$(2SM)_2$			
(26) = $(2) - (21)$	203.42	$M_5$	$M_1$	$(2SM)_2$			
(27) = $(2) - (22)$	46.84	$M_6$	$M_1$	$(2SM)_2$			
(28) = $(2) - (23)$	250.26	$M_1$	$M_1$	$(2SM)_2$			
(29) = $(2) - (24)$		$P_1$	$\mu_2$	$M_1$			
(30) = $(2) - (25)$		$P_1$	$\mu_2$	$M_1$			
(31) = $(2) - (26)$	-46.84	$P_1$	$\mu_2$	$M_1$			
(32) = $(2) - (27)$	130.73	$P_1$	$\mu_2$	$M_1$			
(33) = $(2) - (28)$	261.46	$P_1$	$\mu_2$	$M_1$			
(34) = $(2) - (29)$	351.46	$P_1$	$\mu_2$	$M_1$			
(35) = $(2) - (30)$		$P_1$	$\mu_2$	$M_1$			
(36) = $(2) - (31)$	334.15	$P_1$	$\mu_2$	$M_1$			
(37) = $(2) - (32)$	40.77	$P_1$	$\mu_2$	$M_1$			
(38) = $(2) - (33)$	-1.70	$P_1$	$\mu_2$	$M_1$			
(39) = $(2) - (34)$	203.30	$M_4$	$P_1$	$S_a$			
(40) = $(2) - (35)$	46.60	$M_4$	$P_1$	$S_a$			
(41) = $(2) - (36)$	0.24	$M_4$	$P_1$	$S_a$			
(42) = $(2) - (37)$	290.08	$M_4$	$P_1$	$S_a$			
(43) = $(2) - (38)$	220.16	$M_4$	$P_1$	$S_a$			

\* Greenwich hour = original hour + ( $5^{\circ} + 15'$ ).

\* Positive for West longitude; negative for East longitude.

FIGURE 14.

**312.** In finding the difference between the longitude of the time meridian ( $S$ ) and the longitude of the place ( $L$ ) consider west longitude as positive and east longitude as negative. In the ordinary use of form 244 it is assumed that civil time has been used in the tabulations of the observations. If, however, the original hourly heights as tabulated in form 362 are in accordance with astronomical time in which the 0 hour represents the noon of the corresponding civil day and the 12th hour the following midnight, form 244 will still be applicable if the longitude of the time meridian ( $S$ ) is taken equal to the civil time meridian plus  $180^\circ$ . For example, if tabulations have been made in astronomical time for a locality where the civil time is based upon the meridian  $15^\circ$  E., the value for  $S$  should be taken as  $-15 + 180$ , or  $165^\circ$ . If tabulations have been in Greenwich astronomical time,  $S$  should be taken as  $180^\circ$ .

**313.** *Form 244a, Log F and arguments for elimination (fig. 15).*—Items (1) to (11) are compiled here for convenience of reference for

FIGURE 15  
DEPARTMENT OF COMMERCE  
U. S. COAST AND GEODETIC SURVEY

### TIDES: Log F and Arguments for Elimination

Station ..... Morro, California		yr.	m.	d.	h.
Length of series	163 days. Series begins	1919	Feb.	13	0
Component	Log F	Component	Log F	Component	Log F
(4 dec.)		(4 dec.)		(4 dec.)	
J <sub>1</sub> . . . . .	0.0201	M <sub>2</sub> . . . . .	9.9726	MK . . . . .	0.0091
K <sub>1</sub> . . . . .	0.0160	N <sub>2</sub> , 2N . . . . .	9.9932	2MK . . . . .	0.0023
K <sub>2</sub> . . . . .	0.0472	O <sub>1</sub> . . . . .	0.0264	MN . . . . .	9.9863
L <sub>2</sub> =Log F(M <sub>2</sub> )+(7)	9.9589	OO . . . . .	0.0929	MS, 2SM . . . . .	9.9932
M <sub>1</sub> =Log F(O <sub>1</sub> )+(8)	9.8856	P <sub>1</sub> . . . . .	0.00000	Mf . . . . .	0.0596
M <sub>2</sub> . . . . .	9.9932	Q <sub>1</sub> , 2Q . . . . .	0.0264	MSf . . . . .	9.9932
M <sub>3</sub> . . . . .	9.9897	R <sub>2</sub> , S <sub>1</sub> , S <sub>2</sub> , S <sub>3</sub> , S <sub>4</sub> , T <sub>2</sub>	0.00000	Mm . . . . .	9.9772
M <sub>4</sub> . . . . .	9.9863	$\lambda_2 \mu_2 \nu_2$ . . . . .	9.9932	Sa, Ssa . . . . .	0.00000
M <sub>8</sub> . . . . .	9.9794	$p_1$ . . . . .	0.0264		

(1)= $N$ =item (6) from Form 244= 245°.11 (2 dec.)

(2)= $I$ =item (7) from Form 244= 21.76 (2 dec.)

(3)= $P$ =item (12) from Form 244= 53.08 (2 dec.)

(4)=( $h - \frac{1}{2}p$ )=item (3)- $\frac{1}{2}$  item (10), from Form 244= 327° (0 dec.)

(5)=( $h - p'$ )=item (3)- $\frac{1}{2}$  item (11), from Form 244= 331 (0 dec.)

(6)=( $h - p_i$ )=item (3)-item (4), from Form 244= 41 (0 dec.)

(7)=Log R<sub>a</sub> from Table 7= 9.9657 (4 dec.)

(8)=Log Q<sub>a</sub> from Table 9= 9.8592 (4 dec.)

(9)=Natural number from Log F(K<sub>1</sub>)= 1.038 (3 dec.)

(10)=Log f(K<sub>2</sub>)=10-Log F(K<sub>1</sub>)= 9.9528 (4 dec.)

(11)=Natural number f(K<sub>2</sub>) from (10)= 0.897 (3 dec.)

**EXPLANATION.**—For all tables see Special Publication No. 98. First fill in items (1) to (8). Then obtain values of log F for all components excepting L<sub>2</sub> and M<sub>1</sub> from Table 12. Log F(L<sub>2</sub>)=log F(M<sub>2</sub>)+log R<sub>a</sub>, and log F(M<sub>1</sub>)=log F(O<sub>1</sub>)+log Q<sub>a</sub>. Items (9) to (11) are obtained after the rest of the form has been filled out.

FIGURE 15.

this and form 452. Items (1) to (6) are obtained from values given in form 244. Item (7) is obtained from table 7, using items (2) and (3) as arguments, and item (8) is obtained from table 9, using item (3) as argument. Items (9) to (11) are obtained after the rest of the form has been filled out.

**314.** The log  $F$  for each of the listed constituents, except  $L_2$  and  $M_1$  and those for which the logarithm is given as zero, may be obtained from table 12, using item (2) as the argument. For constituents  $L_2$  and  $M_1$

$$\text{Log } F(L_2) = \text{log } F(M_2) + \text{item (7)} \quad (448)$$

$$\text{Log } F(M_1) = \text{log } F(O_1) + \text{item (8)} \quad (449)$$

If the tidal series analyzed was observed between the years 1900 and 2000, the log  $F(L_2)$  and log  $F(M_1)$  may be taken directly from

Form 194  
DEPARTMENT OF COMMERCE  
COAST AND GEODETIC SURVEY

### TIDES: HARMONIC ANALYSIS

Station	Morro, California				Lat.	35° 22' N.	Long.	120° 51' W.
Component	M	Script begins	1919	Feb.	m.	d.	h.	
			13			0		Length of series 163 Time Mcr. 120°

(DAYS)

Hourly Means from Form 142.												
(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)	(11)	(12)	(13)
(1) Hours 0 to 11...	3.08	2.53	2.12	2.02	2.25	2.84	3.53	4.13	4.53	4.56	4.25	3.76
(2) Hours 12 to 23...	3.18	2.67	2.29	2.12	2.31	2.83	3.48	4.08	4.45	4.49	4.19	3.67
(3) = (1) - (2)...	-0.10	-0.14	-0.17	-0.10	-0.06	0.01	0.05	0.05	0.05	0.08	0.07	0.09
(4) = Last 5 values of (3) reversed...	0.00	0.09	0.06	0.07	0.08	0.05	0.00	0.00	0.00	0.00	0.06	0.09
(5) = (1)+(2)...	6.26	5.20	4.41	4.14	4.56	5.67	7.01	8.21	8.98	9.05	8.44	7.43
(6) = Last half of (5)...	7.01	8.21	8.98	9.05	8.44	7.43						

(7)	(8)	(9)	(10)	(11)	(12)	(13)	(14)	(15)	(16)	(17)	(18)	(19)	(20)	(21)
Sin. $n \times 15^\circ$	(3)+(4)	(7)×(6)	Cos. $n \times 15^\circ$	(3)-(4)	(10)×(11)	Sin. $n \times 45^\circ$	(8)×(13)	Cos. $n \times 45^\circ$	(11)×(15)	Sin. $n \times 30^\circ$	(3)-(6)	(17)×(15)	Cos. $n \times 30^\circ$	(18)×(20)
12 $\omega$	12 $\omega$	12 $\omega$	12 $\omega$	12 $\omega$	12 $\omega$	12 $\omega$	12 $\omega$	12 $\omega$	12 $\omega$	12 $\omega$	12 $\omega$	12 $\omega$	12 $\omega$	
.000	-0.10	0.000	1.000	-0.10	-0.100	.000	0.000	1.000	-0.100	.000	-0.75	0.000	1.000	-0.750
.259	-0.05	-0.013	.996	-0.23	-0.222	.707	-0.035	.707	-0.163	.509	-3.01	-1.505	.856	-2.607
.500	-0.11	-0.055	.996	-0.23	-0.199	.707	-0.110	.707	-0.050	.566	-4.57	-3.958	.500	-2.285
.707	-0.03	-0.021	.707	-0.17	-0.120	.707	-0.021	.707	-0.120	.100	-4.91	-4.910	.000	0.000
.866	0.02	0.017	.500	-0.14	-0.070	.000	0.000	1.000	-0.140	.866	-3.88	-3.360	.500	+1.940
.955	0.06	0.058	.259	-0.04	-0.010	.707	-0.042	.707	-0.028	.500	-1.76	-0.880	.866	+1.524
1.000	0.05	0.050	.000	0.05	0.000	1.000	-0.050	.000	0.000	12 $\omega$	-14.613	12 $\omega$	-2.178	
		+0.036			12 $\omega$	-0.721	12 $\omega$	-0.258	12 $\omega$		0.025			

(22)	(23)	(24)	(25)	(26)	(27)	(28)	(29)	(30)	(31)	(32)	(33)	(34)	(35)	(36)	(37)
Sin. $n \times 90^\circ$	(15)×(22)	Cos. $n \times 90^\circ$	(18)×(24)	(5)+(6)	(26)-(27)	(28)×(29)	Sin. $n \times 60^\circ$	(25)×(31)	(26)+(27)	(28)×(34)	Sin. $n \times 120^\circ$	(33)×(34)	Cos. $n \times 120^\circ$	(33)×(36)	12 $\omega$
12 $\omega$	12 $\omega$	12 $\omega$	12 $\omega$	1st half.	2d half.	12 $\omega$	12 $\omega$	12 $\omega$	12 $\omega$	12 $\omega$	12 $\omega$	12 $\omega$	12 $\omega$	12 $\omega$	
0	0.000	+1	-0.750	13.27	13.19	0.09	.000	0.000	1.000	0.050	26.46	.000	0.000	1.000	26.460
+1	-3.010	0	0.000	13.41	13.00	0.41	.866	0.365	.500	0.205	26.41	.866	22.871	-3.000	-13.205
0	0.000	-1	4.670	13.39	13.10	0.29	.866	0.251	.500	-0.145	26.49	.866	-22.940	.500	-13.245
-1	4.910	0	0.000	12.98	12.46	0.00	12 $\omega$	0.606	12 $\omega$	0.140	12 $\omega$	-0.063	12 $\omega$	0.010	
0	0.000	+1	-3.880	12 $\omega$	12 $\omega$	0.000									
+1	-1.760	0	0.000	12 $\omega$	12 $\omega$	0.000									
	0.140	12 $\omega$	-0.060												

t' is in 1st quadrant when we have +s and +c.

t' is in 2d quadrant when we have +s and -c.

t' is in 3d quadrant when we have -s and -c.

t' is in 4th quadrant when we have -s and +c.

Component	M	M <sub>s</sub>	M <sub>ss</sub>				
(38)=log 12 $\omega$	8.55630	1.16474	9.41162	9.78247	9.14613	8.83885	
(39)=log 12 $\omega$	9.85794	0.33806	8.39794	9.14613	8.77813	8.00000	
(40)=(38)-(39)=log tan t'	8.69836	0.82668	1.01368	0.63654	0.36798	0.83885	
(41)=t' for beginning of the series	177.14	261.52	275.53	76.99	113.20	278.25	
(42)=local V <sub>s</sub> +u (From Form 244)	326.05	46.84	70.26	93.68	140.52	187.36	
(43)=(41)+(42)=t'	143.19	308.36	345.79	170.67	253.72	105.61	
(44)=log sin t' or	9.99523	9.99797	9.98871	9.96338	9.99548		
(45)=log cos t'	9.99946						
(46)=(35)-(41), or (39)-(45)	9.85848	1.16951	9.41365	9.79376	9.18275	8.84337	
(47)=log (augmenting factor+12)	8.92200	8.92205	8.92204	8.94058	8.96642	9.00332	
(48)=log (reciprocal of 12) for component S	8.92082	8.92082	8.92082	8.92082	8.92082	8.92082	
(49)=(46)+(47), or (46)+(45)=log R'	8.78054	0.09530	8.34569	8.73461	8.14917	7.84669	
(50)=log factor F (From Form 244a)	9.8956	9.9932	9.9897	9.9863	9.9794	9.9728	
(51)=(49)+(50)=log H'	8.66614	0.0858	8.33539	8.72091	8.12857	8.712929	
(52)=natural number from (49)=I'	0.060	1.245	0.022	0.054	0.014	0.007	
(53)=natural number from (51)=J'	0.046	1.226	0.022	0.053	0.013	0.007	

FIGURE 16.

table 13, using the year of observations, together with item (1), as argument.

**315.** *Form 194, Harmonic analysis* (fig. 16).—This form is based, primarily, upon formulas (295), (296), (303), and (304) and is designed for the computations of the first approximate values of the epochs ( $\kappa$ ) and the amplitudes ( $H$ ) of the harmonic constants. Provisions are made for obtaining the diurnal, semidiurnal, terdiurnal, quarter-diurnal, sixth-diurnal and eighth-diurnal constituents, but only such items need be computed as are necessary for the particular constituents sought. For the principal lunar series  $M_1$ ,  $M_2$ ,  $M_3$ ,  $M_4$ ,  $M_6$ , and  $M_8$ , compute all items of the form. For the principal solar series  $S_1$ ,  $S_2$ ,  $S_4$ , and  $S_8$ , items (14), (16), (33), (35), and (37) may be omitted. For the lunisolar constituents  $K_1$  and  $K_2$ , items (14), (16), and (23) to (37) may be omitted. For the diurnal constituents  $J_1$ ,  $O_1$ ,  $OO$ ,  $P_1$ ,  $Q_1$ ,  $2Q$ , and  $\rho_1$ , items (5), (6), and (14) to (37) may be omitted. For the semidiurnal constituents  $L_2$ ,  $N_2$ ,  $2N$ ,  $R_2$ ,  $T_2$ ,  $\lambda_2$ ,  $\mu_2$ ,  $v_2$ , and  $2SM$ , items (3), (4), (8) to (16), and (23) to (37) may be omitted. For terdiurnal constituents  $MK$  and  $2MK$ , items (5), (6), (9), (12), and (18) to (37) may be omitted. For quarter-diurnal constituents  $MN$  and  $MS$ , items (3), (4), (8) to (25), and (35) to (37) may be omitted. In the bottom portion of the form the symbol of the constituent is to be entered at the head of the column or columns indicated by the subscript corresponding to the number of constituent periods in a constituent day, the remaining columns being left blank.

**316.** The hourly means from form 142 (fig. 13) are entered as items (1) and (2) in regular order, beginning with the mean for 0 hour. Item (4) consists of the last five values of item (3) arranged in reverse order. Item (6) consists of the last six values of item (5) in their original order. For the computations of this form the following tables will be found convenient: table 19 of this publication for natural products, Vega's Logarithmic Tables for logarithms of linear quantities, and Bremiker's Funfstellige Logarithmen for logarithms of the trigonometrical functions. In the last table the angular arguments are given in degrees and decimals.

**317.** In choosing between items (44) and (45) the former should be used if the tabular value of (41) in the first quadrant is greater than  $45^\circ$  and the latter if this angle is less than  $45^\circ$ . In referring (41) to the proper quadrant it must be kept in mind that the signs of the natural numbers corresponding to (38) and (39) are respectively the signs of the sine and cosine of the required angles. Therefore (41) will be in the first quadrant if both  $s$  and  $c$  are positive, in the second quadrant if  $s$  is positive and  $c$  negative, in the third quadrant if both  $s$  and  $c$  are negative, and in the fourth quadrant if  $s$  is negative and  $c$  positive. In obtaining (49) use (46)+(47) for all constituents except  $S$ , and (46)+(48) for  $S$ . The log factor  $F$  for item (50) may be obtained from form 244a.

**318.** Form 194 is designed for use when 24 constituent hourly means have been obtained and all the original hourly heights have been used in the summation. If in the summation for a constituent each constituent hour of the observation period received one and only one of the hourly heights, it will be necessary to take the log-augmenting factor from table 20 and add this to the sum of items (46) and (48) to obtain item (49), striking out item (47).

**319.** This form is also adapted for use with the long-period constituents. Assuming that the daily means have been cleared of the effects of the short-period constituents (p. 89), and that these means have been assorted into 24 groups to cover the constituent period, the 24 group means may then be entered in form 194 in place of the 24 hourly means used for the short-period constituents. Then, treating the constituents  $Mm$  and  $Sa$  the same as the diurnal tides and the constituents  $Mf$ ,  $Msf$ , and  $Ssa$  as the semidiurnal tides, the form may be followed except that the log-augmenting factor must be taken from table 20 and then combined with items (46) and (48) to obtain item (49), striking out item (47).

**320.** To obtain  $Sa$  and  $Ssa$  from the monthly means of sea level, or tide level, the following process may be used: Enter the monthly means beginning with that for January in alternate spaces provided for the hourly means in form 194, placing the value for January in the space for the 0 hour. For convenience consider all the intermediate blank spaces as being filled with zero values and make the computations indicated by (3) to (12) and (18) to (21). Correct the coefficients of  $s_1$  and  $c_1$  from 12 to 6, at top and foot of columns (9), (12), (19), and (21). In bottom of form enter  $Sa$  in column having subscript 2 and  $Ssa$  in column with subscript 4 in order to obtain correct augmenting factors and strike out numerals indicating subscripts. For (38) and (39) take the logarithm of twice the values of  $6s$  and  $6c$  as obtained above. The  $\xi$ 's as obtained from (40) must have the following corrections applied in order to refer them to 0 hour of the first day of January—common years,  $Sa$  correction =  $+14.07^\circ$ ,  $Ssa$  correction =  $+28.14^\circ$ ; leap years,  $Sa$  correction =  $+14.94^\circ$ ,  $Ssa$  correction =  $+29.88^\circ$ . For convenience in recording the results it is suggested that the  $\xi$  as directly obtained from (40) be entered (in its proper quadrant) in the space just below the logarithm from which it is obtained, and that the  $\xi$  corrected to the first day of January be entered in the same line in the vacant column just to the right. The  $V+u$ , computed to the first day of January, may then be entered immediately under the corrected  $\xi$ 's and the  $\kappa'$  of (43) readily obtained. For (49) the combination (46)+(47) will be used.

**321.** *Form 452, R,  $\kappa$ , and  $\xi$  from analysis and inference (figs. 17 and 18)*—This form provides for certain computations preliminary to the regular elimination process. The constants for constituents  $K_1$  and  $S_2$  as obtained directly from form 194 may be improved by the application of corrections from tables 21 to 26; and constants for some of the smaller constituents, which have been poorly determined or not determined at all by the analysis, may be obtained by inference. If the series of observations is very short, the inferred values for the constants of some of the constituents may be better than the uneliminated values from form 194.

**322.** Form 452 is based upon paragraphs 229 to 243. It is designed to take account of the diurnal constituent on one side (fig. 17) and the semidiurnal constituents on the other side (fig. 18). The amplitudes and epochs indicated by the accent ('') are to be taken from form 194 and the quantities indicated by the asterisk (\*) from form 244 or 244a. If the series is less than 355 days, values for  $S_1$  and  $2SM$  may be omitted.

**323.** For all short series the values in columns (4) and (8) are to be computed in accordance with the equivalents and factors in columns

(3) and (7) respectively. If the series is 192 days or more in length, the  $\kappa$  of  $M_1$ ,  $P_1$ , and  $K_2$  for column (4), and the  $\log R$  of  $M_1$ ,  $P_1$ , and  $K_2$  for column (8) may be taken directly from form 194, and if the series is 355 days or more in length the  $\kappa$  and  $\log R$  of all the components for which analyses have been made may be taken directly from the same form. When a value is thus taken directly from the analysis, the corresponding equivalent in column (3) and factors in column (7) are to be crossed out.

**324.** The tabular values of items (12) and (13) for the diurnal constituents and items (14) to (18) for the semidiurnal constituents may be obtained from tables 21 to 26 or from plotted curves representing these tables, but for a series of 355 days or more in length the accelerations may be taken as zero and the resultant amplitude factors as unity.

Form 452  
DEPARTMENT OF COMMERCE  
U. S. COAST AND GEODETIC SURVEY

### TIDES: $R$ , $\kappa$ , AND $\xi$ , FROM ANALYSIS AND INFERENCE.

Station		Morro, California		Length of Series				Series begins				1919, February 13													
				163				days.				DIURNAL COMPONENTS.													
<b>FROM ANALYSIS.</b>																									
Components.		Components.		FROM ANALYSIS AND INFERENCE.				Components.				FROM ANALYSIS AND INFERENCE.													
$R'$		$t'$		Components.		$\kappa$		$V_s + u^*$		$t = (4) - (5)$		$R$		(7)											
(1)		(2)		Components.		(3)		(4)		(5)		(8)		(4 dec.)											
Ft. (3 dec.)		° (2 dec.)		Components.		Equivalent.		° (1 dec.)		° (1 dec.)		Factors.		(4 dec.)											
$J_1$		.		$J_1$		$J_1 + 0.5 \times (4)$		116.7		150.9		326		$J_1$											
$K_1$		0.967		$K_1$		$K_1' + 103.8 + (12)$		110.9		60.0		51		$\log R(J_1)$											
$M_1$		0.060		$M_1$		$M_1' - 0.5 \times (4)$		105.1		326.0		139		$\log R(F(J_1))$											
$O_1$		0.569		$O_1$		$O_1' + (12)$		99.3		342.7		117		8.6589											
$OO$		.		$OO$		$OO' + (4)$		122.5		325.6		157		$R(J_1)$											
$P_1$		0.299		$P_1$		$P_1' + (12)$		110.9		306.8		164		$R(K_1)$											
$Q_1$		0.107		$Q_1$		$Q_1' - 1.5 \times (14)$		93.5		255.9		198		$R(K_1)$											
$2Q$		.		$2Q$		$2Q' - 2.0 \times (14)$		87.7		169.1		279		$R(K_1)$											
$S_1$		.		$S_1$		$S_1' + (12)$		-		-		-		$R(M_1)$											
$P_2$		.		$P_2$		$P_2' - 1.43 \times (14)$		94.3		116.1		338		$R(M_1)$											
(9) $= P^* = -53.03$ (2 dec.); (10) $= F(K_1) = -1.038$ (3 dec.); (11) $= (K - 1)^* = 327$ (0 dec.)																									
(12) = acceleration in $K_1$ due to $P_1 = F(K_1) \times$ Table 21 = $-2.9$ (1 dec.)																									
(13) = resultant amplitude, $K_1$ and $P_1 = 1 + [F(K_1) \times (\text{Table 22})] = 1.03$ (2 dec.)																									
(14) $= (K_1' - O_1') = -11.6$ (1 dec.)																									
<i>Explanation.</i> —Obtain from Form 194 the amplitudes and epochs indicated by the accent ('), and from Form 244 or 244a the quantities indicated by the asterisk (*).																									
If the series is less than 355 days, omit $S_1$ . For all short series, the values in columns (4) and (8) will be computed in accordance with the Equivalents and Factors in columns (3) and (7), respectively; but if the series is 192 days or more in length, the $\kappa$ of $M_1$ and $P_1$ for column (4), and the $\log R$ of $M_1$ and $P_1$ for column (8) may be taken directly from Form 194; and if the series is 355 days or more in length, the $\kappa$ and $\log R$ of all the components for which analysis has been made may be taken directly from the same form. When a value is thus taken directly from the analysis, the corresponding equivalent in column (3) and factors in column (7) should be crossed out.																									
The tabular values for (12) and (13) may be obtained from Tables 21 and 22 in Special Publication No. 98 or from plotted curves representing this table. For a series of 355 days or more, (12)=0, and (13)=1.																									
Obtain the $\kappa$ of $K_1$ by applying (12) to $(K_1')^*$ from Form 194, and use this corrected $\kappa$ in computing (14). If the two angles in (14) differ by more than $180^\circ$ , add $360^\circ$ to the smaller before taking the difference, which may be either positive or negative.																									
In computing column (8) it will be noted that the corrected $\log R(K_1)$ is to be used when inferring $P_1$ .																									
<b>REMARKS:</b> _____																									
$R(S_1) = R'(S_1)$																									
$R(O_1) = R'(O_1)$																									
$R(P_1) = R'(P_1)$																									
$R(Q_1) = R'(Q_1)$																									
$R(2Q) = R'(2Q)$																									
$R(S_2) = R'(S_2)$																									
$R(O_2) = R'(O_2)$																									
$R(P_2) = R'(P_2)$																									
$R(Q_2) = R'(Q_2)$																									

FIGURE 17.

**325.** The  $\kappa$ 's of  $K_1$  and  $S_2$  are to be corrected by the accelerations as indicated before entering in column (4), and in computing item (14) for the diurnal constituents and (21) for the semidiurnal constituents the corrected  $\kappa$ 's are to be used. If the two angles in item (14) for the diurnal constituents, or in items (20) or (21) for the semidiurnal constituents, differ by more than  $180^\circ$ , the smaller angle should be increased by  $360^\circ$  before taking the difference, which may be either positive or negative. In computing column (8) it will be noted that the corrected log  $R$ 's of  $K_1$  and  $S_2$  are to be used in inferring other constituents depending upon them.

**326.** *Form 245, Elimination of component effects (fig. 19).*—This form is based upon formulas (389) and (390). One side of the form is designed for the elimination of the effects of the diurnal constituents upon each other and the other side for use with the semidiurnal constituents, the two sides being similar except for the listing of the

Form 452  
DEPARTMENT OF COMMERCE  
U. S. COAST AND GEODETIC SURVEY

### TIDES: $R$ , $\kappa$ , AND $\xi$ , FROM ANALYSIS AND INFERENCE.

Station ..... Morro, California

Length of Series ..... 163 days. Series begins ..... 1919, February 13

#### SEMIIDIURNAL COMPONENTS.

Components.	FROM ANALYSIS.		FROM ANALYSIS AND INFERENCE.				Components.	FROM ANALYSIS AND INFERENCE.			
	$R'$	$\kappa'$	(3)	(4)	(5)	(6)		(7)	(8)		
	(1)	(2)						Factors.	(4 dec.)		
Ft. (3 dec.)	% (2 dec.)	Components.	Equivalent.	% (1 dec.)	% (1 dec.)	% (0 dec.)					
$K_2$	0.090	20.63	$K_2$	304.2	299.6	5	$K_2$	log. 0.272	+ 9.4346		
$L_2$	-0.065	15.67	$L_2$	329.9	302.3	28		log. $R(S_2)$	+ 9.5017		
$M_2$	1.245	261.52	$M_2$	( $M_2$ )'	308.4	46.8	262	log. $F(K_2)$ *	- 0.0472		
$N_2$	0.263	328.84	$N_2$	( $N_2$ )'	266.9	320.1	327	log. $R(K_2)$	8.8891		
$2N$			$2N$	$N_2^*$ -(20)	265.4	233.3	32	$R(K_2)$			
$R_2$			$R_2$	$S_2^*$	304.2	219.1	85	$L_2$	log. 0.143	+ 9.1553	
$S_2$	0.314	306.43	$S_2$	( $S_2^*$ )' 304.8+(16)	304.2	356.3	306	log. $R(N_2)$	+ 9.4201		
$T_2$			$T_2$	$S_2^*$	304.2	317.5	347	log. $R(L_2)$	- 9.9657		
$\lambda_2$			$\lambda_2$	$M_2^*+0.461\times(21)$	306.5	93.5	213	$R(L_2)$	8.6097		
$P_2$	0.015	99.20	$P_2$	$M_2^*-(21)$	312.6	93.4	219	$M_2$	log. $R'(M_2)$	0.0953	
$V_2$	0.310	118.79	$V_2$	$M_2^*-0.866\times(20)$	289.8	180.2	110	$R(M_2)$	$R(M_2)=R'(M_2)$	9.4201	
2SM			2SM	(2SM)'	.....	.....	.....	2N	log. 0.133	+ 9.1239	
									log. $R(N_2)$	+ 9.4201	
									log. $R(2N)$	8.5440	
									$R(2N)$		
									$R_2$	log. 0.008	+ 7.9031
									$R(S_2)$	log. $R(S_2)$	+ 9.5017
									$R(T_2)$	7.4048	
									$S_2$	log. $R'(S_2)$	+ 9.4972
									$(V_2)$	log. $R(V_2)$	- 9.9955
									$R(S_2)$	9.5017	
									$T_2$	log. 0.059	+ 8.7709
									$R(T_2)$	$R(T_2)=R'(T_2)$	+ 9.5017
									$\lambda_2$	log. 0.007	+ 7.8451
									$R(\lambda_2)$	log. $R(\lambda_2)$	+ 0.0953
									$R(\lambda_2)$	7.9404	
									$P_2$	log. 0.024	+ 8.3802
									$R(P_2)$	log. $R(P_2)$	+ 0.0953
									$R(P_2)$	8.4755	
									$V_2$	log. 0.134	+ 9.2879
									$R(V_2)$	log. $R(V_2)$	+ 9.4201
									$R(V_2)$	8.7079	
									2SM	log. $R'(2SM)$	
										$R(2SM)=R'(2SM)$	

*Explanation.—* Obtain from Form 194 the amplitudes and epochs indicated by the accent (\*); and from Form 244 or 244a the quantities indicated by the asterisk (\*).

If the series is less than 355 days, omit 2SM. For all short series, the values in columns (4) and (8) will be computed in accordance with the Equivalents and Factors in columns (3) and (7), respectively; but if the series is 352 days or more in length, the  $\kappa$  of  $K_1$  for column (4) and the  $\log. R$  of  $K_1$  for column (8) may be taken directly from Tables 23 to 26, and if the series is 355 days or more, the  $\log. R$  of  $K_1$  for all of the components for which analysis has been made may be taken directly from the same form. When a value is thus taken directly from the analysis, the corresponding equivalent in column (3) and factors in column (7) should be crossed out.

The tabular values for (14) to (18) may be obtained from Tables 23 to 26 in Special Publication No. 98, or from plotted curves representing these tables. For a series of 355 days or more, (14)-(15)-(16)-(17)=(17)-(18)-(19); and (19)-(20)-(21)=(21)-(22)-(23).

Obtain the  $\log. R$  of  $S_2$  from Form 194, and use this corrected  $\kappa$  in computing (21). If the two angles in either (20) or (21) differ by more than  $180^\circ$ , add  $360^\circ$  to the smaller before taking the difference, which may be either positive or negative.

In computing column (8) it will be noted that the corrected  $\log. R$  of  $S_2$  is to be used in inferring other components depending upon this one.

REMARKS: \_\_\_\_\_

FIGURE 18.

constituents. The symbol  $A$  represents the constituent to be cleared, and the symbol  $B$  is the general designation for the disturbing constituents. The symbol applying to constituent  $A$  is to be crossed out in column (1) and entered in column (8). The values for items (9) and (19) are to be taken from columns (1) and (2) of form 452.

**327.** For obtaining column (2) it will be found convenient to copy the logarithms of the  $R$ 's of  $B$  from column (8) of form 452 on a horizontal strip of paper spaced the same as table 29. Applying this strip successively to the upper line of the tabular values for each con-

FORM 452  
DEPARTMENT OF COMMERCE  
COAST AND GEODETIC SURVEY

### TIDES: ELIMINATION OF COMPONENT EFFECTS

Station: Morro, California

Length of series: 163 days. Series begins: yr. 1919 m. Feb. d. 13 h. 0

(1) $B_1$	(2) $R(B_1)$ X Table 29	(3) Nat. No. (2) Table 27	(4) Table 29 -( $R(B_1)$ )	(5) (4)+(19)	(6) (3) X > 0 (5) Table 30	(7) (3) X < 0 (5) Table 30	(8) RESULTS
	log (4 dec)	ft. (3 dec)	% (ao dec)	% (ao dec)	(3 dec)	(3 dec)	
$K_2$	----	----	----	----	----	----	Component $A_2 = \text{K}_2 \dots$
$L_2$	6.0462	0.001	137	39	+0.001	+0.001	(9) = $R'(A_2)$ from analysis = 0.090
$M_2$	7.8481	0.007	290	311	-0.005	+0.005	(10) = (9) - (7) = 0.080
$N_2$	7.3351	0.002	240	261	-0.002	-0.002	(11) = log (6) = 8.6721
$2N$	6.5019	----	----	----	----	----	(12) = log (10) = 8.9051
$R_2$	7.2518	.002	195	216	+0.001	+0.002	(13) = (11) - (12) = log tan $\delta$ $\zeta$ = 9.7690
$S_2$	8.5740	.037	253	274	+0.037	+0.003	(14) = $\delta$ $\zeta$ = -30.4
$T_2$	7.5905	.004	312	333	+0.002	+0.004	(15) = log cos $\delta$ $\zeta$ = 9.9356
$\lambda_2$	6.3638	----	----	----	----	----	(16) = (12) - (15) = log $R(A_2)$ = 8.9675
$\mu_2$	5.5703	----	----	----	----	----	(16a) = log $F(A_2)$ = 0.0472
$\nu_2$	6.2207	----	----	----	----	----	(17) = (16) + log $F(A_2)$ = log $H(A_2)$ = 9.0147
$2SM$	----	----	----	----	----	----	(18) = $H(A_2)$ = 0.103
					Sums =	+0.047	(19) = $\gamma'(A_2)$ from analysis = 20.6
							(20) = (14) + (19) = $\gamma'(A_2)$ = 350.2
							(20a) = ( $V_o + u$ ) = 299.6
							(21) = (20) + ( $V_o + u$ ) = $\kappa(A_2)$ = 289.8
$K_2$	6.3256	----	----	----	----	----	Component $A_1 = \text{L}_2 \dots$
$L_2$	6.2443	0.018	293	309	-0.014	+0.011	(9) = $R'(A_2)$ from analysis = 0.065
$M_2$	7.5542	0.004	243	259	-0.004	-0.001	(10) = (9) - (7) = 0.054
$N_2$	6.6512	----	----	----	----	----	(11) = log (6) = 8.0000
$2N$	6.1512	----	----	----	----	----	(12) = log (10) = 8.7324
$R_2$	7.8741	.007	76	92	+0.007	-0.007	(13) = (11) - (12) = log tan $\delta$ $\zeta$ = 9.2676
$S_2$	7.0540	.001	135	151	-0.001	-0.001	(14) = $\delta$ $\zeta$ = -10.5
$T_2$	7.3292	.002	5	21	+0.001	+0.002	(15) = log cos $\delta$ $\zeta$ = 9.9927
$\lambda_2$	5.5935	----	----	----	----	----	(16) = (12) - (15) = log $R(A_2)$ = 8.7397
$\mu_2$	6.2592	----	----	----	----	----	(16a) = log $F(A_2)$ = 9.9589
$2SM$	----	----	----	----	----	----	(17) = (16) + log $F(A_2)$ = log $H(A_2)$ = 8.6986
					Sums =	-0.010	(18) = $H(A_2)$ = 0.050
							(19) = $\gamma'(A_2)$ from analysis = 15.7
							(20) = (14) + (19) = $\gamma'(A_2)$ = 5.2
							(20a) = ( $V_o + u$ ) = 302.3
							(21) = (20) + ( $V_o + u$ ) = $\kappa(A_2)$ = 307.5
$K_2$	6.6419	----	----	----	----	----	Component $A_2 = \text{M}_2 \dots$
$L_2$	6.7592	0.001	137	39	+0.001	+0.001	(9) = $R'(A_2)$ from analysis = 1.245
$M_2$	7.5696	0.004	223	130	+0.003	-0.003	(10) = (9) - (7) = 1.245
$N_2$	6.5791	----	----	----	----	----	(11) = log (6) = 7.6021
$2N$	5.8470	----	----	----	----	----	(12) = log (10) = 0.0955
$R_2$	7.0530	.001	81	323	-0.001	+0.001	(13) = (11) - (12) = log tan $\delta$ $\zeta$ = 7.5066
$S_2$	6.7316	.001	120	22	-0.001	-0.001	(14) = $\delta$ $\zeta$ = 0.2
$T_2$	6.3128	----	----	----	----	----	(15) = log cos $\delta$ $\zeta$ = 0.0000
$\lambda_2$	6.0268	----	----	----	----	----	(16) = (12) - (15) = log $R(A_2)$ = 0.0955
$\mu_2$	7.0803	.001	228	130	+0.001	-0.001	(16a) = log $F(A_2)$ = 9.9932
$2SM$	----	----	----	----	Sums =	+0.004	(17) = (16) + log $F(A_2)$ = log $H(A_2)$ = 0.0887
							(18) = $H(A_2)$ = 1.227
							(19) = $\gamma'(A_2)$ from analysis = 261.5
							(20) = (14) + (19) = $\gamma'(A_2)$ = 261.7
							(20a) = ( $V_o + u$ ) = 46.8
							(21) = (20) + ( $V_o + u$ ) = $\kappa(A_2)$ = 308.5

\*  $\delta$  or (14) is in the 1st quadrant when (6) is + and (10) is +.  
 \*\*  $\delta$  or (14) is in the 2d quadrant when (6) is + and (10) is -.  
 \*\*\*  $\delta$  or (14) is in the 3d quadrant when (6) is - and (10) is -.  
 \*\*\*\*  $\delta$  or (14) is in the 4th quadrant when (6) is - and (10) is +.  
 When (6) is 0 and (10) is positive, (11) & (12) = - $\infty$ , and (14) & (15) = 0.

Computed by I.A.A. Feb. 28, 1921  
(Date)  
L.P.D. " " " (Date)  
Verified by

FIGURE 19.

stituent the logarithms of the resulting products for column (2) may be readily obtained. Similarly, for column (4), the  $\xi$ 's of  $B$  from column (6) of form 452 may be copied on a strip of paper and applied to the bottom line of the tabular values for each constituent and the differences obtained. The natural numbers for column (3) corresponding to the logarithms in column (2) can usually be obtained most expeditiously from table 27, this table giving the critical logarithm for each change of 0.001 in the corresponding natural number. If the logarithm is less than 6.6990, the natural number will be too small to appear in the third decimal place, and the effects of the corresponding constituent may be considered as nil. The products for columns (6) and (7) may be conveniently obtained from table 30. In column (8) the references to (6) and (7) are to the sums of these columns. The values of  $\log F(A)$  and  $(V_0 + u)$  for column (8) may be obtained from forms 244 and 244a.

**328.** In the use of this form it will be noted that the  $R$ 's and  $\xi$ 's referring to constituent  $B$  are to be the best known values whether derived from the analysis or by inference, but the  $R'$  and  $\xi'$  of constituent  $A$ , entered as items (9) and (19), respectively, must be the unmodified values as obtained directly by form 194.

#### ANALYSIS OF TIDAL CURRENTS

**329.** Tidal currents are the periodic horizontal movements of the waters of the earth's surface. As they are caused by the same periodic forces that produce the vertical rise and fall of the tide, it is possible to represent these currents by harmonic expressions similar to those used for the tides. Constituents with the same periods as those contained in the tides are involved, but the current velocities take the place of the tidal heights. There are two general types of tidal currents, known as the reversing type and the rotary type.

**330.** In the reversing type the current flows alternately in opposite directions, the velocity increasing from zero at the time of turning to a maximum about 3 hours later and then diminishes to zero again, when it begins to flow in the opposite direction. By considering the velocities as positive in one direction and negative in the opposite direction, such a current may be expressed by a single harmonic series, such as

$$V = A \cos(at + \alpha) + B \cos(bt + \beta) + C \cos(ct + \gamma) + \text{etc.} \quad (450)$$

in which  $V$ =velocity of the current in the positive direction at any time  $t$ .

$A, B, C$ , etc.=maximum velocities of current constituents.

$a, b, c$ , etc.=speeds of constituents.

$\alpha, \beta, \gamma$ , etc.=initial phases of constituents.

**331.** In the rotary type the direction of the current changes through all points of the compass, and the velocity, although varying in strength, seldom becomes zero. In the analysis of this type of current it is necessary to resolve the observed velocities in two directions at right angles to each other. For convenience the north and east directions are selected for this purpose, velocities toward the south and west being considered as negatives of these. For the harmonic

representation of such currents it is, therefore, necessary to have two series—one for the north and the other for the east component.

332. For the analysis of either type of current the original hourly velocities or the resolved hourly velocities are tabulated in the same form used for the hourly heights of the tide. To avoid the inconvenience of negative readings in this tabulation, a constant, such as 3 knots, is added to all velocities. These hourly velocities are then summed with the same stencils that are used for the tides, and the hourly mean velocities are analyzed in the same manner as the hourly heights of the tide. The same forms are used for the currents, with the necessary modifications in the headings. The rotary currents will be represented by a double set of constants, one for the north components and the other for the east components.

333. For a 29-day series of observations, it is recommended that the analysis be made for the M series, the S series, and for  $N_2$ ,  $K_1$ , and  $O_1$ . For longer series additional constituents may be included. In the analysis of current velocities, the harmonics of the higher degrees such as  $M_4$  and  $M_6$  may be expected to be of relatively greater magnitude than they are in the tides. From theoretical considerations it may also be shown that the magnitude of the diurnal constituents as compared with the semidiurnal constituents in a simple tidal oscillation is only about one-half as great in the current as in the tide. However, because of the complexity of the tidal and current movement, the actual relation between the various constituents as determined by the analysis is subject to wide variations. The constituent  $S_1$ , which is usually negligible in the tides, may be found to be of appreciable magnitude in offshore currents because of the effect of daily periodic land and sea breezes. However, as this constituent has a speed very nearly the same as that of  $K_1$  it can be separated from the latter only by a long series of observations, preferably a year or more.

334. Form 723 (fig. 20) provides for the determination of harmonic constants from a series of current observations by comparison with corresponding constants from a tidal series covering the same period of time. This comparison is to be used if the series of observations is less than 29 days and may be used for longer series if desired. For the purpose of this comparison the hourly predicted heights at the tide station are usually to be preferred to actual observations since meteorological irregularities appearing in observed tides do not necessarily appear in a similar manner in the observed currents. In this work both currents and tides for the simultaneous period are to be summed for constituents M, S, N, K, and O; and the analysis is then carried through form 194 (Tides: Harmonic Analysis) to obtain the values of  $R'$  and  $\xi'$  for each constituent. The harmonics  $M_4$ ,  $M_6$ , and  $M_8$  are to be obtained for the current series, but may be omitted in the tidal series.

335. Enter in Form 723 the accepted  $H$  and  $\kappa$  of the principal tidal constituents for the reference station and also the values of  $R'$  and  $\xi'$  obtained from the analyses of the simultaneous series of tides and currents. The necessary calculations in the form are self-explanatory. The corrected velocity amplitude of each current constituent is obtained by a ratio on the assumption that for each constituent the relation of the corrected amplitude to the uncorrected amplitude is the same for both tide and current. The ratio derived for the con-

Form 723  
DEPARTMENT OF COMMERCE  
COAST AND GEODETIC SURVEY  
Ed. M., 1935

## CURRENTS: HARMONIC COMPARISON

(A) Current Station No. 1, Bolivar Roads, Galveston Bay, Tex. Lat.  $29^{\circ} 20.8' N$  Long.  $94^{\circ} 46.1' W = L(A) 94.77^{\circ}$   
 (B) Tide Station Galveston, Tex. (predictions) Lat.  $29^{\circ} 18.7' N$  Long.  $94^{\circ} 47.6' W = L(B) 94.79^{\circ}$

Series begins Nov. 28, 1936

Use one decimal for angles and three decimals for other quantities

Component	(1)		(2)		(3)		(4)		(5)		(6)		(7)		(8)		(9)		(10)		(11)		
	Accepted II at (B)	R' at (B)	Feet	Knots	From Simultaneous Observations	R' at (A)	Ratio (1) + (2)	Accepted H at (A)	Corrected H at (A) X (4)	Accepted κ at (B)	Corrected H at (A) X (4)	From Simultaneous Observations	f' at (B)	f' at (A)	Degrees								
M <sub>1</sub>	0.271	0.382	0.869	0.709	0.609	117.7	101.3	94.4	16.4	28.1	28.1	28.1	28.1	28.1	28.1	28.1	28.1	28.1	28.1	28.1	28.1	28.1	
M <sub>4</sub>	.....	.....	.....	.....	0.135	.....	.....	.....	.....	.....	.....	197.7	32.8	32.8	32.8	32.8	32.8	32.8	32.8	32.8	32.8	32.8	32.8
M <sub>6</sub>	.....	.....	.....	.....	0.078	.....	.....	.....	.....	.....	.....	231.2	49.2	49.2	49.2	49.2	49.2	49.2	49.2	49.2	49.2	49.2	49.2
M <sub>8</sub>	.....	.....	.....	.....	0.042	.....	.....	.....	.....	.....	.....	67.2	65.6	65.6	65.6	65.6	65.6	65.6	65.6	65.6	65.6	65.6	65.6
S <sub>1</sub>	0.084	0.380	0.862	0.221	0.191	118.7	140.2	132.4	21.5	0.0	0.0	110.8	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....
N <sub>1</sub>	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	23.8	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
K <sub>1</sub>	0.066	0.365	0.814	0.181	0.147	97.1	73.2	67.1	28.8	0.0	0.0	91.0	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....
.....	0.365	0.809	1.665	0.451	0.751	320.8	332.6	300.5	-11.8	0.0	0.0	288.7	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....
O <sub>1</sub>	0.343	0.868	1.868	0.355	0.738	311.3	284.4	256.0	26.9	0.0	0.0	282.9	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....

Remarks: Direction of flood (positive velocities)  $287^{\circ}$  (true).

Direction of ebb (negative velocities)  $111^{\circ}$  (true).

FIGURE 20.

stinent  $M_2$  is used also for the higher harmonics of  $M$ , this being considered more reliable than ratios determined directly from the much smaller amplitudes of these harmonics. The corrected epoch ( $\kappa$ ) for each current constituent is calculated on the assumption that the difference between the corrected and uncorrected epoch is the same for tide and current. For convenience the zetas ( $\zeta$ ) rather than the kappas from the simultaneous observations are used in the form and a longitude correction, column (10), is introduced to allow for this fact. Differences in column (9) for the higher harmonics of  $M_2$  are derived from the difference for that constituent because of the uncertainty in the determination of epochs of constituents of very small amplitudes.

**336.** Short series of current observations are frequently taken at half-hourly intervals. As individual observations are somewhat rough, the utilization of the half-hourly observations will add materially to the accuracy of the results obtained from an analysis. Moreover, the closer spacing of the half-hourly values will give a better development of the higher harmonics of  $M$  which are of greater relative importance in the currents than in the tides. Special stencils have been prepared for the summation of these observations. Observations taken on the exact hour are tabulated in form 362 as usual, while observations on the half-hour are offset to the right on the intermediate lines. As the series of observations under consideration are short, provisions have been made for obtaining only the diurnal constituents  $K_1$  and  $O_1$ ; the semidiurnal constituents  $M_2$ ,  $S_2$ , and  $N_2$ ; and the higher harmonics of  $M$ .

**337.** For the diurnal constituents, the special stencils provide for the same distribution, with the inclusion of the half-hourly values, as is obtained with the standard stencils used for the hourly values only. Hourly means for the constituents are obtained and entered in form 194 and all subsequent computations are the same as those based upon the use of the standard stencils.

**338.** For the semidiurnal constituents  $M_2$ ,  $S_2$ , and  $N_2$ , the semidiurnal period is divided into 24 parts. Special stencils for the constituents  $M_2$  and  $N_2$  provide for the distribution of the observed half-hourly velocities into the 24 groups indicated by this division. No stencil is required for the constituent  $S_2$ , the necessary grouping being accomplished by combining sums for afternoon observations with those for the forenoon observations of corresponding hours. Thus, the noon observations will be included with those taken at midnight, and the observations at 12:30 p. m. with those taken at 0:30 a. m.

**339.** The resulting means obtained for the semidiurnal constituents by the method described above are in reality half-hourly means, but in adapting form 194 for the analysis, these means may be entered in order in the spaces provided for the hourly means. Then, after doubling all subscripts in the form, the necessary computations may be carried out as indicated. Thus, all computations for the semidiurnal constituents will be made in the spaces originally designed for the diurnal constituents. The computations for all higher harmonics of even subscripts may be carried out in the same form using the spaces originally designed for the harmonics with subscripts one-half as great. In this adaptation of the form no provision is made for the computation of a harmonic of odd subscript which is here of rela-

tively little importance. Other forms which are used in connection with the analysis will not be affected by the use of the special stencils for the half-hourly velocities.

**340.** Observations on the half-hour may also be analyzed separately from those on the exact hour, using the standard stencils for the summation. In this case the stencils are moved to the right one column and dropped one line, thus covering the hourly values and exposing those occurring on the half-hour. Allowance must be made for the difference of a half hour in the beginning of the series when computing the  $(V_o+u)$ 's in form 244. This may be conveniently done by assuming a time meridian a half-hour or  $7\frac{1}{2}^{\circ}$  westerly from the actual time meridian used so that the first half-hourly observation will correspond to the 0 hour of the assumed time meridian. The difference of 15 minutes for the middle of the series has a negligible effect in the computations and may be disregarded. In other respects the analysis is carried on in the same manner as the analysis for the hourly observations, and the results obtained afford a useful check on the latter.

## PREDICTION OF TIDES

### HARMONIC METHOD

**341.** The methods for the prediction of the tides may be classified as harmonic and nonharmonic. By the harmonic method the elementary constituent tides, represented by harmonic constants, are combined into a composite tide. By the nonharmonic method the predictions are made by applying to the times of the moon's transits and to the mean height of the tide systems of differences to take account of average conditions and various inequalities due to changes in the phase of the moon and in the declination and parallax of the moon and sun. Without the use of a predicting machine the harmonic method would involve too much labor to be of practical service, but with such a machine the harmonic method has many advantages over the nonharmonic systems and is now used exclusively by the Coast and Geodetic Survey in making predictions for the standard ports of this country.

**342.** The height of the tide at any time may be represented harmonically by the formula

$$h = H_0 + \sum f H \cos [at + (V_0 + u) - \kappa] \quad (451)$$

in which

$h$ =height of tide at any time  $t$ .

$H_0$ =mean height of water level above datum used for prediction.

$H$ =mean amplitude of any constituent  $A$ .

$f$ =factor for reducing mean amplitude  $H$  to year of prediction.

$a$ =speed of constituent  $A$ .

$t$ =time reckoned from some initial epoch such as beginning of year of predictions.

$(V_0 + u)$ =value of equilibrium argument of constituent  $A$  when  $t=0$ .

$\kappa$ =epoch of constituent  $A$ .

In the above formula all quantities except  $h$  and  $t$  may be considered as constants for any particular year and place, and when these constants are known the value of  $h$ , or the predicted height of the tide, may be computed for any value of  $t$ , or time. By comparing successive values of  $h$  the heights of the high and low waters, together with the times of their occurrence, may be approximately determined. The harmonic method of predicting tides, therefore, consists essentially of the application of the above formula.

**343.** The exact value of  $t$  for the times of high and low waters will be roots of the first derivative of formula (451) equated to zero, which may be written—

$$\frac{dh}{dt} = -\sum af H \sin [at + (V_0 + u) - \kappa] = 0 \quad (452)$$

Although formula (452) cannot, in general, be solved by rigorous methods, it may be mechanically solved by a tide-predicting machine of the type used in the office of the Coast and Geodetic Survey.

**344.** The constant  $H_o$  of formula (451) is the depression of the adopted datum below the mean level of the water at the place of prediction. For places on the open coast the mean water level is identical with mean sea level, but in the upper portions of tidal rivers that have an appreciable slope the mean water level may be somewhat higher than the mean sea level. The datum for the predictions may be more or less arbitrarily chosen but it is customary to use the low-water plane that has been adopted as the reference for the soundings on the hydrographic charts of the locality. For all places on the Atlantic and Gulf coasts of the United States, including Puerto Rico and the Atlantic coast of the Panama Canal Zone, this datum is mean low water. For the Pacific coast of the United States, Alaska, Hawaii and the Philippines, the datum is in general mean lower low water. For the rest of the world, the datum is in general mean low water springs, although there are many localities where somewhat lower planes are used. After the datum for any particular place has been adopted its relation to the mean water level may be readily obtained from simple nonharmonic reductions of the tides as observed in the locality. The value of  $H_o$ , thus determined is a constant that is available for future predictions at the stations.

**345.** The amplitude  $H$  and the epoch  $\kappa$  for each constituent tide to be included in the predictions are the harmonic constants determined by the analysis discussed in the preceding work. Each place will have its own set of harmonic constants, and when once determined will be available for all times, except as they may be slightly modified by a more accurate determination from a better series of observations or by changes in the physical conditions at the locality such as may occur from dredging, by the depositing of sediment, or by other causes.

**346.** The node factor  $f$  (par. 77) is introduced in order to reduce the mean amplitude to the true amplitude depending upon the longitude of the moon's node. The factor  $f$  for any single constituent, therefore, passes through a cycle of values. The change being slow, it is customary to take the value as of the middle of the year for which the predictions are being made and assume this as a constant for the entire year. The error resulting from this assumption is practically negligible. Each constituent has its own set of values for  $f$ , but these values are the same for all localities and have been compiled for convenient use in table 14 for the middle of each year from 1850 to 1999.

**347.** The quantity  $a$  represents the angular speed of any constituent per unit of time. In the application of formulas (451) and (452) to the prediction of tides this is usually given in degrees per mean solar hour, the unit of  $t$  being taken as the mean solar hour. The values of the speeds of the different constituents have been calculated from astronomical data by formulas derived from the development of the tide-producing force which has already been discussed. These speeds have been compiled in table 2 and are essentially constant for all times and places. The quantity  $(V_0 + u)$  is the value of the equilibrium argument of a constituent at the initial instant from which the value of  $t$  is reckoned; that is, when  $t$  equals zero. In the prediction

of tides this initial epoch is usually taken at the midnight beginning the year for which the predictions are to be made. In strictness the  $V_0$ , or uniformly varying portion of the argument alone, refers to the initial epoch, while the  $u$ , or slow variation due to changes in the longitude of the moon's node, is taken as of the middle of the period of prediction and assumed to have this value as a constant for the entire period. The quantity  $(V_0+u)$  is different for each constituent and is also different for each initial epoch and for different longitudes on the earth. In table 15 there have been compiled the values of this quantity for the beginning of each year from 1850 to 2000 for the longitude of Greenwich. The values may be readily modified to adapt them to other initial epochs and other longitudes.

**348.** Let

$L$ =west longitude in degrees of station for which predictions are desired.

$S$ =west longitude in degrees of time meridian used at this station.

For east longitude,  $L$  and  $S$  will have negative values.

Now let

$p=0$  when referring to the long-period constituents.

1 when referring to the diurnal constituents.

2 when referring to the semidiurnal constituents, etc.

then  $p$  will be the coefficient of the quantity  $T$  in the equilibrium arguments. Now,  $T$  is the hour angle of the mean sun and is the only quantity in these arguments that is a function of the longitude of the place of observation or of prediction. At any given instant of time the difference between the values of the hour angle  $T$  at two stations will be equal to the difference in longitude of the stations. If, therefore, the value of the argument  $(V_0+u)$  for any constituent at any given instant has been computed for the meridian of Greenwich, the correction to refer this argument for the same instant to a place in longitude  $L^\circ$  west of Greenwich will be  $-pL$ , the negative sign being necessary as the value of  $T$  decreases as the west longitude increases.

**349.** The instant of time to which each of the tabular values of the Greenwich  $(V_0+u)$ 's of table 15 refers is the 0 hour of the Greenwich mean civil time at the beginning of a calendar year. In the predictions of the tides at any station it is desirable to take as the initial epoch the 0 hour of the standard or local time customarily used at that station. If, therefore, the longitude of the time meridian used is  $S^\circ$  west of Greenwich, the initial epoch of the predictions will usually be  $S/15$  mean solar hours later than the instant to which the tabular Greenwich  $(V_0+u)$ 's are referred.

**350.** In formulas (451) and (452) the symbol  $a$  is the general designation of the speed of any constituent; that is to say, it is the hourly rate of change in the argument. The difference in the argument due to a difference of  $S/15$  hours in the initial epoch is therefore  $aS/15$  degrees. The total correction to the tabular Greenwich  $(V_0+u)$  of any year in order to obtain the local  $(V_0+u)$  for a place in longitude  $L^\circ$  west at an initial epoch of 0 hours of time meridian  $S^\circ$  west at the beginning of the same calendar year is

$$\frac{aS}{15} - pL. \quad (453)$$

The general expression for the angles of (451) and (452) may now be written

$$at + (V_0 + u) - \kappa = at + \text{Greenwich } (V_0 + u) + \frac{aS}{15} - pL - \kappa \quad (454)$$

**351.** In order to avoid the necessity of applying the corrections for longitude and initial epoch to the Greenwich  $(V_0 + u)$ 's for each year, these corrections may be applied once for all to the  $\kappa$ 's. Let

$$\frac{aS}{15} - pL - \kappa = -\kappa' \quad (455)$$

Then (454) may be written

$$at + (V_0 + u) - \kappa = at + \text{Greenwich } (V_0 + u) - \kappa' \quad (456)$$

Thus, by applying the corrections indicated in (455) to the  $\kappa$ 's for any station, a modified set of epochs is obtained. These will remain the same year after year and permit the direct use of the tabular Greenwich  $(V_0 + u)$ 's in determining the actual constituent phases at the beginning of each calendar year.

**352.** Let

$$\text{Greenwich } (V_0 + u) - \kappa' = \alpha \quad (457)$$

then formulas (451) and (452) may be written

$$h = H_0 + \sum fH \cos (at + \alpha) \quad (458)$$

for height of tide at any time, and

$$\sum afH \sin (at + \alpha) = 0 \quad (459)$$

for times of high and low waters. Formula (458) may be easily solved for any single value of  $t$ , but for many values of  $t$  as are necessary in the predictions of the tides for a year at any station the labor involved by an ordinary solution would be very great. Formula (459) can not, in general, be solved by rigorous methods. The invention of tide-predicting machines has rendered the solution of both formulas a comparatively simple matter.

#### TIDE-PREDICTING MACHINE

**353.** The first tide-predicting machine was designed by Sir William Thomson (afterwards Lord Kelvin) and was made in 1873 under the auspices of the British Association for the Advancement of Science. This was an integrating machine designed to compute the height of the tide in accordance with formula (458). It provided for the summation of 10 of the principal constituents, and the resulting predicted heights were registered by a curve automatically traced by the machine. This machine is described in part I of Thomson and Tait's *Natural Philosophy*, edition of 1879. Several other tide-predicting machines designed upon the same general principles but providing for an increased number of constituents were afterwards constructed.

**354.** The first tide-predicting machine used in the United States was designed by William Ferrel, of the U. S. Coast and Geodetic Survey. This machine, which was completed in 1882, was based upon modified formulas and differed somewhat in design from any other machine that has ever been constructed. No curve was traced, but both the times and heights of the high and low waters were indicated directly by scales on the machine. The intermediate heights of the tide could be obtained only indirectly. A description of this machine is given in the report of the Coast and Geodetic Survey for the year 1883.

**355.** The first machine made to compute simultaneously the height of the tide and the times of high and low waters as represented by formulas (458) and (459), respectively, was designed and constructed in the office of the Coast and Geodetic Survey. It was completed in 1910 and is known as the United States Coast and Geodetic Survey tide-predicting machine No. 2. The machine sums simultaneously the terms of formulas (458) and (459) and registers successive heights of the tide by the movement of a pointer over a dial and also graphically by a curve automatically traced on a moving strip of paper. The times of high and low waters determined by the values of  $t$  which satisfy equation (459) are indicated both by an automatic stopping of the machine and also by check marks on the graphic record.

**356.** The general appearance of the machine is illustrated by figure 21. It is about 11 feet long, 2 feet wide, and 6 feet high, and weighs approximately 2,500 pounds. The principal features are: First, the supporting framework; second, a system of gearing by means of which shafts representing the different constituents are made to rotate with angular speeds proportional to the actual speeds of the constituents; third, a system of cranks and sliding frames for obtaining harmonic motion; fourth, summation chains connecting the individual constituent elements, by means of which the sums of the harmonic terms of formulas (458) and (459) are transmitted to the recording devices; fifth, a system of dials and pointers for indicating in a convenient manner the height of the tide for successive instants of time and also the time of the high and low waters; sixth, a tide curve or graphic representation of the tide automatically constructed by the machine. The machine is designed to take account of the 37 constituents listed in table 38, including 32 short-period and 5 long-period constituents.

**357.** The heavy cast-iron base of the machine, which includes the operator's desk, has an extreme length of 11 feet and is 2 feet wide. This forms a very substantial foundation for the superstructure, increasing its stability and thereby diminishing errors that might result from a lack of rigidity in the fixed parts. On the left side of the desk is located the hand crank for applying the power (1, fig. 24), and under the desk are the primary gears for setting in motion the various parts of the machine. The superstructure is in three sections, each consisting of parallel hard-rolled brass plates held from 6 to 7 inches apart by brass bolts. Between these plates are located the shafts and gears that govern the motion of the different parts of the machine.

**358.** The front section, or dial case, rests upon the desk facing the operator and contains the apparatus for indicating and registering the results obtained by the machine. The middle section rests upon a depression in the base and contains the mechanism for the harmonic motions for the principal constituents  $M_2$ ,  $S_2$ ,  $K_1$ ,  $O_1$ ,  $N_2$ , and  $M_4$ . The

rear section contains the mechanism for the harmonic motions for the remaining 31 constituents for which the machine provides.

**359.** The angular motions of the individual constituents, as indicated by the quantity  $at$  in formulas (458) and (459), are represented in the machine by the rotation of short horizontal shafts having their bearings in the parallel plates. All of these constituent shafts are connected by a system of gearing with the hand crank at the left of the dial case and also with the time-registering dials, so that when the machine is in operation the motion of each of these shafts will be proportional to the speed  $a$  of the corresponding constituent, and for any interval of time or increment in  $t$  as indicated by the time dials the amount of angular motion in any constituent shaft will equal the increment in the product  $at$  corresponding to that constituent.

**360.** Since the corresponding angles in formulas (458) and (459) are identical for all values of  $t$ , the motion provided by the gearing will be applicable alike to the solution of both formulas. The mechanism for the summation of the terms of formula (458) is situated on the side of the machine at the left of the operator, and for convenience this side of the machine is called the "height side" (fig. 21), and the mechanism for the summation of the terms of formula (459) is on the right-hand side of the machine, which is designated as the "time side" (fig. 22).

**361.** In table 37 are given the details of the general gearing from the hand-operating crank to the main vertical shafts, together with the details of all the gearing in the front section or dial case. It will be noted that  $S-6$  (fig. 25) is the main vertical shaft of the dial case and is connected through the releasable gears to the hour hand, the minute hand, and the day dial, respectively. The releasable gears permit the adjustment of these indicators to any time desired. After an original adjustment is made so that the hour and minute hand will each read 0 at the same instant that the day dial indicates the beginning of a day, further adjustment will, in general, be unnecessary, as the gearing itself will cause the indicators to maintain a consistent relation throughout the year, and by use of the hand-operating crank the entire system may be made to indicate any time desired. The period of the hour-hand shaft is 24 dial hours, and the hand moves over a dial graduated accordingly (3, fig. 23). The minute-hand shaft, with a period of 1 dial hour, moves over a dial graduated into 60 minutes (2, fig. 23).

**362.** The day dial, which is about 10 inches in diameter, is graduated into 366 parts to represent the 366 days in a leap year. The names of the months and numerals to indicate every fifth day of each month are inscribed on the face of the dial. This dial is located just back of the front plate or face of the machine, in which there is an arc-shaped opening through which the graduations representing nearly two months are visible at any one time (4, fig. 23). The progress of the days as the machine is operated is indicated by the rotation of this dial past an index or pointer just below the opening (6, fig. 23). This pointer is secured to a short shaft which carries at its inner end a lever arm with a pin reaching under the lower edge of the day dial, against which it is pressed by a light spring. A portion of the edge of the dial equal to the angular distance from January 1 to February 28 is of a slightly larger radius, so that the pin pressing against it rises and throws the day pointer to the right one day when this portion has passed by. On

Special Publication No. 98





FIGURE 21.—COAST AND GEODETIC SURVEY TIDE-PREDICTING MACHINE.

FIGURE 22.—TIDE-PREDICTING MACHINE, TIME SIDE.



Special Publication No. 98




FIGURE 23.—TIDE-PREDICTING MACHINE, RECORDING DEVICES.

Special Publication No. 98




FIGURE 24.—TIDE-PREDICTING MACHINE, DRIVING GEARS.

Special Publication No. 98




FIGURE 25.—TIDE-PREDICTING MACHINE. DIAL CASE FROM HEIGHT SIDE.

Special Publication No. 98




FIGURE 26.—TIDE-PREDICTING MACHINE. DIAL CASE FROM TIME SIDE.

Special Publication No. 96




FIGURE 27.—TIDE-PREDICTING MACHINE. VERTICAL DRIVING SHAFT OF MIDDLE SECTION.

Special Publication No. 98




FIGURE 28.—TIDE-PREDICTING MACHINE. FORWARD DRIVING SHAFT OF REAR SECTION.

Special Publication No. 98




FIGURE 29.—TIDE-PREDICTING MACHINE, REAR END.

Special Publication No. 98




FIGURE 30.—TIDE-PREDICTING MACHINE, DETAILS OF RELEASABLE GEAR.




FIGURE 31.—TIDE-PREDICTING MACHINE, DETAILS OF CONSTITUENT CRANK.

the last day of December this pointer will move back one day to its original position.

**363.** On the same center with the day pointer there is a smaller index (7, fig. 23) which may be turned either to the right toward a plate inscribed "Common year," or to the left to a plate inscribed "Leap year." When this smaller index is turned toward the right, the day pointer is free to move in accordance with the change in radius of the edge of the dial. If the smaller index is turned toward the left, the day pointer is locked and must hold a fixed position throughout the year. For the prediction of the tides for two or more common years in succession the day dial must be set forward one day at the close of the year in order that the days of the succeeding year may be correctly registered. The day dial can be released for setting by the nut (5, fig. 23) immediately above the large dial ring. A slower movement of the day dial is provided by a releasable gear on the vertical shaft  $S-6$  (fig. 25).

**364.** There are three main vertical shafts  $S-13$  (fig. 27),  $S-14$  (fig. 28), and  $S-16$  (fig. 29), to which are connected the gearing for the individual constituents. The period of rotation of each is 12 dial hours, and all move clockwise when viewed from above the machine. The connections between these main shafts and the individual constituent crankshafts are, in general, made by two pairs of bevel gears and an intermediate horizontal shaft, except that for the slow moving constituents  $Sa$ ,  $Ssa$ ,  $Mm$ ,  $Mf$ , and  $MSf$ , a worm screw and wheel and a pair of spur gears are in each case substituted for a pair of bevel gears. In each case the gear on the main vertical shaft is releasable so that each crankshaft can be set independently.

**365.** Main shaft  $S-13$  in the middle section of the machine drives 9 individual crankshafts representing 6 constituents, 3 of them being provided with two crankshafts each. These 6 constituents are  $M_2$ ,  $S_2$ ,  $K_1$ ,  $N_2$ ,  $M_4$ , and  $O_1$ , the first three having the double crankshafts. Main shaft  $S-14$  at the front of the rear section of the machine drives 16 crankshafts representing one constituent each. These are  $M_6$ ,  $MK$ ,  $S_4$ ,  $MN$ ,  $\nu_2$ ,  $S_6$ ,  $\mu_2$ , and  $2N$  in the upper range, and  $MS$ ,  $M_8$ ,  $K_2$ ,  $2MK$ ,  $L_2$ ,  $M_3$ ,  $2SM$ , and  $P_1$  in the lower range. Main shaft  $S-16$  at the back of the rear section drives 15 crankshafts. The constituents represented are  $OO$ ,  $\lambda_2$ ,  $S_1$ ,  $M_1$ ,  $J_1$ ,  $Mm$ , and  $Ssa$ , in the upper range, and  $2Q$ ,  $R_2$ ,  $T_2$ ,  $Q_1$ ,  $\rho_1$ ,  $Mf$ ,  $MSf$ , and  $Sa$  in the lower range.

**366.** For each of the five long-period constituents motion is communicated from the intermediate shaft by a worm screw and wheel to a small shaft on which is mounted a sliding spur gear. The latter engages a spur gear on the crankshaft, but may be easily disconnected by drawing out a pin on the time side of the machine, thus permitting the crankshaft to be turned freely when setting the machine.

**367. Gear speeds.**—The relative angular motion of each constituent crankshaft must correspond as nearly as possible to the theoretical speed of the constituent represented. The period of rotation of each of the three main vertical shafts being 12 dial hours, the angular motion of each of these shafts is  $30^\circ$  per dial hour. Table 38 contains the details of the gearing from the main vertical shafts to the individual crankshafts, the number of teeth in the different gears for each constituent being given in columns I, II, III, and IV. In designing the predicting machine it was necessary to find such values for these

columns as would give gear speeds approximating as closely as possible with the theoretical speeds of the constituents. By comparing the gear speeds as obtained with the corresponding theoretical speeds it will be noted that the accumulated errors of the gears for an entire dial year for all the constituents are negligible in the prediction of the tides.

**368. Releasable gears.**—Releasable gears (52, fig. 27) on the main vertical shafts permit the independent adjustment of the time indicators and individual crankshafts. The details of these gears are illustrated in figure 30. A collar *C*, with a thread at its upper end and a flange at the bottom, is fastened to the shaft by means of three steel screws. The gear wheel *A* fits closely upon this collar and rests upon the flange. It has sunk into its upper surface a recess *a*, which is filled by the flange of collar *B*. When in place, the latter is prevented from turning by a small steel screw reaching into a vertical groove *c* in the collar *C*. The lower surface of collar *B* is slightly dished, and the collar is split twice at right angles nearly to the top. When the milled nut *D* is screwed down with a small pin wrench, the edge of the collar *B* is pressed against the edge of the recess *a* with such force as to make slipping practically impossible. When the nut is loosened, the gear may be turned independently of the main driving shaft. A small wrench (56, fig. 28) is used for setting these gears. Each of the three main driving shafts is provided with a clamp (55, fig. 28) to secure the shaft from turning when the nut of the releasable gear is being loosened or tightened.

**369. Constituent cranks.**—Secured to the ends of the constituent crank shafts, which projects through the brass plates on both sides of the machine, are brass cranks (40, fig. 25) which are provided for the constituent amplitudes. Those on the left or height side of the machine are designated as the constituent height cranks and are used for the coefficients of the cosine terms of formula (458), and those on the right or time side of the machine are designated as the constituent time cranks and are used for the coefficients of the sine terms of formula (459). The time crank on each constituent crank shaft is attached  $90^{\circ}$  in advance (in the direction of rotation) of the height crank on the same shaft. For the constituents *Sa* and *Ssa* no time cranks are provided, as the coefficients of the sine terms corresponding to these constituents are too small to be taken into account. The direction of rotation of each constituent crank shaft with its constituent cranks is clockwise when viewed from the time side of the machine and counterclockwise when viewed from the height side. The details of a constituent crank are shown in figure 31. The pointer *a* is rigidly attached to the crank as an index for reading its position on a dial. In each crank there is a longitudinal groove *b* with flanges in which a crank pin *d* may be clamped in any desired position. The crank pin has a small rectangular block as a base which is designed to fit the groove in the crank, and through the center of the crank pin there is a threaded hole for the clamp screw *f*. Attached to the under side of the crank-pin block is a small spring *c* that presses the block outward against the flanges of the groove, keeping it from slipping out of place when unclamped and at the same time permitting it to be moved along the groove when setting the machine. The crank pin may be securely fastened in any desired position by tightening up on the clamp screw, which, pressing

against the small spring at the back, forces the crank-pin block outward against the flanges of the groove with sufficient pressure to prevent any slipping. A milled head wrench *B* is used for tightening the clamp screw. A small rectangular block *e* of hardened steel is fitted to turn freely upon the finely polished axle of the crank pin. This block is designed to fit into and slide along the slot of the constituent frame.

**370. Positive and negative direction.**—All the constituent crank shafts and cranks may be grouped into two ranges—those above the medial horizontal plane of the framework being in the upper range and those below this plane in the lower range. In the following discussion direction toward this medial plane is to be considered as negative and direction away from the plane as positive; that is to say, for all constituents in the upper range the positive direction will be upward and the negative direction downward, while for the constituents in the lower range the positive direction will be downward and the negative direction upward.

**371. Constituent dials.**—To indicate the angular positions of the constituent crank shafts, the pointer (*a*, fig. 31) moves around a dial (41, fig. 25) which is graduated in degrees. These dials are fastened to the frame of the machine back of the constituent cranks on both sides of the machine, those on the time side being graduated clockwise and those on the height side counterclockwise. These dials and pointers are so arranged that the angular position of a constituent crank shaft at any time will be the same whether read from the dial on the height side or from the dial on the time side of the machine, and at the zero reading for any constituent the height crank will be in a positive vertical position and the corresponding time crank in a horizontal position. At a reading of  $90^\circ$  the height crank will be horizontal and the time crank in a negative vertical position.

**372.** With the face of the machine registering the initial epoch, such as January 1, 0 hour, of any year, the value of  $t$  then being taken as zero, each constituent crank shaft may be set, by means of its releasable gear, so that the dial readings will be equal to the  $\alpha$  of the corresponding constituent as represented in formulas (458) and (459). If the machine is then put in operation, the dial readings will, for successive values of  $t$ , continuously correspond to the angle  $(at + \alpha)$  of the formulas, as the gearing already described will provide for the increment  $at$ .

**373. Constituent sliding frames.**—For each constituent crank there is a light steel frame (42, fig. 25) fitted to slide vertically in grooves in a pair of angle pieces attached to the side plates of the machine. At the top of the frame there is a horizontal slot in which the crank pin slides. As the machine is operated the rotation of the crank shafts with their cranks cause each crank pin to move in the circumference of a circle, the radius of which depends upon the setting of the pin on the crank. This motion of the pin, acting in the horizontal slot of the sliding frame, imparts a vertical harmonic motion to that frame. The frame is in its zero position when the center horizontal line of the slot intersects the axis of the crank shaft. Positive motion is the direction away from the medial horizontal plane of the machine and negative motion is toward the medial plane. The displacement of each constituent height frame from its zero position will always equal the product of the amplitude setting of the crank pin by the cosine

of the constituent dial reading, and the displacement of each constituent time frame will always equal the product of the amplitude setting by minus the sine of the constituent dial reading.

**374. Constituent pulleys.**—Each constituent frame is connected with a small movable pulley (43, fig. 25). For all constituents except  $M_2$ ,  $S_2$ ,  $N_2$ ,  $K_1$ ,  $O_1$ , and  $Sa$  on the height side and  $M_2$ ,  $S_2$ ,  $N_2$ , and  $M_4$  on the time side this connection is by a single steel strip, so that the pulley has the same vertical motion as the corresponding frame.

**375. Doubling gears.**—Because of the very large amplitudes of some of the constituents two methods were used in order to keep the lengths of the cranks within practical limits. For  $M_2$ ,  $S_2$ , and  $K_1$  two sets of shafts and cranks were provided, so that the amplitudes of these constituents may be divided when necessary and a portion set on each. A further reduction in the length of the cranks for these and the other large constituents is accomplished by the use of doubling gears between the sliding frame and movable pulley. Two spur gears with the ratio of 1:2 (48, fig. 25) are arranged to turn together on the same axis. The smaller gear engages a rack (46) attached to the sliding frame and the larger gear engages a rack (47) attached to the constituent pulley. Each rack is held against its gear by a flange roller (49), and counterpoise weights are provided to take up the backlash in the gears. Through the action of these doubling gears any motion in the sliding frame causes a motion twice as great in the constituent pulley. Doubling gears are provided on the height side of the machine for constituents  $M_2$ ,  $S_2$ ,  $N_2$ ,  $K_1$ ,  $O_1$ , and  $Sa$  and on the time side for constituents  $M_2$ ,  $S_2$ ,  $N_2$ , and  $M_4$ .

**376. Scales for amplitude settings.**—The scales for setting the constituent amplitudes are attached to the frame of the machine and are, in general, graduated into units and tenths (44, fig. 25). The scales are arranged to read in a negative direction; that is, downward for the constituents of the upper range and upward for the constituents in the lower range. On a small adjustable plate (45) attached to each constituent pulley there is an index line which is set to read zero on the scale when the sliding frame is in its zero position. For setting the crank pins for the constituent amplitudes the cranks to be set are first turned to a negative vertical position. For the cranks on the height side of the machine this position corresponds to a dial reading of  $180^\circ$  and for the cranks on the time side to a reading of  $90^\circ$ .

**377.** The scales on the height side of the machine, which are used in setting the coefficients of formula (458), are graduated uniformly one-half inch to the unit. On the time side of the machine the scales are modified in order to automatically take account of the additional factor involving the speed of the constituent which appears in each of the coefficients of formula (459). Dividing the members of this formula by  $m$ , the speed of constituent  $M_2$ , it becomes

$$\sum \frac{a}{m} fH \sin (at + \alpha) = 0 \quad (460)$$

The modified scales are graduated  $0.5 a/m$  inch to the unit. The use of the modified scales on the time side of the machine permits both the height and time crank for any constituent to be set in accord with the factor  $fH$  which is common to the coefficients of both formulas (458) and (459). There are also provided for special use on the time

side of the machine unmodified scales graduated uniformly to read in a positive direction.

**378. Summation chains.**—The summations of the several cosine terms in formula (458) and of the several sine terms in formula (459) are carried on simultaneously by two chains, one (27, fig. 25) on the height side and the other (28, fig. 26) on the time side of the machine. The chains are of the chronometer fuse type, of tempered steel, and have 125 links per foot. The total length of the height chain is 27.6 feet and of the time chain 30.6 feet. A platinum point is attached to one of the links of the time chain 3.5 feet from its free end for an index.

**379.** Each of these chains is fastened at one end near the back part of the machine by a pair of adjusting screws (53, fig. 29, and 54, fig. 22). From these adjusting screws each chain passes alternately downward under a constituent pulley of the lower range and upward over a constituent pulley of the upper range, spanning the space between the rear and middle section of the machine by two idler pulleys and continuing until every constituent pulley on each side of the machine is included in the system. The movable pulleys are so arranged that the direction of the chain in passing from one to another is always vertical and parallel to the direction of the motion of the sliding frames.

**380. Summation wheels.**—The free or movable end of each of the chains is attached to a threaded grooved wheel (29, 30, fig. 25), 12 inches in circumference and threaded to hold more than seven turns of the chain, or about 90 inches in all. These are called the height and time summation wheels. Each is mounted on a shaft that admits a small lateral motion, and by means of a fixed tooth attached to the framework of the machine and reaching into the threads of a screw fastened to the shaft the latter when rotating is forced into a screw motion with a pitch equal to that of the thread groove of the summation wheel; so that the path of the chain as it is wound or unwound from the summation wheel remains unchanged.

**381.** The height summation wheel (29, fig. 25) is located near the front edge of the middle section of the machine, where it receives the height summation chain directly from the nearest constituent pulley. The time summation pulley (30) is located inside the dial case near the lower left side, and three fixed pulleys are used to carry the time chain from the end constituent pulley to the summation wheel. Counterpoise weights are connected with the shafts containing the summation wheels in order to keep the summation chains taut.

**382.** When all of the sliding frames on either side of the machine are in their zero positions, the corresponding summation wheel is approximately half filled by turns of the summation chain. Any motion of a sliding frame in a positive direction will tend to unwind the chain from the wheel, and any motion in the negative direction will tend to slacken the chain so that it will be wound up by the counterpoise weight. With several of the sliding frames on either side of the machine moving simultaneously, the resultant motion, which is the algebraic sum of all, will be communicated to the summation wheel. The motion of the sliding frame being transmitted to the chain through a movable pulley, the motion of the free end of the chain must be twice as great as that in the pulley. The scale of the pulley motion is one-half inch to the unit of amplitude, and there-

fore the scale of the chain motion is 1 inch to the unit, and one complete rotation of the summation wheel represents a change of 12 units of amplitude.

**383.** The zero position of the height summation wheel is indicated by the conjunction of an index line (50, fig. 25) on the arm attached to the wheel and an index line (51, fig. 25) on a bracket attached to the framework of the machine just below the summation wheel, the wheel itself being approximately one-half filled with the summation chain. The length of the chain is adjusted so that the summation wheel will be in its zero position when all the sliding frames on the height side of machine are in their zero positions. It will be noted that the conjunction of the index lines will not alone determine the zero position of the wheel, since such conjunctions will occur at each turn of the wheel, while there is only one zero position, which is that taken when the constituent frames are set at zero.

**384.** The zero position of the time summation wheel is indicated by the conjunction of an index point (11, fig. 23) attached to the time summation chain and a fixed index (12, fig. 23) in the middle of the horizontal opening near the bottom of the dial case, and the length of the time summation chain is so adjusted that this conjunction will occur when all sliding frames on the time side of machine are in their zero positions.

**385. Predicted heights of the tide.**—When the machine is in operation, the sum of all the cosine terms of formula (458) included in the settings for a station will be transmitted through the height summation wheel to the face of the machine and there indicated in two ways—first by a pointer moving over a circular height scale (8, fig. 23) and second by the ordinates of a tide curve that is automatically traced on a roll of paper (15, fig. 23). The motion of the height summation wheel is transmitted by a gear ratio of 30 : 100 to a horizontal shaft which is located just back of the dial case. One complete rotation of this shaft represents 40 units in the height of the tide. From this shaft the motion is carried by two separate systems of gearing to the height pointer on the face of the machine and to the pen that traces the tide curve.

**386. Height scale.**—The height pointer is geared to make one complete revolution for a change of 40 units in the height of the tide. A height scale, with its circumference divided into 40 equal parts and each of these unit parts subdivided into tenths, provides for the direct registering of the sum of the cosine terms of formula (458) as communicated through the summation wheel. This scale has its zero graduation at the top and is graduated positively to the right and negatively to the left. The height pointer can easily be adjusted to any position by means of a small milled nut (10, fig. 23) at the end of its shaft. If it should be desired to refer the predicted heights to mean sea level, this pointer must be adjusted to read zero at the same time that the summation wheel is in its zero position; but if it is desired to refer to some other datum, the pointer will be adjusted according to the elevation of mean sea level above this datum. For the value of  $h$  in formula (458) the pointer will be adjusted to a reading corresponding to the adopted value of  $H_0$  at the time the summation wheel is in its zero position, then this value of  $H_0$  will be automatically included with the sum of the cosine terms of that formula. As the

machine is operated the height pointer will indicate the predicted height of the tide corresponding to the time shown on the time dials.

**387.** In order to increase the working scale of the machine when predicting tides with smaller ranges, two additional circular height scales are provided, one with the circle divided into 20 units and the other into 10 units, with the units subdivided into tenths. These scales may be easily removed or replaced on the machine, the scale in use being secured in place by a small button at the top (9, fig. 23). The 20-unit scale may be conveniently used when the extreme range of the predicted tide at any place is between 10 and 20 feet, and the 10-unit scale when the extreme range is less than 10 feet. If the 20-unit scale is to be used, the value of each coefficient of both the cosine and the sine terms must be doubled before setting the component cranks, and if the 10-unit scale is used these original coefficients must first be multiplied by 4 before setting the values in the machine. If the extreme tide is less than 4 feet, the 40-unit dial may be readily used as a 4-unit scale by considering the original unit graduations as tenths of units in the larger scale. In this case the coefficients of the cosine and sine terms of the formula must be multiplied by 10 before entering in the machine. The factor used for multiplying the coefficients to adapt them to the different height scales is called the working scale of the machine. Working scales of 1, 2, 4, and 10 are now in general use to take account of the different ranges of tide at the places for which predictions are made.

**388. Predicted times of the tide.**—Simultaneously with the summation of the cosine terms of formula (458) on the height side of the machine, the summation of the sine terms of formula (460), which was derived from formula (459), is being effected on the time side. Being concerned only with the time at which the sum of the sine terms is zero, no provision is made for registering the sum except at this time, which is indicated on the machine by the conjunction of the index point on the time chain and the fixed platinum index in the dial case. Near the time of a high water the index on the chain moves from right to left and near the time of a low water from left to right. The conjunction of the movable and fixed index is visible to the operator of the machine and he may note the corresponding dial readings for the time and height of the high or low water.

**389. Automatic stopping device.**—This device provides for automatically stopping the machine at each high and low water. Secured to the hand-crank shaft is a ratchet wheel and just above the ratchet wheel is a steel pawl (25, fig. 24) operated by an electromagnet (26) mounted under the desk top. The electric circuit for the electromagnet is closed by a contact spring that rests upon a hard-rubber cylinder (31, fig. 25) on the rear end of the shaft on which the time summation wheel is mounted. A small platinum plug in this rubber cylinder comes in contact with the spring, which is fitted with a fine motion adjustment, when the time summation chain registers zero. This closes the circuit and draws the pawl against the ratchet wheel, thereby automatically stopping the machine. The lateral screw motion of the shaft on which the rubber cylinder is mounted prevents the platinum plug from coming in contact with the spring on any revolution other than the one which brings the time chain to its zero position. The circuit is led through an insulated ring on the hub of the hand crank where a contact is kept closed by a spring. After

the operator has noted the time and height readings of the high or low water he may easily break the circuit at the crank hub by a slight inward pressure against the crank handle, thus releasing the armature and pawl and permitting the machine to be turned forward to the next stop. By means of a small switch (23, fig. 24) just below the crank the circuit may be held open to prevent the automatic device from operating when so desired.

390. *Nonreversing ratchet*.—Upon the crank shaft, close to the bearing in the desk frame, there is a small ratchet wheel and above this there is a pawl (24, fig. 24) that is lifted away from the wheel by friction springs when the machine is being turned forward but which is instantly thrown into engagement when the crank is accidentally turned backward. By pushing in one of the small buttons (22, fig. 24) just above the crank the pawl is locked so that it cannot engage the ratchet, thus permitting the machine to be turned backward when desired. Pressure on another button releases the pawl.

391. *Tide curve*.—The tide curve which graphically represents the rise and fall of the predicted tide is automatically traced on a roll of paper by the machine at the same time that the results are being indicated on the dials. The curve is the resultant of a horizontal movement of the paper, corresponding to the passing of time, and a vertical movement of a fountain pen (13, fig. 23), corresponding to the rise and fall of the tide. The paper is 6 inches wide with about 380 feet to the roll, which is sufficient to include a little more than a full year of record of the predicted tides at a station. The paper should be about 0.0024 inch thick in order that the complete roll may be of a suitable size for use in the machine.

392. Within the dial case, near the upper right-hand corner, is a mandrel (33, fig. 25), which can be quickly removed and replaced. It is designed to hold the blank roll of paper, the latter being wound upon a wooden core especially designed to fit on the mandrel. At the bottom of the mandrel is an adjustable friction device to provide tension on the paper. From the blank roll the paper is led over an idler roller (34, fig. 25), mounted in the front plate of the dial case, then across the face of the machine for a distance of about 13 inches to a feed roller (35, fig. 25), then over the feed roller to the receiving roller (36, fig. 25), upon which it is wound.

393. The feed roller governs the motion of the paper across the face of the machine and is provided near each end with 12 fine needle points to prevent the paper from slipping. The feed roller is controlled by the main vertical shaft of the dial case through gearing of such ratio that the feed roller will turn at the same rate as the main vertical shaft; that is to say, one complete turn of the feed roller will represent 12 dial hours in time. The feed roller being 6 inches in circumference the paper will be moved forward at the rate of one-half inch to the dial hour. A ratchet and pawl (37, fig. 25) are so placed as to leave the paper at rest when the machine is turned backward. If desired, the paper feed can be thrown out of action altogether by turning a small milled head on the ratchet gear.

394. To provide for the winding up of the paper on the receiving roller there is a sprocket wheel (38, fig. 25) held by adjustable friction to the upper end of the feed roller. Fitted to the top of the receiving roller is a smaller sprocket which is driven by a chain from the feed-roller sprocket. The ratio of the sprockets is such as to force the

receiving roller to wind up all the paper delivered by the feed roller, the tension on the paper being kept uniform by the friction device. To remove a completed roll of record the smaller sprocket is lifted from the receiving roller and a pin (39, fig. 25) at the back of the dial case is drawn out, releasing the upper bearing bracket. The bracket can then be raised and the receiving roller with its record removed. A similar bracket secured by a pin is provided for the removal of the mandrel on which the blank roll of paper is placed.

**395. Marigram gears.**—The pen that traces the tide curve is mounted in a carriage which is arranged to slide vertically on a pair of guiding rods and is controlled from a horizontal shaft at the back of the dial case. On this shaft there is mounted a set of three sliding change gears (18, fig. 26), which are designed to mesh, respectively, with three fixed gears mounted on a shaft just above. By sliding the change gears in different positions any one of them may be brought into mesh with its corresponding fixed gear. These gears provide for ratios of 1:1, 2:1, and 3:2, according to whether the innermost, the middle, or the outer gears are in mesh. At the outer end of the shaft containing the fixed gears is a thread-grooved wheel 4 inches in circumference (19, fig. 26), to which is attached one end of the pen-carriage chain (20, fig. 26). The chain is partly wound upon the wheel and from it passes through the dial case to the front of the machine, then upward over a pulley near the top to a counterpoise weight within the dial case. The pen carriage is secured to this chain by means of a clamp and can be adjusted to any desired position.

**396. Scale of tide curve.**—With a working scale of unity, the rotation of the height summation wheel, as transmitted through marigram gear ratio of 1:1 to the curve-line pen, will move the latter vertically 0.1 inch for each unit change in the sum of the harmonic terms and this may be taken as the basic or natural scale of the graphic record. This scale may be enlarged by the factor 3/2 or 2 through the use of one of the other gear ratios and may be further modified to any desired extent by the introduction of an arbitrary working scale-factor. Letting  $G$  equal the marigram gear ratio (1, 3/2, or 2) and  $S$  equal the working scale factor applied to the amplitude settings, the vertical scale of the graphic record may be expressed as follows:

$$1 \text{ inch of graph represents } 10/GS \text{ units of summation} \quad (461)$$

$$1 \text{ summation unit is represented by } GS/10 \text{ inches in graph} \quad (462)$$

The scale ratio of the graph will differ with different units used in the predictions. Thus

$$\text{Graph scale (amplitude settings in feet)} = GS/120 \quad (463)$$

$$\text{Graph scale (amplitude settings in meters)} = GS/393.7 \quad (464)$$

$$\text{Graph scale (amplitude settings in decimeters)} = GS/39.37 \quad (465)$$

**397.** In selecting the marigram gear ratio and scale factor for the predictions at any station, it is the general aim to secure as large a scale as possible while keeping the graph within the limits of the paper. Some consideration must be given also to the limits of the height dial scale and in some instances to the mechanical limits of the individual amplitude settings. The marigram gear ratio affects the graph only but the scale factor affects also the amplitude settings and the height dial readings. The extreme amplitude of the graphic

record is limited by the width of the paper which extends 3 inches on either side of the medial line, but for mechanical reasons it is desirable in general to keep the record within a band  $2\frac{1}{2}$  inches on either side of the medial line. The following table suggests suitable scale, dial, and gear combinations for different tidal ranges and different current velocities. The tabular marigram scales are applicable only when the foot or knot has been used as the unit for machine settings. The marigram amplitude limits given in the last column are expressed in the same unit that is used in setting the machine regardless of what unit that may have been.

*Working scale, height dial, marigram gear, and scale*

Tidal range limits	Current velocity limits	Working scale factor	Height dial	Mari- gram gear	Marigram scale		Mari- gram ampli- tude limit
					Settings in feet	Settings in knots	
<i>Feet</i>	<i>Knots</i>				<i>Ratio</i>	<i>Knots per inch</i>	<i>Units</i>
0.0- 2.5	0.0- 1.0	10	4	2:1	1: 6	0.50	1.5
2.6- 3.5	1.1- 1.5	10	4	3:2	1: 8	0.67	2.0
3.6- 4.0	1.6- 2.0	10	4	1:1	1:12	1.00	3.0
4.1- 6.0	2.1- 3.0	4	10	2:1	1:15	1.25	3.7
6.1- 8.0	3.1- 4.0	4	10	3:2	1:20	1.67	5.0
8.1-10.0	4.1- 5.0	4	10	1:1	1:30	2.50	7.5
10.1-12.5	5.1- 6.0	2	20	2:1	1:30	2.50	7.5
12.6-16.5	6.1- 8.0	2	20	3:2	1:40	3.33	10.0
16.6-20.0	8.1-10.0	2	20	1:1	1:60	5.00	15.0
20.1-25.0	10.1-12.5	1	40	2:1	1:60	5.00	15.0
25.1-32.5	12.6-16.0	1	40	3:2	1:80	6.67	20.0
32.6-	16.1-	1	40	1:1	1:120	10.00	30.0

When height dial readings are not required, and amplitude settings are in feet, a convenient graph scale of 1:10 can be obtained by using any one of the following combinations; scale factor 12 with gear ratio 1:1, scale factor 8 with gear ratio 3:2, or scale factor 6 with gear ratio 2:1.

**398.** When the tide-predicting machine is used for the prediction of the tide-producing force, the graph scale to be adopted will depend upon the unit in which the force is to be expressed. Assume that the sum of all terms in the vertical component of the force (par. 79) is desired. Referring to paragraph 43, it will be noted that the extreme value of this component due to the combined action of moon and sun is approximately  $0.2 \times 10^{-6}$  with the unit of force taken as  $g$ , the mean acceleration of gravity. In this case a convenient scale relation which will bring the graph within the desired limits on the paper is obtained by adopting a working scale factor of  $6 \times 10^7$  with the marigram gear ratio of 2:1. With this combination 0.1 foot of graph ordinate will represent  $10^{-7} g$  units of force. In practice the scale factor would be combined with the general coefficient common to all terms in the formulas.

**399. Pens.**—The curve-line pen (13, fig. 23) and the datum-line pen (14) are each of the ordinary fountain type. Each is fitted with a metal lock joint, so that it may be quickly removed and replaced in the same position, and is pressed against the paper by a light coil spring when in use. The curve-line pen is mounted in a swivel arm on a light carriage which slides vertically along two rods. The datum-line pen is mounted in a swivel arm that may be adjusted so that the mean sea-level line will be traced midway between the upper and lower edges of the paper.

**400. Hour-marking device.**—The arm for the datum-line pen is secured to the outer end of a shaft which carries two armatures, one for the upper and the other for the lower of two electromagnets (17, fig. 26). A spring keeps the armatures at equal distances from their respective electromagnets. The upper electromagnet is designed for indicating the hours on the datum line and is in a circuit that is opened and closed by a platinum-tipped contact spring resting upon the edge of an ivory disk in which are embedded, equally spaced, 24 narrow strips of platinum (32, fig. 25). The ivory disk is mounted on the shaft of the hour pointer, and as this rotates the platinum strips successively make an electric contact that throws the datum-line pen downward for an instant, making a corresponding jog in the datum line, the downward stroke of the pen indicating the exact hour. An extra strip of platinum placed close to the one representing the midnight hour causes a double jog for the beginning of each day, the downward stroke of the second jog indicating the zero hour.

**401. High and low water marking device.**—The lower electromagnet is in a circuit that is closed when the platinum index on the time chain (11, fig. 23) is in contact with the fixed platinum index (12); that is to say, at the times of high and low waters. When this contact is made, the electromagnet attracts the armature, which throws the datum-line pen upward, causing a corresponding upward jog in the datum line, and thus automatically marking the time of the high or low water. A small switch (21, fig. 24) just above the hand-crank shaft permits the cutting out of the current from the two electromagnets.

**402. Adjustment of machine.**—The adjustment of the machine should be tested at least once each year and at any other time when there is any reason for believing that a change may have taken place. The following adjustments are required.

**403. Height-chain adjustment.**—All amplitudes should be set at zero, so that the turning of each constituent crank shaft will produce no motion in the height chain. This should bring the summation wheel to its zero position, but on account of a certain amount of backlash and flexures in the machine this wheel may not be in an exact zero position even when the chain is in adjustment. Now, set a single constituent with a very small amplitude and operating the machine with the hand crank, note whether the index of the summation wheel oscillates equal distances on both sides of its zero position. If not, the chain should be adjusted by the adjusting nut at its fixed end at the back part of the machine.

**404. Time-chain adjustment.**—The adjustment of the time chain is similar to that of the height chain. The zero position is indicated by the conjunction of a small triangular-shaped index on the chain and a fixed platinum index in the middle of the horizontal opening in the dial face. A small amplitude being set on one of the constituent time cranks and the machine operated by the hand crank, the chain index should oscillate equal distances on both sides of the platinum point. If it does not, the necessary adjustment may be made at the fixed end of the chain.

**405. Hour-hand adjustment.**—This must be so adjusted that it will register the exact hour at the same instant the circuit for the electromagnet is closed for the hour mark on the marigram, which is indicated by a downward stroke of the datum-line pen. It is also neces-

sary that the zero hour or beginning of the day shall correspond to the double hour mark on the marigram. This adjustment may be accomplished by moving the hour hand on its shaft after releasing its set screw. A finer adjustment may be effected by changing the position of the contact spring back of the dial face.

**406. Minute-hand adjustment.**—This is to be adjusted to read zero on the exact hour indicated by the hour hand and the closing of the electric circuit for the hour mark. The adjustment may be accomplished either by moving the minute hand on its shaft after releasing its set screw or by means of the releasable gears on the main vertical shaft of the dial case. The adjustments just described are those which need be made only occasionally. Other adjustments are taken into account each time the machine is set for a station.

**407. Setting predicting machine.**—The time indicators on the face of the machine are first set to represent the exact beginning of the period for which predictions are to be made, which will usually be 0 hour of January 1 of some year. The hour and minute hands should always be brought into place by the turning of the operating crank in order that the adjustment of these hands relative to the electromagnet circuit may not be affected. The date dial may, however, if desired, be set independently, using the binding nut just above the large dial ring for releasing and clamping. If only a small motion of the date dial is necessary, it is generally preferable to set it by the operating crank. The year index should be set to indicate the kind of year.

**408.** In the usual operation of the machine a ratchet prevents the operating crank from being turned backward, but this ratchet may be released when desired by pressing on a button in the side of the machine just above the crank. After the face of the machine has been thus set to register the beginning of the predictions the three main vertical shafts should be clamped to prevent them from turning.

**409. To set the height amplitudes.**—All the constituent cranks on the left or height side of the machine are first turned, by means of the releasable gears on the main vertical shafts, to a vertical position, the cranks of the upper range of constituents pointing downward and those in the lower range upward, in which position all angles will read  $180^\circ$ . For the long-period constituents the cranks can be more quickly brought to the vertical position by drawing out small knobs on the time side of the machine, thus disconnecting the gearing. The cranks are then turned by hand to the desired position and the knobs pushed back into place. The amplitudes may now be set according to the scales attached to the sides of the machine. The crank pin is unclamped by a small milled head wrench and is then moved along its groove until the index at the scale registers the amplitude setting given in Form 445, when it is clamped in this position. If no amplitude is given for any constituent, the corresponding crank must be set at zero.

**410. To set time amplitudes.**—The process is similar to that for the height amplitudes, the cranks on the time side of the machine being first turned to a vertical position with all angles reading  $90^\circ$ . The cranks are to be set with the same amplitudes as were used for the height side, the modified scales automatically taking account of the true differences in the amplitudes. For the constituents Sa and Ssa the amplitudes are set on the height side only.

**411. To set constituent angles.**—After the amplitudes have been set and checked on both sides of the machine the angles are set for the beginning of the period of predictions, these settings being given in Form 445. The angles may be set from either side of the machine, except for constituents  $S_a$  and  $S_{sa}$ , for which there are no dials on the time side, as the readings are the same for both sides. As each constituent angle is set its releasable gear is clamped to the main vertical shaft. After all the angles have been thus set the three main vertical shafts must be unclamped to permit them to turn.

**412. Changing height scale.**—There are three interchangeable height scales, known as the 40-foot, the 20-foot, and the 10-foot scale. The 40-foot ring may also be conveniently used as a 4-foot scale. The scale to be used for any station is indicated in Form 445. In removing a scale from the machine a small button at the top is turned to release the ring, which is then lifted slightly as it is being removed. The desired scale is then placed on the machine and secured in place by a button. Before removing or replacing the height scale it is desirable that the height pointer be set approximately  $45^\circ$  to the left of its zero position in order to interfere least with the removal or replacement of the scale.

**413. The datum or plane of reference.**—The hand-operating crank should be turned forward or backward until the index of the summation wheel on the height side of the machine indicates mean sea level. It must be kept in mind, however, that as the index lines may come in conjunction at each complete rotation of the summation wheel there is a possibility of being misled in regard to the mean sea-level position. When in doubt, the operating crank should be turned forward to obtain a number of conjunctions, the corresponding height dial reading for each being noted. The conjunction that corresponds most closely with the average of such height readings will be the one that applies to the true zero position. Each complete turn of the height summation wheel will cause a change in the height reading of 12 units, 6 units, or 3 units, respectively, according to whether the 40-unit, 20-unit, or 10-unit dial is used. The height hand, which can be released by the milled nut on the face of the machine, may now be set to the scale reading that corresponds to the height of mean sea level above the datum which has been adopted for the predictions, this value being given in Form 445.

**414. The marigram gear.**—There are three gear combinations, designated as the 1:1, 3:2, and 2:1 ratios. The gear ratio to be used for any station is indicated in Form 445. When it is necessary to change the gear ratio, the machine should be first turned to its mean sea-level position. The change is then effected by sliding the lower set of gears horizontally, being careful to hold the upper set with one hand to prevent it from turning when the gears are released. Before engaging the gears in their new ratios the counterpoise for the pen carriage should be brought to a position approximately midway between the limits of its range of motion. The 1:1 ratio is obtained by sliding the lower set of gears as far as possible toward the height side of the machine, thus engaging the innermost gears; the 3:2 ratio by moving these gears toward the time side until the outer gears are engaged, and the 2:1 ratio by engaging the middle gear of each set.

415. In setting up the machine for successive stations there is a mechanical advantage in making the necessary gear changes before setting the new amplitudes if the gear changes are in the order of 2:1, 3:2, 1:1, and after setting the amplitudes if the gear changes are in the reverse order. This precaution will lessen the chances of jamming the curve pen carriage and throwing the height chain off its pulleys when setting the amplitudes.

416. *Inserting paper roll.*—To place the paper on the machine, remove the mandril that is mounted within the dial case near the upper right-hand corner and slip the roll of paper over the mandril, the roll being so placed that the winding is clockwise when viewed from above and when on the machine the paper unwinds from the outer side of the roll. In placing the roll on the mandril care should be taken to see that the small projection on the base of the latter enters the cavity in the wooden core, so that the roll will fit flat against the base. After the mandril with the roll of paper has been returned to the machine and secured in place, the end of the paper is passed around a roller to the face of the machine, across the face, and over the feed roller at the left of the machine. The end is then inserted into the slit in the receiving roller, which is given a few turns to take up the slack paper and make it secure. Before passing the paper over the feeding roller and on the receiving roller these rollers should be released to permit them to turn independently, the release being effected by turning the small milled head on a ratchet stud gear near the base of the feeding roller and by lifting off from the top of the receiving roller the small knob holding the connecting chain. After the paper has been secured to the receiving roller these connections should be restored.

417. *Curve pen adjustment.*—With the machine in its mean sea-level position, the curve pen must be adjusted to bring the pen point on the mean sea-level line as drawn by the base-line pen. This adjustment may be effected by releasing the pen carriage from the operating chain and moving it to the desired position, where it is clamped in place by the binding screw.

418. *Verification of machine settings.*—Each step in the adjustment and setting of the machine should be carefully checked before proceeding with the next step. After the setting of the machine for any station has been completed an excellent check on the work is afforded, if the predictions for the same station for the preceding year are available, by turning the machine backward several days and then comparing the predicted tides with those previously obtained.

419. *Predicting.*—The datum and curve fountain pens are filled and put in place, the electric cut-out switch under the base of the machine closed, and the ratchet of the operating crank set to prevent the machine from being turned backward. If the predicted height of the tide for any given time is desired, the machine may be turned forward until the required time is registered on the time dials and the corresponding height read off of the height dial.

420. If the predicted high and low waters for the year are desired, the operating crank is turned forward until the machine is automatically stopped by the brake at a high or low water. To avoid the strain on the machine due to sudden stops, the operator should watch the small index on the time chain, and as this approaches the fixed index in the center of the opening on the face of the machine, turn the

crank more slowly until the machine is stopped as the indexes come in contact with each other. The time and height may then be read directly from the dials on the face of the machine. The movement of the height pointer before the stopping of the machine and also the tide curve will clearly indicate whether the tide is a high or low water. After the tide has been recorded an inward pressure on the crank handle will release the brake and the machine can be turned forward to the next tide, the process being repeated until all the tides of the year have been predicted and recorded.

#### FORMS USED WITH TIDE-PREDICTING MACHINE

**421. Form 444, standard harmonic constants for predictions (fig. 32).**—This form provides for the compilation of the harmonic con-

DEPARTMENT OF COMMERCE  
U.S. COAST AND GEODETIC SURVEY  
Form No. 444

#### TIDES CURRENTS | STANDARD HARMONIC CONSTANTS FOR PREDICTION

STATION Morro, California | Lat.  $35^{\circ} 22' N.$   
Long.  $120^{\circ} 51' W.$   
Long.  $120.85^{\circ} W.$

COMPONENT	H AMPLITUDE	A EPOCH	$\frac{A}{\pi - \kappa}$	B $\frac{B}{4 \times H}$	C $360^{\circ} - \alpha'$	D $\alpha'$	REMARKS
M <sub>2</sub>	1.227	308.5	+ 9.8	4.91	+ 41.7	- 318.3	Time meridian 120° = 8.0 h.
S <sub>2</sub>	0.320	304.3	+ 1.7	1.28	+ 54.0	- 306.0	Extreme range 8.8 ft.
N <sub>2</sub>	0.260	284.5	+ 14.2	1.04	+ 61.3	- 298.7	Dial 10 mm.
K <sub>1</sub>	1.001	111.2	+ 0.5	4.00	+ 248.3	- 111.7	Marigram gear 3:2.
M <sub>4</sub>	0.053	170.7	+ 19.6	0.21	+ 169.7	- 190.3	Marigram scale 1:20
O <sub>1</sub>	0.608	99.1	+ 9.3	2.43	+ 251.8	- 108.4	A <sub>0</sub> 2.40 ft.
M <sub>6</sub>	0.013	253.7	+ 29.5	0.05	+ 76.8	- 283.2	Permanent current kn.
(MK) <sub>2</sub>					+	-	The DATUM is a plane
S <sub>4</sub>	0.009	176.6	+ 3.4	0.04	+ 180.0	- 180.0	below mean { low water springs
(MN) <sub>2</sub>					+	-	lower low water
J <sub>2</sub>	0.065	285.0	+ 13.6	0.26	+ 61.4	- 298.6	
S <sub>6</sub>	0.006	148.3	+ 5.1	0.02	+ 206.6	- 153.4	
M <sub>2</sub>	0.025	174.2	+ 18.0	0.10	+ 167.8	- 192.2	
(2N) <sub>2</sub>	(0.035)	260.5	+ 18.5	0.14	+ 81.0	- 279.0	
(OO) <sub>1</sub>	(0.026)	123.3	- 8.3	0.10	+ 245.0	- 115.0	
A <sub>2</sub>	(0.009)	306.6	+ 6.0	0.04	+ 47.4	- 312.6	
S <sub>1</sub>					+	-	
M <sub>1</sub>	0.041	132.4	+ 4.9	0.16	+ 222.7	- 137.3	
J <sub>1</sub>	(0.048)	117.2	- 3.8	0.19	+ 246.6	- 113.4	
M <sub>m</sub>					+	-	
S <sub>ea</sub>					+	-	
S <sub>a</sub>					+	-	
MS <sub>1</sub>					+	-	
M <sub>f</sub>					+	-	
P <sub>1</sub>	(0.023)	93.9	+ 13.1	0.09	+ 253.0	- 107.0	
Q <sub>1</sub>	.0.107	98.9	+ 13.7	0.43	+ 247.4	- 112.6	
T <sub>2</sub>	(0.019)	304.5	+ 2.0	0.08	+ 53.5	- 308.5	
R <sub>2</sub>	(0.003)	304.1	+ 1.4	0.01	+ 54.5	- 305.5	
(2Q <sub>1</sub> )	(0.016)	87.1	+ 18.0	0.06	+ 254.9	- 105.1	
P <sub>1</sub>	0.274	107.7	+ 1.2	1.10	+ 251.1	- 108.9	
(2SM) <sub>2</sub>					+	-	
M <sub>3</sub>	0.022	345.8	+ 14.7	0.09	+ 359.5	- 0.5	
L <sub>2</sub>	0.050	307.5	+ 5.5	0.20	+ 47.0	- 313.0	
(2MK) <sub>2</sub>					+	-	
K <sub>2</sub>	0.0103	289.8	+ 1.0	0.41	+ 69.2	- 290.8	
M <sub>8</sub>	0.007	105.6	+ 39.3	0.03	+ 215.1	- 144.9	
(MS) <sub>1</sub>					+	-	

Source of constants from observed hourly heights for 163 days beginning February 13, 1919.

Compiled by LaP. Da. March 29, 1923. Verified by F. J. H. March 29, 1923.  
(Date) (D-12)

FIGURE 32.

stants for use in the prediction of the tides and also for certain permanent preliminary computations to adapt the constants for use with the U. S. Coast and Geodetic Survey tide-predicting machine No. 2. The form is used in a loose-leaf binder.

**422.** The constituents are listed in an order that conforms to the arrangement of the corresponding constituent shafts and cranks on the predicting machine. The accepted amplitudes and epochs are to be given in the columns provided for the purpose. At the bottom of the page a space is provided for indicating the source from which the constants were derived.

**423.** The column of Remarks provides for miscellaneous information pertaining to the predictions. This includes the kind of time in which the predictions are to be given, the approximate extreme range of tide at the place for determining the proper scale to be used, the height dial, the marigram gear, the marigram scale, and the datum to which the predicted heights are to be referred.

**424.** The extreme range may be estimated from the predictions for a preceding year or may be taken approximately as twice the sum of the amplitudes of the harmonic constants. The height dial, marigram gear, and marigram scale which are recommended for use with different extreme ranges are given in the table on page 138.

**425.** The principal hydrographic datums in general use are as follows: Mean low water for the Atlantic and Gulf coasts of the United States and Puerto Rico. Mean lower low water for the Pacific coast of the United States, Canada, and Alaska, and the Hawaiian and Philippine Islands. Approximate low water springs for the rest of the world, with a few exceptions. For use on the predicting machine the datum must be defined by its relation to the mean sea level, and this relation is usually determined from a reduction of the high and low waters.

**426.** Column A of Form 444 is designed for the differences by which the epochs of the constituents are adapted once for all for use with the unmodified Greenwich ( $V_0 + u$ )'s of each year. These differences take account of the longitude of the station and also of the time meridian used for the predictions, and are computed by the formula

$$\kappa' - \kappa = pL - \frac{aS}{15} \quad (466)$$

in which

$\kappa'$ —adapted epoch—true epoch.

$p$ =subscript of constituent, which indicates number of periods in one constituent day. For the long-period constituents Mm, Ssa, Sa, MSf, and Mf,  $p$  should be taken as zero.

$L$ =longitude of station in degrees; + if west, — if east.

$a$ =speed of constituent in degrees per solar hours.

$S$ =longitude of time meridian in degrees; + if west, — if east.

The values of the products  $\frac{aS}{15}$  for the principal time meridians may be taken from table 35. For any time meridian not given in the table the products may be obtained by direct multiplication, taking the values for the constituent speeds ( $a$ ) from table 2.

**427.** Column B is designed for the reduction of the amplitudes to the working scale of the machine. The scale is unity when the 40-

foot height dial is used, 2 for the 20-foot height dial, 4 for the 10-foot height dial, and 10 for a 4-foot height dial. The working scale should be entered at the head of the column and used as a factor with the amplitudes in order to obtain the values for this column.

428. Columns *C* and *D* are designed to contain the adapted epochs in positive and negative forms which may be used additively with the Greenwich ( $V_0 + u$ )'s. It will be found most convenient to compute column *D* first, by applying the difference in column *A* to the  $\kappa$  in the preceding column and entering the result with the negative sign. If the direct application of the difference should give a negative result, this must be subtracted from  $360^\circ$  before entering in column *D*.

DEPARTMENT OF COMMERCE  
U. S. COAST AND GEODETIC SURVEY  
Form No. 445

**TIDES** } SETTINGS FOR TIDE PREDICTING MACHINE  
**CURRENTS** }

STATION Morro, California

YEAR 19.23

COMPONENT	AMPLITUDE SETTING	DIAL SETTING			REMARKS
		JAN. 1, 0 <sup>a</sup>	FEB. 1, 0 <sup>a</sup>	DEC. 31, 24 <sup>b</sup>	
$M_2$	$5.10$	82.9	47.1	183.7	
$S_2$	$1.30$	54	54	54	
$N_2$	$1.10$	214	133	226	
$K_1$	$3.55$	255	286	255	
$M_4$	$0.20$	252	180	94	
$O_1$	$2.00$	287	221	28	
$M_8$	$0.05$	200	93	142	
(MK) <sub>2</sub>	---	---	---	---	
$S_4$	$0.05$	180	180	180	
(MN) <sub>4</sub>	---	---	---	---	
$\mu_2$	$0.25$	33	6	323	
$S_2$	---	---	---	---	
$\mu_2$	$0.10$	251	179	93	
(2N) <sub>2</sub>	$0.15$	345	219	268	
(OO) <sub>1</sub>	$0.05$	40	167	298	
$\lambda_2$	$0.05$	338	293	249	
$S_1$	---	---	---	---	
$M_1$	$0.25$	347	329	217	
$J_1$	$0.15$	140	216	228	
$M_m$	---	---	---	---	
$Ssa$	---	---	---	---	
$Sa$	---	---	---	---	
MSF	---	---	---	---	
MF	---	---	---	---	
$P_1$	$0.05$	219	162	150	
$Q_1$	$0.35$	35	284	47	
$T_2$	$0.10$	56	25	56	
$R_2$	---	---	---	---	
(2Q) <sub>1</sub>	$0.05$	154	358	78	
$P_1$	$1.10$	242	211	242	
(2SM) <sub>2</sub>	---	---	---	---	
$M_2$	$0.10$	61	7	32	
$L_1$	$0.20$	149	158	338	
(2MK) <sub>2</sub>	---	---	---	---	
$K_2$	$0.30$	264	325	264	
$M_2$	$0.05$	20	237	63	
(MS) <sub>4</sub>	---	---	---	---	

Computed by L.P.D. March 28, 1923 Verified by F.J.H. March 29, 1923

Predicted by

Date

FIGURE 33.

The values for column *C* may then be obtained by applying  $360^\circ$  to the negative values in column *D*.

**429.** *Form 445, settings for tide-predicting machine* (fig. 33).—This form is designed for the computations of the settings for the predicting machine for the beginning of each year of predictions. The forms are bound in books, a separate book being used for each year of predictions. This form is used in connection with Form 444, and for convenience the order of arrangement of the constituents is identical in the two forms. The name of the station, the time meridian, the height dial, marigram gear, marigram scale, and datum plane are copied directly from Form 444.

**430.** For the amplitude settings the amplitudes of column *B* of Form 444 are multiplied by the factors *f* from table 14 for the year for which the predictions are to be made. A convenient way to apply these factors is to prepare a strip of paper with the same vertical spacing as the lines on Form 444 and enter the factors *f* for the required year on this strip. The strip may then be placed alongside of column *B* of Form 444 and the multiplication be performed. The same strip will serve for every station for which predictions are to be made for the given year. It has been the recent practice to enter the amplitude settings to the nearest 0.05 foot as being sufficiently close for all practical purposes.

**431.** For the dial settings for January 1, 0 hour, the Greenwich equilibrium arguments of  $(V_o + u)$ 's from table 15 are to be applied, according to the indicated sign, to the angles of column *C* or *D* of Form 444, using the angle in column *D* if it is less than the argument, otherwise using the angle in column *C*. For the application of the  $(V_o + u)$ 's a strip similar to that used for the factors *f* should be prepared. The same strip will serve for all stations for the given year. For the dial settings it is customary to use whole degrees, except for constituent  $M_2$ , for which the setting is carried to the first decimal of a degree.

**432.** The settings for February 1 and December 31 are used for checking purposes to ascertain whether there has been any slipping of the gears during the operation of the machine. To obtain the dial settings for February 1, 0<sup>h</sup>, and December 31, 24<sup>h</sup>, prepare strips similar to those for the *f*'s and  $(V_o + u)$ 's. On one enter the angular motion of the constituents from January 1, 0<sup>h</sup>, to February 1, 0<sup>h</sup>; on a second and a third strip, the angular motion for February 1, 0<sup>h</sup>, to December 31, 24<sup>h</sup>, for a common and leap year, respectively. For checking purposes a fourth and fifth strip may contain the angular changes for a complete common and a complete leap year, respectively. The values for these strips may be obtained from table 36. These strips will be found more convenient if arranged with two columns each, one column containing the values in a positive form and the other column containing the equivalent negative value which is obtained by subtracting the first from  $360^\circ$ . These strips are good for all years, distinction being made between the common and leap years. By applying the first strip to the dial settings for January 1 the values for February 1 are readily obtained, and by applying the second or third strip to the latter settings those for the end of the year are obtained. The values obtained by applying the fourth or fifth strips to the settings for January 1 should also give the correct setting for the end of the year, and thus serve as a check. The

angular changes for computing the settings for any day of the year may be obtained from tables 16 and 17.

#### PREDICTION OF TIDAL CURRENTS

**433.** Since the tidal current velocities in any locality may be expressed by the sum of a series of harmonic terms involving the same periodic constituents that are found in the tides, the tide-predicting machine may be used for their prediction. For the currents, however, consideration must be given to the direction of flow, and in the use of the machine some particular direction must be assumed. At present the machine is used for the prediction of reversing currents in which the direction of the flood current is taken as positive and the maximum velocity in this direction corresponds to the high water of the predicted tide. The ebb current is then considered as having a negative velocity with its maximum corresponding to the low water of the predicted tide. Rotary currents may be predicted by taking the north and east components separately but the labor of obtaining the resultant velocities and directions from these components would be very great without a machine especially designed for the purpose. Predictions can, however, be made along the main axis of a rotary movement without serious difficulties. Formulas for referring the harmonic constants of the north and east components to any desired axis are given in Coast and Geodetic Survey Special Publication No. 215, Manual of Current Observations.

**434.** The harmonic constants for the prediction of current velocities are derived from current observations by an analysis similar to that used in obtaining the harmonic constants from tide observations. In the current harmonic constants, however, the amplitudes are expressed in a unit of velocity, usually the knot, instead of the linear unit that is used for the tidal harmonic constants. Forms 444 and 445 for the computation of the settings for the tide-predicting machine are applicable for the current predictions and the procedure in filling out these forms is essentially the same as described in paragraphs 421-432 for the tide predictions. The node factors ( $f$ ) and arguments ( $V_0 + u$ ) are the same as for the tides. The height dial, marigram gear and scale suitable to the current velocity can be obtained from the table on page 138. Instead of a sea level elevation there should be entered in the column of "Remarks" the velocity of any permanent current along the axis in which the predictions are to be made. This velocity should be marked plus (+) or minus (-) according to whether the permanent current is in the flood or ebb direction.

**435.** The predicting machine is set with the current harmonic constants in the same manner as for the tidal harmonic constants. To take account of the permanent current the height summation wheel should be brought to its zero position and the height hand then set at a dial reading corresponding to the velocity of the permanent current, the hand being set to the right of the scale zero if the permanent current is in the flood direction and to the left if in the ebb direction. The hand crank should be then turned to bring the height hand to its zero position and the curve-pen set at the medial line of the paper, this line now representing zero velocity or slack water.

**436.** The operation of the machine for the prediction of the currents is similar to that for the prediction of the tides. The machine automatically stops at each maximum flood and ebb velocity and the corresponding times and velocities are then recorded, the flood velocities being read to the right and the ebb velocities to the left of the scale zero. In the prediction of the currents the times of slack water are also desired. These are indicated by the zero position of the recording hand as well as by the intersections of the curve and medial line in the graphic record. The velocity of the current at any intermediate time can be read directly from the height dial when the machine has been turned to the time desired and it may be also scaled from the graphic record.

**437.** Predictions of hydraulic currents in a strait, based upon the difference in the tidal head at the two entrances, may be made by means of harmonic constants derived from the tidal constants for the entrances. Differences in tidal range or in the times of the high and low waters at the two ends of a strait will cause the water surface at one end alternately to rise above and fall below that at the other end, thus creating a periodic reversing current in the strait. Theoretically, disregarding friction or inertia, the velocity of the current would vary as the square root of the difference in head, being zero when the surface is at the same level at both ends and reaching a maximum when the difference is greatest. Actually there will generally be a lag of some minutes in the response of the current movement to the difference in head which must be determined from observations.

**438.** Let the two ends of the strait be designated by *A* and *B*, with the flow from *A* toward *B* considered as flood or positive and the flow in the opposite direction as ebb or negative. With the waterway receiving the tide from two sources, the application of the terms "flood" and "ebb" will be somewhat arbitrary, and care must be taken to indicate clearly the direction assumed for the flood movement. In the following discussion tidal constants pertaining to entrances *A* and *B* will be distinguished by subscripts *a* and *b*, respectively, and those pertaining to the difference in tidal head by the subscript *d*. Since the usual constituent epochs known as "kappas" refer to the local meridian, it will be necessary for the purpose of comparison between places on different meridians to use the Greenwich epochs "*G*" (par. 226), these being independent of local time and longitude.

**439.** For any one constituent let *T* represent time as expressed in degrees of the constituent reckoned from the phase zero of its Greenwich equilibrium argument. Also let  $Y_a$  and  $Y_b$  represent the height of the constituent tide for any time *T* as referred to the mean level at locations *A* and *B*, respectively; and let  $Y_d$  equal the difference ( $Y_a - Y_b$ ). Formulas for heights and difference may now be written

$$Y_a = H_a \cos (T - G_a) \text{ for location "A"} \quad (467)$$

$$Y_b = H_b \cos (T - G_b) \text{ for location "B"} \quad (468)$$

$$\begin{aligned} Y_d &= H_a \cos (T - G_a) - H_b \cos (T - G_b) \\ &= (H_a \cos G_a - H_b \cos G_b) \cos T + (H_a \sin G_a - H_b \sin G_b) \sin T \\ &= H_d \cos (T - G_d) \end{aligned} \quad (469)$$

in which

$$H_d = [H_a^2 + H_b^2 - 2H_a H_b \cos (G_b - G_a)]^{\frac{1}{2}} \quad (470)$$

$$G_d = \tan^{-1} \frac{H_a \sin G_a - H_b \sin G_b}{H_a \cos G_a - H_b \cos G_b} \quad (471)$$

The proper quadrant for  $G_d$  is determined by the signs of the numerator and denominator of the above fraction, these being the same, respectively, as for the sine and cosine of the angle. Formulas (470) and (471) may be solved graphically (fig. 34) by drawing from any point  $C$  a line  $CD$  to represent in length and direction  $H_a$  and  $G_a$ , respectively; from the point  $D$  a line  $DE$  to represent in length and direction  $H_b$  and  $(G_b \pm 180^\circ)$ , respectively. The connecting line from




FIGURE 34.

$C$  to  $E$  will represent by its length the amplitude  $H_d$  and by its direction the epoch  $G_d$ .

**440.** Formulas (470) and (471) may be modified to adapt them for use with tables 41 and 42. From (470) we may obtain

$$H_d/H_a = [1 + (H_b/H_a)^2 + 2(H_b/H_a) \cos (G_b - G_a \pm 180^\circ)]^{\frac{1}{2}} \quad (472)$$

or

$$H_d/H_b = [1 + (H_a/H_b)^2 + 2(H_a/H_b) \cos (G_a - G_b \pm 180^\circ)]^{\frac{1}{2}} \quad (473)$$

and from (471) we have

$$\tan (G_d - G_a) = \frac{(H_b/H_a) \sin (G_b - G_a \pm 180^\circ)}{1 + (H_b/H_a) \cos (G_b - G_a \pm 180^\circ)} \quad (474)$$

or

$$\tan(G_a - G_b \pm 180^\circ) = \frac{(H_a/H_b) \sin(G_a - G_b \pm 180^\circ)}{1 + (H_a/H_b) \cos(G_a - G_b \pm 180^\circ)} \quad (475)$$

Formulas (472) and (474) are to be used when the ratio  $H_b/H_a$  does not exceed unity. In this case take argument  $r$  of the tables =  $H_b/H_a$ , and argument  $x = (G_b - G_a \pm 180^\circ)$ . If the ratio  $H_b/H_a$  exceeds unity use formulas (473) and (475) and take argument  $r = H_a/H_b$  and argument  $x = (G_a - G_b \pm 180^\circ)$ . The tabular values will give the ratios and angular differences represented in the first terms of the formulas. Therefore, in order to obtain the amplitude  $H_d$ , the tabular value from table 41 must be multiplied by  $H_a$  if the ratio  $H_b/H_a$  does not exceed unity, or by  $H_b$  if this ratio does exceed unity. Also to obtain the epoch  $G_d$ , the tabular value from table 42 must be increased by  $G_a$  if the ratio does not exceed unity or by  $(G_b \pm 180^\circ)$  if the ratio is greater than unity.

**441.** By the formulas given above separate computations are made for each of the tidal constituents. The values obtained for  $H_d$  and  $G_d$  are the corresponding amplitudes and Greenwich epochs in an harmonic expression for the continually changing difference in elevation of the water surface at the two entrances to the strait. When only a single time zone is involved, the small  $g$ 's or modified kappas ( $\kappa'$ ) pertaining to that zone may be substituted for the Greenwich epochs ( $G$ ) in the formulas. For the prediction of the current, further modifications are necessary in the amplitudes to reduce to velocity units and in the epochs to allow for the lag in the response of the current to the changing difference in water level at the two entrances to the strait.

**442.** Since the velocity of an hydraulic current is theoretically proportional to the square root of the difference in head, we may write

$$(\text{Velocity})^2 = \text{constant } (C) \times \text{height difference} \quad (476)$$

If we now let  $V$  equal the average velocity of the current at time of strength as determined from actual observations and assume that the corresponding difference in water level is 1.02 times the difference resulting from the principal constituent  $M_2$ , we may obtain an approximate value for the constant ( $C$ ) by the formula

$$C = V^2 / (1.02M_2) \quad (477)$$

in which  $M_2$  is the amplitude of the constituent  $M_2$  in the harmonic expression for the difference in head. The application of the factor ( $C$ ) to all the constituent amplitudes in this expression has the effect of changing the height units into units representing the square of the velocity of the resulting current.

**443.** The lag in the current is usually determined by a comparison of the times of strengths and slacks from actual observations with preliminary predictions of the corresponding phases based upon the harmonic constants derived by the method just described. This lag expressed in hours is multiplied successively by the speed of each constituent and the result applied to the preliminary epoch for that constituent.

**444.** In order that the magnitude of the constituent amplitudes may be adapted for use with the predicting machine, a scale factor ( $S$ ) is

introduced. This factor, which depends upon the velocity of the current, is selected with the view of obtaining a reasonably large working scale without exceeding the limitations of the predicting machine. The following scale factors are suggested:

Average velocity of current at time of strength:	Scale factor
Less than 0.3 knot	20
From 0.3 to 0.5 knot	10
From 0.5 to 1.0 knot	5
From 1.0 to 1.5 knots	3
From 1.5 to 2.0 knots	2
From 2.0 to 3.0 knots	1
From 3.0 to 4.0 knots	0.5
From 4.0 to 5.0 knots	0.25
From 5.0 to 10.0 knots	0.1

In practice, the scale factor is usually combined with the factor ( $C$ ) and the product applied to each of the constituent amplitudes in the expression for difference in head.

**445.** Using the harmonic constants, modified in the manner described above, in the predicting machine, the resulting dial readings will represent the square of the current velocity. In order to avoid the necessity of extracting the square root of each individual reading, a square-root scale may be improvised and substituted for the regular height dial on the machine. From a consideration of the construction of this machine, it can be shown that with a scale factor of unity the angular position of a velocity graduation as measured in degrees from the zero point will be  $9^\circ \times (\text{velocity})^2$ . Thus the 1-knot graduation will be spaced  $9^\circ$  from the zero, the 2-knot graduation at  $36^\circ$ , the 3-knot graduation at  $81^\circ$ , etc. For any scale factor ( $S$ ), the formula for constructing the square-root scale becomes

$$\text{Angular distance from dial zero} = 9^\circ \times S \times (\text{velocity})^2 \quad (478)$$

**446.** To take account of any nontidal current not attributed to difference in head at the two entrances to the strait, a special graduation of the square-root scale is necessary. Let  $V_0$  represent the nontidal current velocity, positive or negative according to whether it sets in the flood or ebb direction, and let  $V$  represent the resultant velocity as indicated by any scale graduation, positive or negative according to whether it is flood or ebb. The angular distance of any scale graduation as measured from an initial point, usually marked by an arrow, may then be expressed by the following formula:

$$\text{Angle in degrees} = 9 \times S \times (V - V_0)^2 \quad (479)$$

The required angle is to be measured to the right or to the left of the initial point according to whether the angle  $(V - V_0)$  is positive or negative. When setting the predicting machine the velocity pointer must be at the initial point marked by the arrow when the sum of the harmonic terms is zero.

**447.** In the graphic representation of the summation of the harmonic terms by the predicting machine, the scale of the marigram depends upon the marigram gear ratio as well as upon any scale factor which may have been introduced. With a gear ratio of unity, the scale of the marigram is 0.1 inch per unit of machine setting. In the summation for the hydraulic currents, the marigram read by a natural

scale would indicate the square of the velocity. A special square-root reading scale for taking the velocities direct from the marigram may be prepared as follows: Let  $Y$ =distance of any velocity graduation from zero of scale. Then

$$Y \text{ (in inches)} = 0.1 \times (\text{scale factor}) \times (\text{gear ratio}) \times (\text{velocity})^2 \quad (480)$$

With the scale factor and gear ratio each unity, 1 knot of velocity would be represented by 0.1 inch on the marigram, 2 knots by 0.4 inch, 3 knots by 0.9 inch, etc. With a scale thus constructed the velocity of the tidal current may be taken directly from the marigram. Any nontidal current which is to be included may afterward be applied.

**TABLES**  
**EXPLANATION OF TABLES**

Table 1. *Fundamental astronomical data.*—This table includes fundamental constants and formulas with references which form the basis for the computation of other tables contained in this volume. Because of the smallness of the solar and lunar parallax no distinction is made between the parallax and its sine. The eccentricity of the earth's orbit and the obliquity of the ecliptic are given for epoch January 1, 1900. The former changes about 0.000042 per century and the latter about 0.013 of a degree in a century. The values given may therefore be considered as applicable to the present century.

The formulas for longitude of both sun and moon are the same as used in the previous edition of this book and are from the work of Simon Newcomb. In a later work by Ernest W. Brown, slightly different values are obtained for the elements of the moon's orbit but the differences may be considered negligible in so far as the tidal work of the present century is concerned. In these formulas it will be noted that  $T$  is the number of Julian centuries reckoned from Greenwich mean noon on December 31, 1899, of the Gregorian calendar which corresponds to December 19, 1899, by the Julian calendar. In the application of these formulas to early dates special care must be taken to make suitable allowances for the particular calendar in use at the time. See page 4 for information in regard to calendars.

Table 1 includes the numerical values of the mean longitude of the solar and lunar elements for the beginning of the century years 1600 to 2000 and also the rate of change in these longitudes as of January 1, 1900. As the variations in these rates are very small, they are applicable without material error for all modern times. This table includes also the principal astronomical periods depending on the solar and lunar elements with formulas showing how they are derived. In these formulas the longitude symbol is used to represent its own rate of change according to the unit in which the period is expressed.

Table 2. *Harmonic constituents.*—This table includes the arguments, speeds, and coefficients of the constituent harmonic terms obtained in the development of the tide-producing forces of the moon and sun. They are grouped with reference to the formulas of the text from which they are derived, the long-period constituents first, followed by the diurnal, semidiurnal, and terdiurnal terms. The reference numbers in the first column correspond to the numbered terms of the formulas of the text, the letter  $A$  indicating a term in the lunar development and the letter  $B$  a term in the solar development. In the second column the usual symbols are given for the principal constituents, parentheses being used when the term only partially represents the constituent.

For an explanation of the constituent argument (E) see page 22. The argument consists of two parts—the  $V$  which contains the

uniformly changing elements and determines the speed and period of the constituent, and the  $u$  which is a function of the moon's node with slow variations and which is treated as a constant for a limited series of observations. Because of the very small change in the element  $p_1$  it may for practical purposes also be treated as a constant with a mean value of  $282^\circ$  for the present century.

The constituent speeds are obtained by adding the hourly rates of change in the elements appearing in the  $V$  of the arguments. The hour angle ( $T$ ) of the mean sun changes at the rate of  $15^\circ$  per hour. The hourly rate of change for each of the other elements will be found in table 1.

For an explanation of the constituent coefficients (C) see page 24. The coefficients of the solar terms include the solar factor  $S'$  (paragraph 118), and coefficients of the lunar terms involving the 4th power of the moon's parallax include the factor  $a/c$  (paragraph 108); in order that all terms may be comparable when used with the common basic factor  $U$ . It is to be noted that in the present system of coefficients for the terms of the principal tide-producing force there is included a factor "2" which was formerly incorporated in the general coefficient. For the terms involving the 4th power of the parallax there is a corresponding factor of "3" in order that all terms may be comparable in respect to the vertical component force.

In general the coefficients have been computed in accordance with the coefficient formulas of the text, but exceptions were made for the evectional and variational constituents  $\rho_1$ ,  $\nu_2$ ,  $\lambda_2$ , and  $\mu_2$ , the coefficients of which are based upon computations by Professor J. C. Adams who was associated with Darwin in the investigation of harmonic analysis and who carried the development of the lunar theory to a higher order of precision than is provided in this work. (See pp. 60-61 of the Report of British Association for the Advancement of Science for year 1883.)

The node factor ( $f$ ) is explained on page 25. The last column of table 2 contains references to the formulas for the node factors of the various constituents.

Table 2a. *Shallow-water constituents*.—In this table there are listed the overtides and compound tides which are described on page 47.

Table 3. *Latitude factors*.—This table includes numerical values of the latitude ( $Y$ ) functions which appear in the text as factors in formulas representing component tidal forces and the equilibrium height of the tide. The combination symbol at the head of each column is taken to suggest the formula to which it applies. Thus, the letters  $v$ ,  $s$ , and  $w$  refer respectively to the vertical, south, and west components of the force, the letter  $v$  being applicable also to the formulas for the equilibrium height of the tide which have the same latitude factors as the corresponding terms in the vertical component of the force. The first numeral "3" or "4" indicates whether the formula is from the development of the principal force involving the cube of the parallax or from the development involving the 4th power of the parallax of the tide-producing body. The last digit "0," "1," "2," or "3" refers to the species of the constituents and indicates whether they are long-period, diurnal, semidiurnal, or terdiurnal. In several cases the same latitude factor is applicable to a number of different groups as indicated at the head of the column in the table.

The following formulas were used in computing the latitude factors. The maximum value (irrespective of sign) with corresponding latitude is also given for each function.

$Y_{v30} = (1/2 - 3/2 \sin^2 Y)$	maximum - 1 when $Y = \pm 90^\circ$ .
$Y_{v31} = \sin 2Y$	maximum $\pm 1$ when $Y = \pm 45^\circ$ .
$Y_{s30}, Y_{s32}$ , and $Y_{w42}$ , same	as $Y_{v31}$
$Y_{v32} = \cos^2 Y$	maximum + 1 when $Y = 0$ .
$Y_{w43}$ same as $Y_{v32}$	
$Y_{s31} = \cos 2Y$	maximum $\pm 1$ when $Y = 0$ or $\pm 90^\circ$ .
$Y_{w31} = \sin Y$	maximum $\pm 1$ when $Y = \pm 90^\circ$ .
$Y_{w32} = \cos Y$	maximum + 1 when $Y = 0$ .
$Y_{v40} = \sin Y (\cos^2 Y - 2/5)$	maximum $\mp 0.4$ when $Y = \pm 90^\circ$ .
$Y_{v41} = \cos Y (\cos^2 Y - 4/5)$	maximum - 0.2754 when $Y = \pm 58.91^\circ$ .
$Y_{s40}$ same as $Y_{v41}$	
$Y_{v42} = \sin Y \cos^2 Y$	maximum $\pm 0.3849$ when $Y = \pm 35.26^\circ$ .
$Y_{s43}$ same as $Y_{v42}$	
$Y_{v43} = \cos^3 Y$	maximum + 1 when $Y = 0$ .
$Y_{s41} = \sin Y (\cos^2 Y - 4/15)$	maximum $\mp 0.2667$ when $Y = \pm 90^\circ$ .
$Y_{s42} = \cos Y (\cos^2 Y - 2/3)$	maximum - 0.2095 when $Y = \pm 61.87^\circ$ .
$Y_{w41} = (\cos^2 Y - 4/5)$	maximum - 0.8 when $Y = \pm 90^\circ$ .

Table 4. *Mean longitude of lunar and solar elements.*—This table contains the mean longitude of the moon ( $s$ ), of the lunar perigee ( $p$ ), of the sun ( $h$ ), of the solar perigee ( $p_1$ ), and of the moon's ascending node ( $N$ ), for January 1, 0 hour, Greenwich mean civil time, for each year from 1800 to 2000, the dates referring to the Gregorian calendar.

These values are readily derived from table 1, the rate of change in the mean longitude of the elements for the epoch January 1, 1900, being applicable without material error to any time within the two centuries 1800 to 2000 covered by table 4. The same rate of change may also be used, without introducing any errors of practical importance, to extend table 4 to dates beyond these limits. In extending the table, care should be taken to distinguish between the common and leap years, and for the earlier dates due consideration should be given to the kind of calendar in use. (See p. 4 for discussion of calendars.) It will be noted that each Julian century contains 36,525 days, while the common Gregorian century contains only 36,524 days, with an additional day every fourth century.

Table 5. *Differences to adapt table 4 to any month, day, and hour.*—These differences are derived from the daily and hourly rate of change of the elements as given in table 1, multiples of  $360^\circ$  being rejected when they occur. The table is prepared especially for common years, but is applicable to leap years by increasing the given date by one day if it is between March 1 and December 31, inclusive. The correction for the hour of the day refers to the Greenwich hour, and if the hour for which the elements are desired is expressed in another kind of time the equivalent Greenwich hour must be used for the table.

Table 6. *Values of  $I$ ,  $v$ ,  $\xi$ ,  $v'$ , and  $2v''$  for each degree of  $N$ .*—Referring to figure 1 (page 6), note that by construction arc  $\Omega\Upsilon'$  equals arc  $\Omega\Upsilon$ . Then in the spherical triangle  $\Omega\Upsilon A$ , the three sides are  $N$ ,  $v$ , and  $(N-\xi)$ , and the opposite angles are respectively  $(180^\circ - I)$ ,  $i$ , and  $\omega$ .

Therefore we have the following relations which may be used in computing the values of  $I$ ,  $\nu$ , and  $\xi$  in the table:

$$\begin{aligned}\cos I &= \cos i \cos \omega - \sin i \sin \omega \cos N \\ &= 0.91370 - 0.03569 \cos N\end{aligned}$$

$$\tan \frac{1}{2}(N - \xi + \nu) = \frac{\cos \frac{1}{2}(\omega - i)}{\cos \frac{1}{2}(\omega + i)} \tan \frac{1}{2}N = 1.01883 \tan \frac{1}{2}N$$

$$\tan \frac{1}{2}(N - \xi - \nu) = \frac{\sin \frac{1}{2}(\omega - i)}{\sin \frac{1}{2}(\omega + i)} \tan \frac{1}{2}N = 0.64412 \tan \frac{1}{2}N$$

For the computation of  $\nu'$  and  $2\nu''$ , formulas (224) and (232) on pages 45-46 may be used. The tabular values themselves were taken from the preceding edition of this work where they were based upon formulas differing slightly from those given here but any differences arising from the use of the latter may be considered as negligible.

Table 7. *Values of  $\log R_a$  for amplitude of constituent  $L_2$ .*—Values in this table are based upon formula (213) on page 44.

Table 8. *Values of  $R$  for argument of constituent  $L_2$ .*—Values in this table are derived from formula (214) on page 44.

Table 9. *Values of  $\log Q_a$  for amplitude of constituent  $M_1$ .*—Values in this table are based upon formula (197) on page 41.

Table 10. *Values of  $Q$  for argument of constituent  $M_1$ .*—Values in this table are derived from formula (203) on page 42.

Table 11. *Values of  $u$  for equilibrium arguments.*—This table is based upon the  $u$ -formulas in table 2 and includes values for the principal lunar constituents for each degree of  $N$ . The  $u$ 's of  $L_2$  and  $M_1$ , which are functions of both  $N$  and  $P$  are given separately in table 13 for the years 1900 to 2000.

Table 12. *Log factor  $F$  for each degree of  $I$ .*—The factor  $F$  is the reciprocal of the node factor  $f$  to which references are given in table 2. The values in table 12 are based upon the formulas for these factors and are given for all the lunar constituents used in the tide-predicting machine, excepting values for  $L_2$  and  $M_1$  which are given separately in table 13.

Table 13. *Values of  $u$  and  $\log F$  for  $L_2$  and  $M_1$ .*—From a comparison of the  $u$ 's of constituents  $L_2$ ,  $M_1$ , and  $M_2$  in table 2, it will be noted that the following relations exist:

$$\begin{aligned}u \text{ of } L_2 &= (u \text{ of } M_2) - R \\ u \text{ of } M_1 &= \frac{1}{2}(u \text{ of } M_2) + Q\end{aligned}$$

Also, the following relations may be derived from formula (215) on page 44 and formula (207) on page 43 since the factor  $F$  is the reciprocal of the node factor  $f$ :

$$\begin{aligned}\log F(L_2) &= \log F(M_2) + \log R_a \\ \log F(M_1) &= \log F(O_1) + \log Q_a\end{aligned}$$

The values for table 13 were computed by the above formulas, the component parts being taken from tables 7 to 12, inclusive. The values for  $\log F(M_1)$  in this table are in accord with Darwin's original

formula from which a factor of approximately 1.5 was inadvertently omitted (see page 43).

Table 14. *Node factor f for middle of each year 1850 to 1999.*—The factor  $f$  is the reciprocal of factor  $F$ . The values for the years 1850 to 1950 were taken directly from the Manual of Tides, by R. A. Harris, and the values for 1951 to 1999 were derived from tables 12 and 13.

Table 15. *Equilibrium argument ( $V_o+u$ ) for beginning of each year 1850 to 2000.*—The equilibrium argument is discussed on page 22. The tabular values are computed by the formulas for the argument in table 2, the  $V_o$  referring to the value of  $V$  on January 1, 0 hour Greenwich mean civil time, for each year, and the  $u$  referring to the middle of the same calendar year; that is, Greenwich noon on July 2 in common years and the preceding midnight in leap years. The value of the  $T$  of the formulas is  $180^\circ$  for each midnight, and the values of the other elements for the  $V$  may be obtained from table 4. The  $u$  of the argument may be obtained from tables 11 and 13 after the value of  $N$  has been determined for the middle of each year from tables 4 and 5. In constructing table 15 the values for the years 1850 to 1950 were taken directly from the Manual of Tides, by R. A. Harris, and the values for the years 1951 to 2000 were computed as indicated above.

Tables 16, 17, and 18.—These tables give the differences to adapt table 15 to any month, day, and hour, and are computed from the hourly speeds of the constituents as given in table 2. The differences refer to the uniformly varying portion  $V$  of the argument, it being assumed that for practical purposes the portion  $u$  is constant for the entire year.

The approximate Greenwich ( $V_o+u$ ) for any desired Greenwich hour may be obtained by applying the appropriate differences from tables 16, 17, and 18 to the value for the first of January of the required year, as given in table 15. To refer this Greenwich ( $V_o+u$ ) to any local meridian, it is necessary to apply a further correction equal to the product of the longitude in degrees by the subscript of the constituent, which represents the number of periods in a constituent day. West longitude is to be considered as positive and east longitude as negative, and the subscripts of the long-period constituents are to be taken as zero. This correction is to be subtracted.

The ( $V_o+u$ ) obtained as above will, in general, differ by a small amount from the value as computed by Form 244, because in the former case the  $u$  refers to the middle of the calendar year and in the latter case to the middle of the series of observations.

Table 19. *Products for Form 194.*—This is a multiplication table especially adapted for use with Form 194, the multipliers being the sines of multiples of  $15^\circ$ .

Table 20. *Augmenting factors.*—A discussion of augmenting factors is given on page 71. The tabular values for the short-period constituents are obtained by formulas (308) and (309) on page 72, and those for the long-period constituents by formulas (403) and (404) on page 92. For constituents  $S_1$ ,  $S_2$ , etc. the augmenting factor is unity.

Tables 21 to 26.—These tables represent perturbations in  $K_1$  and  $S_2$  due to other constituents of nearly equal speeds. They are based upon formulas (359) to (364), inclusive, on page 83.

Table 27. *Critical logarithms for Form 245.*—This table was designed for quickly obtaining the natural numbers to three decimal places for column (3) of Form 245 from the logarithms of column (2). The logarithms are given for every change of 0.001 in the natural number. Each logarithm given in this table is derived from the natural number that is 0.0005 less than the tabular number to which it applies. Intermediate logarithms, therefore, apply to the same natural number as the preceding tabular logarithm. For example, logarithms less than 6.6990 apply to the natural number 0.000 and logarithms from 6.6990 to 7.1760 apply to the natural number 0.001, etc.

Table 28. *Constituent speed differences.*—The constituent speeds as given in table 2 were used in the computation of this table.

Table 29. *Elimination factors.*—These tables provide for certain constant factors in formulas (389) and (390). Separate tables for each length of series and different values for each term of the formulas are required. The tabular values are arranged in groups of three, determined as follows:

$$\text{First value} = \text{logarithm of } \frac{180}{\pi} \frac{\sin \frac{1}{2}(b-a)\tau}{\frac{1}{2}(b-a)\tau}$$

Second value = natural number  $\frac{180}{\pi} \frac{\sin \frac{1}{2}(b-a)\tau}{\frac{1}{2}(b-a)\tau}$  always taken as positive.

$$\text{Third value} = \frac{1}{2}(b-a)\tau, \text{ if } \frac{\sin \frac{1}{2}(b-a)\tau}{\frac{1}{2}(b-a)\tau} \text{ is positive,}$$

$$\text{or } \frac{1}{2}(b-a)\tau \pm 180, \text{ if } \frac{\sin \frac{1}{2}(b-a)\tau}{\frac{1}{2}(b-a)\tau} \text{ is negative.}$$

Table 30. *Products for Form 245.*—This table is designed for obtaining the products for columns (6) and (7) of Form 245.

Table 31. *For construction of primary stencils.*—This table gives the differences to be applied to the solar hours in order to obtain the constituent hours to which they most nearly coincide. Each difference applies to several successive solar hours, but for brevity only the first solar hour of each group to which the difference applies is given in the table.

An asterisk (\*) indicates that the solar hour so marked is to be used twice or rejected according to whether the constituent speed is greater or less than  $15p$ , when in the summation it is desired to assign a single solar hour to each successive constituent hour. For the usual summations in which each solar hour height is assigned to the nearest constituent hour no attention need be given to the asterisk.

The table is computed by substituting successive integral values for  $d$  in formula (243) and reducing the resulting solar hour of series (*shs*) to the corresponding day and hour. The solar hour to be tabulated is the integral hour that immediately follows the value of (*shs*) of the formula. If the fractional part of (*shs*) exceeds 0.5, the tabular solar hour is marked by an asterisk (\*). The successive values of  $d$ , although used positively in formula (243), are to be considered as negative in the application of the table when the speed of the constituent is less than  $15p$ . When the constituent speed is greater than  $15p$ , the difference is to be taken as positive. All tabular differences are brought within the limits +24 hours and -24 hours by rejecting multiples of  $\pm 24$  hours when necessary, and for convenience in use all differences are given in both positive and negative forms.

The following example will illustrate the use of the table: To find constituent 2Q hours corresponding to solar hours 12 to 23 on 16th day of series. By the table we see that solar hour 12 of the 16th day of series is within the group beginning on solar hour 8 of the same day with the tabular difference of +19 or -5 hours, and that the difference changes by -1 hour on solar hours 15 and 21, the latter being marked by an asterisk. Applying the differences indicated, we have for these solar hours on the 16th day of series:

Solar hour	12, 13, 14*, 15, 16, 17, 18, 19, 20, 21*, 22, 23
Difference	-5 -5 -5 -6 -6 -6 -6 -6 -6 -7 -7 -7

Constituent	
2Q hour	7, 8, 9*, 9, 10, 11, 12, 13, 14, 14*, 15, 16

In the results it will be noted that the constituent hours 9 and 14 are each represented by two solar hours. If it should be desired to limit the representation to a single solar hour each, the hours marked with the asterisk should be rejected.

To find constituent OO hours corresponding to solar hours 0 to 18 on the 22d day of series. The 0 hour of the 22d day is in the group beginning on solar hour 14 of the preceding day with the tabular difference of +14 or -10 hours, and changes of +1 hour in the differences occur on solar hours 3 and 17 of the 22d day. It will be noted that the hour 3 is marked by an asterisk. Applying the differences from the table as indicated, we have for the 22d day of series:

Solar hours	0, 1, 2, 3*, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16*, 17, 18
Differences	+14, +14, +14, +15, +15, +15, +15, +15, +15, -9, -9, -9, -9, -9, -9, -9, -8, -8

Constituent	
OO hours	14, 15, 16, 18, 19, 20, 21, 22, 23, 0, 1, 2, 3, 4, 5, 6, 7, 9, 10

In the results it will be noted that constituent hours 17 and 8 are missing. If it is desired to have each of these hours represented also, the solar hours marked by asterisks will be used again. In this table the constituents have been arranged in accordance with the length of the constituent day.

Table 32. *Divisors for primary stencil sums.*—This table contains the number of solar hourly heights included in each constituent hour group for each of the standard length of series when all the hourly heights have been used in the summation.

Table 33. *For construction of secondary stencils.*—Constituent *A* is the constituent for which the original primary summations have been made, and constituent *B* is the constituent for which the sums are to be derived by the secondary stencils. The "Page" refers to the page of the original tabulations of the hourly heights in Form 362. The differences in this table were calculated by formula (252), and the corresponding "Constituent *A* hours" from formula (250), *m* being assigned successive values from 1 to 24 for each page of record. Special allowance was made for page 53 of the record to take account of the fact that in a 369-day series this page includes only 5 days of record. The sign of the difference is given at the top of the column. For K-P and R-T the positive sign is to be used for constituents K and R and the negative sign for constituents P and T.

For brevity all the 24 constituent hours for every page of record are not directly represented in the table. The difference for the omitted hours for any page should be taken numerically one greater

than the difference for the given hours on that page. For an example, take the hours for page 2 for constituent OO as derived from constituent J. According to the table the difference for the constituent hours 10 to 3, inclusive, is 9 hours; therefore the difference for the omitted hours 4 to 9, inclusive, should be taken as 10 hours. For constituent 2Q as derived from constituent O the three differences usually required for each page are given in full.

The use of the table may be illustrated from the example above, as follows:

Page 2—

J-hours-----	0,	1,	2,	3,	4,	5,	6,	7,	8,	9,	10,	11
Difference-----	+9,	9,	9,	9,	10,	10,	10,	10,	10,	10,	9,	9
OO-hours-----	9,	10,	11,	12,	14,	15,	16,	17,	18,	19,	19,	20
J-hours-----	12,	13,	14,	15,	16,	17,	18,	19,	20,	21,	22,	23
Difference-----	+9,	9,	9,	9,	9,	9,	9,	9,	9,	9,	9,	9
OO-hours-----	21,	22,	23,	0,	1,	2,	3,	4,	5,	6,	7,	8

The period 24 hours should be added or subtracted when necessary in order that the resulting constituent hours may be between 0 and 23.

Table 34. *For summation of long-period constituents.*—This table is designed to show the assignment of the daily page sums of the hourly heights to the constituent divisions to which they most nearly correspond. The table is based upon formula (395). The constituent division to which each day of series is assigned is given in the left-hand column. For Mf, MSf, and Mm there will frequently occur two consecutive days which are to be assigned to the same constituent division. In such cases the day which most nearly corresponds to the constituent division is the only one given in the table, and this is marked by an asterisk (\*). The missing day, whether it precedes or follows the one marked by the asterisk, is to be assigned to the same constituent division. For Sa a number of consecutive days of series are assigned to each constituent division. In the table there are given the first and last days of each group.

Table 35. *Products aS/15 for Form 444.*—This table contains the products of constituent speeds and time meridian longitudes for formula (466) which is used in obtaining values of  $(\kappa' - \kappa)$  for column A of Form 444.

Table 36. *Angle differences for Form 445.*—This table gives the differences for obtaining and checking the dial settings for February 1 and December 31, as entered in Form 445. The differences are derived from tables 16 and 17.

Table 37. *Coast and Geodetic Survey tide-predicting machine No. 2—General gears.*—This table gives the details of the general gearing from the hand-operating crank to the main vertical shafts, together with the details of the gearing in the front section or dial case. In this table the gears and shafts are each numbered consecutively for convenience of reference, the gears being designated by the letter G and the shafts by the letter S. In the second column are given the face of each bevel or spur gear and the diameter of each shaft. The next two columns contain the number of teeth and pitch of each bevel and spur gear. The pitch is the number of teeth per inch of diameter of the gear. The worm screw is equivalent to a gear of one tooth, as it requires a complete revolution of the screw to move the engaged wheel

one tooth forward. The period of rotation of each shaft and gear is relative and refers to the time as indicated on the face of the machine, which for convenience is called dial time.

Table 38. *Coast and Geodetic Survey tide-predicting machine No. 2—Constituent gears.*—This table contains the details of the gearing from the main vertical shafts to the individual constituent cranks. Column I gives the number of teeth in the bevel gear on the main vertical shaft; column II, the number of teeth in the gear on the intermediate shaft that meshes with the gear on the vertical shaft; column III, the number of teeth in the gear on the intermediate shaft that meshes with the gear on the constituent crank shaft; and column IV, the number of teeth in the gear on the crank shaft.

For the long-period constituents the worm gear is taken as the equivalent of one tooth. For each of these constituents there is a short secondary shaft on which sliding gears are mounted, but the extra gears do not affect the speed of any of the crank shafts except that for constituent Sa in which case a ratio of 1:2 is introduced.

The crank-shaft speed per dial hour for each constituent is equal to  $30^\circ \times \frac{\text{column I}}{\text{column II}} \times \frac{\text{column III}}{\text{column IV}}$ . For constituent Sa the product of both values appearing in each of the columns II and III is to be taken as the value for the column. The column of "Gear speed per dial hour" contains the speeds as computed by the above formula.

For comparison the table contains also the theoretical speed of each of the constituents and the accumulated error per year due to the difference between the theoretical and the gear speeds.

For convenience of reference the table includes also the maximum amplitude settings of the constituent cranks.

Table 39. *Synodic periods of constituents.*—This table is derived from table 28, the period represented by  $360^\circ$  being divided by the speed difference and the results reduced to days.

Table 40. *Day of year corresponding to any date.*—This table is convenient for obtaining the difference between any two dates and also in finding the middle of any series.

Table 41. *Values of  $h$  in formula  $h = (1 + r^2 + 2r \cos x)^{\frac{1}{2}}$ .*—This table may be used with formulas (472) and (473) on page 149 to obtain constituent amplitudes for the prediction of hydraulic currents.

Table 42. *Values of  $k$  in formula  $k = \tan^{-1} \frac{r \sin x}{1 + r \cos x}$ .*—This table may be used with formulas (474) and (475) on pages 149–150 to obtain constituent epochs for the prediction of hydraulic currents.

Table 1.—*Fundamental astronomical data*

Mean distance, earth to sun-----	92, 897, 416 miles <sup>a</sup>
Mean distance, earth to moon-----	238, 857 miles <sup>a</sup>
Equatorial radius of earth (Hayford's Spheroid of 1909)---	3, 963. 34 miles <sup>a</sup>
Polar radius of earth (Hayford's Spheroid of 1909)-----	3, 949. 99 miles <sup>a</sup>
Mean radius of earth ( $a$ ), (Intern. Ell.) 6,371,269 meters <sup>b</sup>	= 20,903,071 feet = 3, 958. 91 miles
Solar parallax (Paris Conference)-----8.80'' <sup>a</sup>	= 0. 000, 042, 66 radian
Lunar equatorial horizontal parallax (Brown) .57' 2.70'' <sup>a</sup>	= 0. 016, 59 radian
Mean solar parallax in respect to mean radius ( $a/c_1$ )-----	0. 000, 042, 61 radian
Mean lunar parallax in respect to mean radius ( $a/c$ )-----	0. 016, 57 radian
Eccentricity of earth's orbit ( $e_1$ ), epoch Jan. 1, 1900-----	0. 016, 75 <sup>c</sup>
Eccentricity of moon's orbit ( $e$ )-----	0. 054, 90 <sup>d</sup>
Obliquity of the ecliptic ( $\omega$ ), epoch Jan. 1, 1900 23° 27' 8.26'' <sup>c</sup> =	23. 452°
Inclination of moon's orbit to plane of ecliptic ( $i$ ) 5° 08' 43.3546'' <sup>d</sup> =	5. 145°
Ratio of mass of sun to combined mass of earth and moon (Sitter)-----	327, 932 <sup>b</sup>
Ratio of mass of earth to mass of moon (Hinks)-----	81. 53 <sup>b</sup>
Mass of sun/mass of earth ( $S/E$ )-----	331, 954
Mass of moon/mass of earth ( $M/E$ )-----	0. 012, 27
Solar coefficient $U_1 = (S/E) (a/c_1)^3$ -----	.2569 x 10 <sup>-7</sup>
Basic factor $U = (M/E) (a/c)^3$ -----	.5582 x 10 <sup>-7</sup>
Solar factor $S' = U_1/U$ -----	0. 4602

In the following formulas for longitude,  $T$  represents the number of Julian centuries (36525 days) reckoned from Greenwich mean noon, December 31, 1899 (Gregorian Calendar).

Mean longitude of sun ( $h$ ) = 279° 41' 48.04'' + 129,602,768.13'' $T$ + 1.089'' $T^2$ <sup>c</sup>
Longitude of solar perigee ( $p_1$ ) = 281° 13' 15.0'' + 6,189.03'' $T$ + 1.63'' $T^2$ + 0.012'' $T^3$ <sup>c</sup>
Mean longitude of moon ( $s$ ) = 270° 26' 14.72'' + (1336 rev. + 1,108,411.20'') $T$ + 9.09'' $T^2$ + 0.006,8'' $T^3$ <sup>c</sup>
Longitude of lunar perigee ( $p$ ) = 334° 19' 40.87'' + (11 rev. + 392,515.94'') $T$ - 37.24'' $T^2$ - 0.045'' $T^3$ <sup>c</sup>
Longitude of moon's node ( $N$ ) = 259° 10' 57.12'' - (5 rev. + 482,912.63'') $T$ + 7.58'' $T^2$ + 0.008'' $T^3$ <sup>c</sup>
Ratio of mean motion of sun to that of moon ( $m$ )-----0. 074, 804

<sup>a</sup> American Ephemeris and Nautical Almanac for year 1940, p. xx.

<sup>b</sup> Table of astronomical constants by W. de Sitter, published in Bulletin of the Astronomical Institutes of the Netherlands, Vol. VIII, No. 307, July 8, 1938, pp. 230-231.

<sup>c</sup> Astronomical Papers for the American Ephemeris, by Simon Newcomb; Vol. VI, pp. 9-10, and Vol. IX, pt. I, p. 224.

<sup>d</sup> The Solar Parallax and Related Constants, by William Harkness, p. 140.

Table 1.—*Fundamental astronomical data—Continued*

## MEAN LONGITUDE OF SOLAR AND LUNAR ELEMENTS FOR CENTURY YEARS

Epoch, Gregorian calendar Greenwich mean civil time	Sun <i>h</i>	Solar perigee <i>p<sub>1</sub></i>	Moon <i>s</i>	Lunar perigee <i>p</i>	Moon's node <i>N</i>
	°	°	°	°	°
1600, Jan. 1, 0 hour	279.857	276.067	99.725	7.417	301.496
1700, Jan. 1, 0 hour	280.624	277.784	47.604	116.501	167.343
1800, Jan. 1, 0 hour	280.407	279.502	342.313	225.453	33.248
1900, Jan. 1, 0 hour	280.190	281.221	277.026	334.384	259.156
2000, Jan. 1, 0 hour	279.973	282.940	211.744	83.294	125.069

## RATE OF CHANGE IN MEAN LONGITUDE OF SOLAR AND LUNAR ELEMENTS (EPOCH, JAN. 1, 1900) ↵

Elements	Per Julian century (36525 days)	Per common year (365 days)	Per solar day	Per solar hour
	°	°	°	°
Sun ( <i>h</i> )	100 <i>r</i> + 0.769	359.761,28	0.985,647,3	0.041,068,64
Solar perigee ( <i>p<sub>1</sub></i> )	1.719	0.017,18	0.000,047,1	0.000,001,96
Moon ( <i>s</i> )	1336 <i>r</i> + 307.892	13 <i>r</i> + 129.384,82	13.176,396.8	0.549,016,53
Lunar perigee ( <i>p</i> )	11 <i>r</i> + 109.032	40.662,47	0.111,404,0	0.004,641,83
Moon's node ( <i>N</i> )	-5 <i>r</i> - 134.142	-19.328,19	-0.052,953,9	-0.002,206,41

## MEAN ASTRONOMICAL PERIODS

*(Symbols refer to rate of change in mean longitude)*

Sidereal day, $360^\circ/(360^\circ+h)$	Solar days
Lunar day, $360^\circ/(360^\circ+h-s)$	0.997,270 1.035,050
Nodical month, $360^\circ/(s-N)$	27.212,220
Tropical month, $360^\circ/s$	27.321,582
Anomalistic month, $360^\circ/(s-p)$	27.554,550
Synodical month, $360^\circ/(s-h)$	29.530,588
Moon's evectional period, $360^\circ/(s-2h+p)$	31.811,939
Eclipse year, $360^\circ/(h-N)$	346.620,0
Tropical year, $360^\circ/h$	365.242,2
Anomalistic year, $360^\circ/(h-p_1)$	365.259,6
Common year	365.000,0
Mean Gregorian year	365.242,5
Mean Julian year	365.250,0
Leap year	366.000,0
Evective period in moon's parallax, $360^\circ/(h-p)$	411.784,7
Revolution of lunar perigee, $360^\circ/p$	8.85 Julian years
Revolution of moon's node, $360^\circ/N$	18.61 Julian years
Revolution of solar perigee, $360^\circ/p_1$	209 Julian centuries

Table 2.—Harmonic constituents

Ref. No.	Symbol	Argument (E)		Speed per solar hour	Coeffi- cient (C)	Factor-f formula
		V	u			
LUNAR LONG-PERIOD TERMS, FORMULA (62) .						
A <sub>1</sub> ...		Zero (permanent term) . . . . .	zero . . . . .	zero . . . . .	0. 5044	(73)
A <sub>2</sub> ...	Mm	s-p . . . . .	zero . . . . .	0. 544, 374, 7	0. 0827	(73)
A <sub>3</sub> ...		2s-2p . . . . .	zero . . . . .	1. 088, 749, 4	0. 0068	(73)
A <sub>4</sub> ...		s-2h+p . . . . .	zero . . . . .	0. 471, 521, 1	0. 0116	(73)
A <sub>5</sub> ...	MSf	2s-2h . . . . .	zero . . . . .	1. 015, 895, 8	0. 0084	(73)
A <sub>6</sub> ...	Mf	2s . . . . .	-2ξ . . . . .	1. 098, 033, 1	0. 1566	(74)
A <sub>7</sub> ...		3s-p . . . . .	-2ξ . . . . .	1. 642, 407, 8	0. 0303	(74)
A <sub>8</sub> ...		s+p+180° . . . . .	-2ξ . . . . .	0. 553, 658, 4	0. 0043	(74)
A <sub>9</sub> ...		4s-2p . . . . .	-2ξ . . . . .	2. 186, 782, 5	0. 0040	(74)
A <sub>10</sub> ...		3s-2h+p . . . . .	-2ξ . . . . .	1. 569, 554, 3	0. 0043	(74)
A <sub>11</sub> ...		s+2h-p+180° . . . . .	-2ξ . . . . .	0. 626, 512, 0	0. 0006	(74)
A <sub>12</sub> ...		4s-2h . . . . .	-2ξ . . . . .	2. 113, 928, 8	0. 0025	(74)
A <sub>13</sub> ...		2h . . . . .	-2ξ . . . . .	0. 082, 137, 3	0. 0001	(74)
LUNAR LONG-PERIOD TERMS, FORMULA (137)						
A <sub>41</sub> ...		s-90° . . . . .	-ξ . . . . .	0. 549, 016, 5	0. 0399	(141)
A <sub>45</sub> ...		2s-p-90° . . . . .	-ξ . . . . .	1. 093, 391, 2	0. 0065	(141)
A <sub>46</sub> ...		p-90° . . . . .	-ξ . . . . .	0. 004, 641, 8	0. 0022	(141)
A <sub>47</sub> ...		3s-90° . . . . .	-3ξ . . . . .	1. 647, 049, 6	0. 0032	(142)
A <sub>48</sub> ...		4s-p-90° . . . . .	-3ξ . . . . .	2. 191, 424, 3	0. 0009	(142)
SOLAR LONG-PERIOD TERMS, FORMULA (185) .						
B <sub>1</sub> ...		Zero (permanent term) . . . . .	zero . . . . .	zero . . . . .	0. 2340	unity
B <sub>2</sub> ...		h-p <sub>1</sub> . . . . .	zero . . . . .	0. 041, 066, 7	0. 0118	unity
B <sub>3</sub> ...		2h-2p <sub>1</sub> . . . . .	zero . . . . .	0. 082, 133, 4	0. 0003	unity
B <sub>4</sub> ...	Ssa	2h . . . . .	zero . . . . .	0. 082, 137, 3	0. 0728	unity
B <sub>5</sub> ...		3h-p <sub>1</sub> . . . . .	zero . . . . .	0. 123, 204, 0	0. 0043	unity
B <sub>6</sub> ...		h+p <sub>1</sub> +180° . . . . .	zero . . . . .	0. 041, 070, 6	0. 0006	unity
B <sub>7</sub> ...		4h-2p <sub>1</sub> . . . . .	zero . . . . .	0. 164, 270, 6	0. 0002	unity
SOLAR LONG-PERIOD TERM, PARAGRAPH 119						
B <sub>84</sub> ...	Sa	h . . . . .	zero . . . . .	0. 041, 068, 6		unity
LUNAR DIURNAL TERMS, FORMULA (63)						
A <sub>14</sub> ...	O <sub>1</sub> ...	T-2s+h+90° . . . . .	+2ξ-v . . . . .	13, 943, 035, 6	0. 3771	(75)
A <sub>15</sub> ...	Q <sub>1</sub> ...	T-3s+h+p+90° . . . . .	+2ξ-v . . . . .	13, 398, 660, 9	0. 0730	(75)
A <sub>16</sub> ...	(M <sub>1</sub> )...	T-s+h-p-90° . . . . .	+2ξ-v . . . . .	14, 487, 410, 3	0. 0104	(75)
A <sub>17</sub> ...	2Q <sub>1</sub> ...	T-4s+h+2p+90° . . . . .	+2ξ-v . . . . .	12, 854, 286, 2	0. 0097	(75)
A <sub>18</sub> ...	ρ <sub>1</sub> ...	T-3s+3h-p+90° . . . . .	+2ξ-v . . . . .	13, 471, 514, 5	0. 0142	(75)
A <sub>19</sub> ...		T-s-h+p-90° . . . . .	+2ξ-v . . . . .	14, 414, 556, 7	0. 0015	(75)
A <sub>20</sub> ...	σ <sub>1</sub> ...	T-4s+3h+90° . . . . .	+2ξ-v . . . . .	12, 927, 139, 8	0. 0061	(75)
A <sub>21</sub> ...		T-h+90° . . . . .	+2ξ-v . . . . .	14, 958, 931, 4	0. 0003	(75)
A <sub>22</sub> ...	(K <sub>1</sub> )...	T+h-90° . . . . .	-v . . . . .	15, 041, 068, 6	0. 3623	(76)
A <sub>23</sub> ...	(M <sub>1</sub> )...	T-s+h+p-90° . . . . .	-v . . . . .	14, 496, 693, 9	0. 0297	(76)
A <sub>24</sub> ...	J <sub>1</sub> ...	T+s-h-p-90° . . . . .	-v . . . . .	15, 585, 443, 3	0. 0297	(76)
A <sub>25</sub> ...		T-2s+h+2p-90° . . . . .	-v . . . . .	13, 952, 319, 2	0. 0024	(76)
A <sub>26</sub> ...		T+2s+h-2p-90° . . . . .	-v . . . . .	16, 129, 818, 0	0. 0024	(76)
A <sub>27</sub> ...	X <sub>1</sub> ...	T-s+3h-p-90° . . . . .	-v . . . . .	14, 569, 547, 6	0. 0042	(76)
A <sub>28</sub> ...	θ <sub>1</sub> ...	T+s-h+p-90° . . . . .	-v . . . . .	15, 512, 589, 7	0. 0042	(76)
LUNAR DIURNAL TERMS, FORMULA (63)						
A <sub>29</sub> ...	MP <sub>1</sub> ...	T-2s+3h-90° . . . . .	-v . . . . .	14, 025, 172, 9	0. 0030	(76)
A <sub>30</sub> ...	SO <sub>1</sub> ...	T+2s-h-90° . . . . .	-v . . . . .	16, 056, 964, 4	0. 0030	(76)
A <sub>31</sub> ...	OO <sub>1</sub> ...	T+2s+h-90° . . . . .	-2ξ-v . . . . .	16, 139, 101, 7	0. 0163	(77)
A <sub>32</sub> ...	KQ <sub>1</sub> ...	T+3s+h-p-90° . . . . .	-2ξ-v . . . . .	16, 683, 476, 4	0. 0032	(77)
A <sub>33</sub> ...		T+s+h+p+90° . . . . .	-2ξ-v . . . . .	15, 594, 727, 0	0. 0005	(77)
A <sub>34</sub> ...		T+4s+h-2p-90° . . . . .	-2ξ-v . . . . .	17, 227, 851, 1	0. 0004	(77)
A <sub>35</sub> ...		T+3s-h+p-90° . . . . .	-2ξ-v . . . . .	16, 610, 622, 8	0. 0004	(77)
A <sub>36</sub> ...		T+s+3h-p+90° . . . . .	-2ξ-v . . . . .	15, 067, 580, 6	0. 0001	(77)
A <sub>37</sub> ...		T+4s-h-90° . . . . .	-2ξ-v . . . . .	17, 154, 997, 5	0. 0003	(77)
A <sub>38</sub> ...		T+3h-90° . . . . .	-2ξ-v . . . . .	15, 123, 205, 9		(77)

Table 2.—Harmonic constituents—Continued

Ref. No.	Symbol	Argument (E)		Speed per solar hour	Coeffi- cient (C)	Factor-f formula
		V	u			
LUNAR DIURNAL TERMS, FORMULA (138)						
A <sub>69</sub>		T-3s+h	+3ξ-v	13.394,019,0	0.0116	(143)
A <sub>70</sub>		T-4s+h+p	+3ξ-v	12.849,644,4	0.0032	(143)
A <sub>71</sub>	(M <sub>1</sub> )	T-s+h	+ξ-v	14.492,052,1	0.0367	(144)
A <sub>72</sub>		T-2s+h+p	+ξ-v	13.947,677,4	0.0060	(144)
A <sub>73</sub>		T+h-p	+ξ-v	15.036,426,8	0.0020	(144)
A <sub>74</sub>		T+s+h	-ξ-v	15.590,085,2	0.0134	(145)
A <sub>75</sub>		T+2s+h-p	-ξ-v	16.134,459,9	0.0022	(145)
SOLAR DIURNAL TERMS, FORMULA (186)						
B <sub>14</sub>	P <sub>1</sub>	T-h+90°	zero	14.958,931,4	0.1755	unity
B <sub>15</sub>	π <sub>1</sub>	T-2h+p <sub>1</sub> +90°	zero	14.917,864,7	0.0103	unity
B <sub>16</sub>		T-p <sub>1</sub> -90°	zero	14.999,998,0	0.0015	unity
B <sub>17</sub>		T-3h+2p <sub>1</sub> +90°	zero	14.876,798,0	0.0004	unity
B <sub>22</sub>	(K <sub>1</sub> )	T+h-90°	zero	15.041,068,6	0.1681	unity
B <sub>23</sub>		T+p <sub>1</sub> -90°	zero	15.000,002,0	0.0042	unity
B <sub>24</sub>	ψ <sub>1</sub>	T+2h-p <sub>1</sub> -90°	zero	15.082,135,3	0.0042	unity
B <sub>25</sub>		T-h+2p <sub>1</sub> -90°	zero	14.958,935,4	0.0001	unity
B <sub>26</sub>		T+3h-2p <sub>1</sub> -90°	zero	15.123,202,0	0.0001	unity
B <sub>31</sub>	φ <sub>1</sub>	T+3h-90°	zero	15.123,205,9	0.0076	unity
B <sub>32</sub>		T+4h-p <sub>1</sub> -90°	zero	15.164,272,6	0.0004	unity
B <sub>33</sub>		T+2h+p <sub>1</sub> +90°	zero	15.082,139,2	0.0001	unity
B <sub>34</sub>		T+5h-2p <sub>1</sub> -90°	zero	15.205,339,3		unity
SOLAR DIURNAL TERM, PARAGRAPH 119						
B <sub>71</sub>	S <sub>1</sub>	T	zero	15.000,000,0		unity
COMBINATION DIURNAL TERMS, FORMULAS (194), (201), AND (222)						
Note 1	M <sub>1</sub>	T-s+h+p-90°	-v-Q <sub>u</sub>	14.496,693,9	0.0209*	(206)
Note 2	K <sub>1</sub>	T-s+h-90°	+ξ-v+Q		0.5305	(227)
		T+h-90°	-v'	15.041,068,6		
LUNAR SEMIDIURNAL TERMS, FORMULA (64)						
A <sub>39</sub>	M <sub>2</sub>	(2T-2s+2h)	+2ξ-2v	28.984,104,2	0.9085	(78)
A <sub>40</sub>	N <sub>2</sub>	2T-3s+2h+p	+2ξ-2v	28.439,729,5	0.1759	(78)
A <sub>41</sub>	(L <sub>2</sub> )	2T-s+2h-p+180°	+2ξ-2v	29.528,478,9	0.0251	(78)
A <sub>42</sub>	2N <sub>2</sub>	2T-4s+2h+2p	+2ξ-2v	27.895,354,8	0.0235	(78)
A <sub>43</sub>	v <sub>2</sub>	2T-s+4h-p	+2ξ-2v	28.512,583,1	0.0341	(78)
A <sub>44</sub>	λ <sub>2</sub>	2T-s+p+180°	+2ξ-2v	29.455,625,3	0.0066	(78)
A <sub>45</sub>	μ <sub>2</sub>	2T-4s+4h	+2ξ-2v	27.968,208,4	0.0219	(78)
A <sub>46</sub>	(S <sub>2</sub> )	2T	+2ξ-2v	30.000,000,0	0.0006	(78)
A <sub>47</sub>	(K <sub>2</sub> )	2T+2h	-2v	30.082,137,3	0.0786	(79)
A <sub>48</sub>	(L <sub>2</sub> )	2T-s+2h+p	-2v	29.537,762,6	0.0064	(79)
A <sub>49</sub>	KJ <sub>2</sub>	2T+s+2h-p	-2v	30.626,512,0	0.0064	(79)
A <sub>50</sub>		2T-s+2h+2p	-2v	28.993,387,9	0.0005	(79)
A <sub>51</sub>		2T+2s+2h-2p	-2v	31.170,886,7	0.0005	(79)
A <sub>52</sub>		2T-s+4h-p	-2v	29.610,616,2	0.0009	(79)
A <sub>53</sub>		2T+s+p	-2v	30.553,658,4	0.0009	(79)
A <sub>54</sub>		2T-2s+4h	-2v	29.066,241,5	0.0007	(79)
A <sub>55</sub>		2T+2s	-2v	31.098,033,1	0.0007	(79)
A <sub>56</sub>		2T+2s+2h	-2ξ-2v	31.180,170,3	0.0017	(80)
A <sub>57</sub>		2T+3s+2h-p	-2ξ-2v	31.724,545,0	0.0003	(80)
A <sub>58</sub>		2T+s+2h+p+180°	-2ξ-2v	30.655,795,6		(80)
A <sub>59</sub>		2T+4s+2h-2p	-2ξ-2v	32.268,919,7		(80)
A <sub>60</sub>		2T+3s+p	-2ξ-2v	31.651,691,4		(80)
A <sub>61</sub>		2T+s+4h-p+180°	-2ξ-2v	30.708,649,3		(80)
A <sub>62</sub>		2T+4s	-2ξ-2v	32.196,066,1		(80)
A <sub>63</sub>		2T+4h	-2ξ-2v	30.164,274,6		(80)

\*Adapted for use with tabular node factors, theoretical value is 0.0317. See p. 43.

Table 2.—*Harmonic constituents—Continued*

Ref. No.	Symbol	Argument (E)		Speed per solar hour	Coeffi- cient (C)	Factor-f formula
		V	u			
LUNAR SEMIDIURNAL TERMS, FORMULA (139)						
$A_{16}$		$2T - 3s + 2h + 90^\circ$	$+3\xi - 2\nu$	28.435, 087, 7	0.0223	(146)
$A_{77}$		$2T - 4s + 2h + p + 90^\circ$	$+3\xi - 2\nu$	27.890, 713, 0	0.0062	(146)
$A_{78}$		$2T - 2s + 2h - p - 90^\circ$	$+3\xi - 2\nu$	28.979, 462, 4	0.0012	(146)
$A_{79}$		$2T - s + 2h - 90^\circ$	$+\xi - 2\nu$	29.533, 120, 8	0.0209	(147)
$A_{80}$		$2T - 2s + 2h + p - 90^\circ$	$+\xi - 2\nu$	28.988, 746, 0	0.0034	(147)
$A_{81}$		$2T + s + 2h - 90^\circ$	$-\xi - 2\nu$	30.631, 153, 8	0.0019	(148)
SOLAR SEMIDIURNAL TERMS, FORMULA (187)						
$B_{39}$	$S_2$	$2T$	zero	30.000, 000, 0	0.4227	unity
$B_{40}$	$T_2$	$2T - h + p_1$	zero	29.958, 933, 3	0.0248	unity
$B_{41}$	$R_2$	$2T + h - p_1 + 180^\circ$	zero	30.041, 066, 7	0.0035	unity
$B_{42}$		$2T - 2h + 2p_1$	zero	29.917, 866, 6	0.0010	unity
$B_{47}$	$(K_2)$	$2T + 2h$	zero	30.082, 137, 3	0.0365	unity
$B_{48}$		$2T + h + p_1$	zero	30.041, 070, 6	0.0009	unity
$B_{49}$		$2T + 3h - p_1$	zero	30.123, 204, 0	0.0009	unity
$B_{50}$		$2T + 2p_1$	zero	30.000, 003, 9	-----	unity
$B_{51}$		$2T + 4h - 2p_1$	zero	30.164, 270, 6	-----	unity
$B_{66}$		$2T + 4h$	zero	30.164, 274, 6	0.0008	unity
$B_{67}$		$2T + 5h - p_1$	zero	30.205, 341, 2	-----	unity
$B_{68}$		$2T + 3h + p_1 + 180^\circ$	zero	30.123, 207, 9	-----	unity
$B_{59}$		$2T + 6h - 2p_1$	zero	30.246, 407, 9	-----	unity
COMBINATION SEMIDIURNAL TERMS, FORMULAS (212) AND (230)						
Note 3.	$L_2$	$2T - s + 2h - p + 180^\circ$	$+2\xi - 2\nu - R$	29.528, 473, 9	0.0251	(215)
Note 4.	$K_2$	$2T + 2h$	$-2\nu''$	30.082, 137, 3	0.1151	(235)
LUNAR TERDIURNAL TERMS, FORMULA (140)						
$A_{82}$	$M_3$	$3T - 3s + 3h$	$+3\xi - 3\nu$	43.476, 156, 3	0.0178	(149)
$A_{83}$		$3T - 4s + 3h + p$	$+3\xi - 3\nu$	42.931, 781, 6	0.0050	(149)
$A_{84}$		$3T - 2s + 3h - p + 180^\circ$	$+3\xi - 3\nu$	44.020, 531, 0	0.0010	(149)
$A_{85}$		$3T - 5s + 3h + 2p$	$+3\xi - 3\nu$	42.387, 406, 9	0.0009	(149)
$A_{86}$		$3T - 4s + 5h - p$	$+3\xi - 3\nu$	43.004, 635, 2	0.0007	(149)
$A_{87}$		$3T - s + 3h$	$+\xi - 3\nu$	44.574, 189, 4	0.0024	(150)
$A_{88}$		$3T - 2s + 3h + p$	$+\xi - 3\nu$	44.029, 814, 7	0.0004	(150)
SOLAR TERDIURNAL TERM, PARAGRAPH 119						
$B_{82}$	$S_3$	$3T$	zero	45.000, 000, 0	-----	unity

Note 1—Combines terms  $A_{16}$  and  $A_{23}$ .Note 2—Combines terms  $A_{22}$  and  $B_{22}$ .Note 3—Combines terms  $A_{41}$  and  $A_{48}$ .Note 4—Combines terms  $A_{47}$  and  $B_{47}$ .

Table 2a.—*Shallow-water constituents*

Symbol	Argument			Speed	Factor- $f$
	Origin	$V$	$u$		
<i>Semidiurnal</i>					
MNS <sub>2</sub> .....	M <sub>2</sub> +N <sub>2</sub> -S <sub>2</sub> .....	2T-5s+4h+p.....	+4ξ-4ν.....	27.423, 833, 7	$f^2(M_2)$
2SM <sub>2</sub> .....	2S <sub>2</sub> -M <sub>2</sub> .....	2T+2s-2h.....	-2ξ+2ν.....	31.015, 895, 8	$f(M_2)$
<i>Tertiurnal</i>					
MK <sub>3</sub> .....	M <sub>2</sub> +K <sub>1</sub> .....	3T-2s+3h-90°.....	+2ξ-2ν-ν'.....	44.025, 172, 9	$f(M_2) \times f(K_1)$
2MK <sub>3</sub> .....	2M <sub>2</sub> -K <sub>1</sub> .....	3T-4s+3h+90°.....	+4ξ-4ν+ν'.....	42.927, 139, 8	$f^2(M_2) \times f(K_1)$
SK <sub>3</sub> .....	S <sub>2</sub> +K <sub>1</sub> .....	3T+h-90°.....	-ν'.....	45.041, 068, 6	$f(K_1)$
SO <sub>3</sub> .....	S <sub>2</sub> +O <sub>1</sub> .....	3T-2s+h+90°.....	+2ξ-ν.....	43.943, 035, 6	$f(O_1)$
<i>Quarter diurnal</i>					
M <sub>4</sub> .....	2M <sub>2</sub> .....	4T-4s+4h.....	+4ξ-4ν.....	57.968, 208, 4	$f^2(M_2)$
MS <sub>4</sub> .....	M <sub>2</sub> +S <sub>2</sub> .....	4T-2s+2h.....	+2ξ-2ν.....	58.984, 104, 2	$f^2(M_2)$
MN <sub>4</sub> .....	M <sub>2</sub> +N <sub>2</sub> .....	4T-5s+4h+p.....	+4ξ-4ν.....	57.423, 833, 7	$f^2(M_2)$
MK <sub>4</sub> .....	M <sub>2</sub> +K <sub>2</sub> .....	4T-2s+4h.....	+2ξ-2ν-2ν''.....	59.066, 241, 5	$f(M_2) \times f(K_2)$
S <sub>4</sub> .....	2S <sub>2</sub> .....	4T.....	zero.....	60.000, 000, 0	unity
<i>Sixth diurnal</i>					
M <sub>6</sub> .....	3M <sub>2</sub> .....	6T-6s+6h.....	+6ξ-6ν.....	86.952, 312, 7	$f^3(M_2)$
2MS <sub>6</sub> .....	2M <sub>2</sub> +S <sub>2</sub> .....	6T-4s+4h.....	+4ξ-4ν.....	87.968, 208, 4	$f^3(M_2)$
2MN <sub>6</sub> .....	2M <sub>2</sub> +N <sub>2</sub> .....	6T-7s+6h+p.....	+6ξ-6ν.....	86.407, 938, 0	$f^3(M_2)$
2SM <sub>6</sub> .....	2S <sub>2</sub> +M <sub>2</sub> .....	6T-2s+2h.....	+2ξ-2ν.....	88.984, 104, 2	$f^3(M_2)$
MSN <sub>6</sub> .....	M <sub>2</sub> +S <sub>2</sub> +N <sub>2</sub> .....	6T-6s+4h+p.....	+4ξ-4ν.....	87.423, 833, 7	$f^3(M_2)$
S <sub>6</sub> .....	3S <sub>2</sub> .....	6T.....	zero.....	90.000, 000, 0	unity
<i>Eighth diurnal</i>					
M <sub>8</sub> .....	4M <sub>2</sub> .....	8T-8s+8h.....	+8ξ-8ν.....	115.936, 416, 9	$f^4(M_2)$
3MS <sub>8</sub> .....	3M <sub>2</sub> +S <sub>2</sub> .....	8T-6s+6h.....	+6ξ-6ν.....	116.952, 312, 7	$f^3(M_2)$
2(MS) <sub>8</sub> .....	2M <sub>2</sub> +2S <sub>2</sub> .....	8T-4s+4h.....	+4ξ-4ν.....	117.968, 208, 4	$f^2(M_2)$
2MSN <sub>8</sub> .....	2M <sub>2</sub> +S <sub>2</sub> +N <sub>2</sub> .....	8T-7s+6h+p.....	+6ξ-6ν.....	116.407, 938, 0	$f^3(M_2)$
S <sub>8</sub> .....	4S <sub>2</sub> .....	8T.....	zero.....	120.000, 000, 0	unity

Table 3.—Latitude factors

Y	Y <sub>v30</sub>	Y <sub>v31</sub> Y <sub>s30</sub> Y <sub>s32</sub> Y <sub>w42</sub>	Y <sub>v32</sub> Y <sub>w43</sub>	Y <sub>s31</sub>	Y <sub>w31</sub>	Y <sub>w32</sub>	Y <sub>v40</sub>	Y <sub>v41</sub> Y <sub>s40</sub>	Y <sub>v42</sub> Y <sub>s43</sub>	Y <sub>v43</sub>	Y <sub>s41</sub>	Y <sub>s42</sub>	Y <sub>w41</sub>	Y
0	0.500	0.000	1.000	1.000	0.000	1.000	0.000	0.200	0.000	1.000	0.000	0.333	0.200	0
1	.500	.035	1.000	0.999	.017	1.000	.010	.200	.017	1.000	.013	.333	.200	1
2	.498	.070	0.999	.998	.035	0.999	.021	.199	.035	0.998	.026	.332	.199	2
3	.496	.105	.997	.995	.052	.999	.031	.197	.052	.996	.038	.330	.197	3
4	.493	.139	.995	.990	.070	.998	.041	.195	.069	.993	.051	.328	.195	4
5	.489	.174	.992	.985	.087	.996	.052	.192	.086	.989	.063	.324	.192	5
6	.484	.208	.989	.978	.105	.995	.062	.188	.103	.984	.076	.321	.189	6
7	.478	.242	.985	.970	.122	.993	.071	.184	.120	.978	.088	.316	.185	7
8	.471	.276	.981	.961	.139	.990	.081	.179	.136	.971	.099	.311	.181	8
9	.463	.309	.976	.951	.156	.988	.090	.173	.153	.964	.111	.305	.176	9
10	.455	.342	.970	.940	.174	.985	.099	.167	.168	.955	.122	.299	.170	10
11	.445	.375	.964	.927	.191	.982	.108	.161	.184	.946	.133	.291	.164	11
12	.435	.407	.957	.914	.208	.978	.116	.153	.199	.936	.143	.284	.157	12
13	.424	.438	.949	.899	.225	.974	.124	.146	.214	.925	.154	.275	.149	13
14	.412	.469	.941	.883	.242	.970	.131	.137	.228	.914	.163	.267	.141	14
15	.400	.500	.933	.866	.259	.966	.138	.128	.241	.901	.172	.257	.133	15
16	.386	.530	.924	.848	.276	.961	.144	.119	.255	.888	.181	.247	.124	16
17	.372	.559	.915	.829	.292	.956	.150	.110	.267	.875	.189	.237	.115	17
18	.357	.588	.905	.809	.309	.951	.156	.099	.280	.860	.197	.226	.105	18
19	.341	.616	.894	.788	.326	.946	.161	.089	.291	.845	.204	.215	.094	19
20	.325	.643	.883	.766	.342	.940	.165	.078	.302	.830	.211	.203	.083	20
21	.307	.669	.872	.743	.358	.934	.169	.067	.312	.814	.217	.191	.072	21
22	.290	.695	.860	.719	.375	.927	.172	.055	.322	.797	.222	.179	.060	22
23	.271	.719	.847	.695	.391	.921	.175	.044	.331	.780	.227	.166	.047	23
24	.252	.743	.835	.669	.407	.914	.177	.032	.339	.762	.231	.153	.035	24
25	.232	.766	.821	.643	.423	.906	.178	.019	.347	.744	.234	.140	.021	25
26	.212	.788	.808	.616	.438	.899	.179	.007	.354	.726	.237	.127	.008	26
27	.191	.809	.794	.588	.454	.891	.179	-.005	.360	.707	.239	.113	-.006	27
28	.169	.829	.780	.559	.469	.883	.178	-.018	.366	.688	.241	.100	-.020	28
29	.147	.848	.765	.530	.485	.875	.177	-.031	.371	.671	.242	.086	-.035	29
30	.125	.866	.750	.500	.500	.866	.175	-.043	.375	.650	.242	.072	-.050	30
31	.102	.883	.735	.469	.515	.857	.172	-.056	.378	.630	.241	.058	-.065	31
32	.079	.899	.719	.438	.530	.848	.169	-.069	.381	.610	.240	.045	-.081	32
33	.055	.914	.703	.407	.545	.839	.165	-.081	.383	.590	.238	.031	-.097	33
34	.031	.927	.687	.375	.559	.829	.161	-.093	.384	.570	.235	.071	-.113	34
35	.007	.940	.671	.342	.574	.819	.155	-.106	.385	.550	.232	.004	-.129	35
36	-.018	.951	.655	.309	.588	.809	.150	-.118	.385	.530	.228	-.010	-.145	36
37	-.043	.961	.638	.276	.602	.799	.143	-.130	.384	.509	.223	-.023	-.162	37
38	-.069	.970	.621	.242	.616	.788	.136	-.141	.382	.489	.218	-.036	-.179	38
39	-.094	.978	.604	.208	.629	.777	.128	-.152	.380	.469	.212	-.049	-.196	39
40	-.120	.985	.587	.174	.643	.766	.120	-.163	.377	.450	.206	-.061	-.213	40
41	-.146	.990	.570	.139	.656	.755	.111	-.174	.374	.430	.199	-.073	-.230	41
42	-.172	.995	.552	.105	.669	.743	.102	-.184	.370	.410	.191	-.085	-.248	42
43	-.198	.998	.535	.070	.682	.731	.092	-.194	.365	.391	.183	-.096	-.265	43
44	-.224	.999	.517	.035	.695	.719	.082	-.203	.359	.372	.174	-.107	-.283	44
45	-.250	1.000	.500	.000	.707	.707	.071	-.212	.354	.354	.165	-.118	-.300	45

\*In these columns reverse signs for south latitude. Other values are applicable to either north or south latitude.

Table 3.—Latitude factors—Continued

Y	Y <sub>v30</sub>	Y <sub>v31</sub> Y <sub>e30</sub>	Y <sub>v32</sub> Y <sub>e32</sub>	Y <sub>w43</sub> Y <sub>w42</sub>	Y <sub>v31</sub>	Y <sub>w31</sub>	Y <sub>w32</sub>	Y <sub>v40</sub>	Y <sub>v41</sub> Y <sub>s40</sub>	Y <sub>v42</sub> Y <sub>e43</sub>	Y <sub>v43</sub>	Y <sub>s41</sub>	Y <sub>s42</sub>	Y <sub>w41</sub>	Y
45	-0.250	1.000	0.500	0.000	0.707	0.707	0.071	-0.212	0.354	0.354	0.165	-0.118	-0.300	45	o
46	-0.276	0.999	.483	-.035	.719	.695	.059	-.221	.347	.335	.155	-.128	-.317	46	
47	-0.302	0.998	.465	-.070	.731	.682	.048	-.228	.340	.317	.145	-.137	-.335	47	
48	-0.328	0.995	.448	-.105	.743	.669	.035	-.236	.333	.300	.135	-.146	-.352	48	
49	-0.354	0.990	.430	-.139	.755	.656	.023	-.242	.325	.282	.124	-.155	-.370	49	
50	-0.380	0.985	.413	-.174	.766	.643	.010	-.249	.317	.266	.112	-.163	-.387	40	
51	-0.406	0.978	.396	-.208	.777	.629	-.003	-.254	.308	.249	.101	-.170	-.404	51	
52	-0.431	0.970	.379	-.242	.788	.616	-.017	-.259	.299	.233	.089	-.177	-.421	52	
53	-0.457	0.961	.362	-.276	.799	.602	-.030	-.263	.289	.218	.076	-.183	-.438	53	
54	-0.482	0.951	.345	-.309	.809	.588	-.044	-.267	.280	.203	.064	-.189	-.455	54	
55	-0.507	0.940	.329	-.342	.819	.574	-.058	-.270	.269	.189	.051	-.194	-.471	55	
56	-0.531	0.927	.313	-.375	.829	.559	-.072	-.272	.259	.175	.038	-.198	-.487	56	
57	-0.555	0.914	.297	-.407	.839	.545	-.087	-.274	.249	.162	.025	-.202	-.503	57	
58	-0.579	.899	.281	-.438	.848	.530	-.101	-.275	.238	.149	.012	-.204	-.519	58	
59	-0.602	.883	.265	-.469	.857	.515	-.115	-.275	.227	.137	-.001	-.207	-.535	59	
60	-0.625	.866	.250	-.500	.866	.500	-.130	-.275	.217	.125	-.014	-.208	-.550	60	
61	-0.647	.848	.235	-.530	.875	.485	-.144	-.274	.206	.114	-.028	-.209	-.565	61	
62	-0.669	.829	.220	-.559	.883	.469	-.158	-.272	.195	.103	-.041	-.210	-.580	62	
63	-0.691	.809	.206	-.588	.891	.454	-.173	-.270	.184	.094	-.054	-.209	-.594	63	
64	-0.712	.788	.192	-.616	.899	.438	-.187	-.266	.173	.084	-.067	-.208	-.608	64	
65	-0.732	.766	.179	-.643	.906	.423	-.201	-.263	.162	.075	-.080	-.206	-.621	65	
66	-0.752	.743	.165	-.669	.914	.407	-.214	-.258	.151	.067	-.092	-.204	-.635	66	
67	-0.771	.719	.153	-.695	.921	.391	-.228	-.253	.141	.060	-.105	-.201	-.647	67	
68	-0.790	.695	.140	-.719	.927	.375	-.241	-.247	.130	.053	-.117	-.197	-.660	68	
69	-0.807	.669	.128	-.743	.934	.358	-.254	-.241	.120	.046	-.129	-.193	-.672	69	
70	-0.825	.643	.117	-.766	.940	.342	-.266	-.234	.110	.040	-.141	-.188	-.683	70	
71	-0.841	.616	.106	-.788	.946	.326	-.278	-.226	.100	.035	-.152	-.183	-.694	71	
72	-0.857	.588	.095	-.809	.951	.309	-.290	-.218	.091	.030	-.163	-.177	-.705	72	
73	-0.872	.559	.085	-.829	.956	.292	-.301	-.209	.082	.025	-.173	-.170	-.715	73	
74	-0.886	.530	.076	-.848	.961	.276	-.311	-.200	.073	.021	-.183	-.163	-.724	74	
75	-0.900	.500	.067	-.866	.966	.259	-.322	-.190	.065	.017	-.193	-.155	-.733	75	
76	-0.912	.469	.059	-.883	.970	.242	-.331	-.179	.057	.014	-.202	-.147	-.741	76	
77	-0.924	.438	.051	-.899	.974	.225	-.340	-.169	.049	.011	-.211	-.139	-.749	77	
78	-0.935	.407	.043	-.914	.978	.208	-.349	-.159	.042	.009	-.219	-.130	-.757	78	
79	-0.945	.375	.036	-.927	.982	.191	-.357	-.146	.036	.007	-.226	-.120	-.764	79	
80	-0.955	.342	.030	-.940	.985	.174	-.364	-.134	.030	.005	-.233	-.111	-.770	80	
81	-0.963	.309	.024	-.951	.988	.156	-.371	-.121	.024	.004	-.239	-.100	-.776	81	
82	-0.971	.276	.019	-.961	.990	.139	-.377	-.109	.019	.003	-.245	-.090	-.781	82	
83	-0.978	.242	.015	-.970	.993	.122	-.382	-.096	.015	.002	-.250	-.079	-.785	83	
84	-0.984	.208	.011	-.978	.995	.105	-.387	-.082	.011	.001	-.254	-.069	-.789	84	
85	-0.989	.174	.008	-.985	.996	.087	-.391	-.069	.008	.001	-.258	-.057	-.792	85	
86	-0.993	.139	.005	-.990	.998	.070	-.394	-.055	.005	.000	-.261	-.046	-.795	86	
87	-0.996	.105	.003	-.995	.999	.052	-.397	-.042	.003	.000	-.264	-.035	-.797	87	
88	-0.998	.070	.001	-.998	.999	.035	-.399	-.028	.001	.000	-.265	-.023	-.799	88	
89	-1.000	.035	.000	-.999	1.000	.017	-.400	-.014	.000	.000	-.266	-.012	-.800	89	
90	-1.000	.000	.000	-1.000	1.000	.000	-.400	-.000	.000	.000	-.267	-.000	-.800	90	

\*In these columns reverse signs for south latitude. Other values are applicable to either north or south latitude.

Table 4.—*Mean longitude of lunar and solar elements at Jan. 1, 0 hour, Greenwich mean civil time, of each year from 1800 to 2000*[s=mean longitude of moon; p=mean longitude lunar perigee; h=mean longitude of sun; p<sub>1</sub>=mean longitude solar perigee; N=longitude of moon's node]

Year.	s	p	h	p <sub>1</sub>	N	Year.	s	p	h	p <sub>1</sub>	N
	o	o	o	o	o		o	o	o	o	o
1800	342.31	225.45	280.41	279.50	33.25	1852	28.44	181.24	279.82	280.40	107.55
1801	111.70	266.12	280.17	279.52	13.92	1853	171.00	222.02	280.57	280.41	88.16
1802	241.08	306.78	279.93	279.54	354.59	1854	330.38	262.68	280.33	280.43	68.54
1803	10.47	347.44	279.69	279.55	335.26	1855	69.77	303.34	280.09	280.45	49.51
1804	139.85	28.10	279.45	279.57	315.93	1856	199.15	344.00	279.85	280.46	30.18
1805	282.41	68.88	280.20	279.59	296.55	1857	341.72	24.78	280.60	280.48	10.80
1806	51.80	109.54	279.96	279.61	277.23	1858	111.10	65.44	280.36	280.50	351.47
1807	181.18	150.20	279.72	279.62	257.90	1859	240.49	106.10	280.12	280.52	332.14
1808	310.57	190.86	279.48	279.64	238.57	1860	9.87	146.77	279.88	280.53	312.81
1809	93.13	231.64	280.23	279.66	219.19	1861	152.43	187.54	280.63	280.55	293.43
1810	222.51	272.30	279.99	279.67	199.86	1862	281.82	228.20	280.39	280.57	274.10
1811	351.90	312.96	279.75	279.69	180.53	1863	51.20	268.87	280.15	280.58	254.78
1812	121.28	353.63	279.51	279.71	161.20	1864	180.59	309.53	279.91	280.60	235.45
1813	263.84	34.40	280.26	279.73	141.82	1865	323.15	350.30	280.66	280.62	216.07
1814	33.23	75.06	280.02	279.74	122.49	1866	92.53	30.96	280.42	280.64	196.74
1815	162.61	115.73	279.78	279.76	103.17	1867	221.92	71.63	280.18	280.65	177.41
1816	292.00	156.39	279.54	279.78	83.84	1868	351.30	112.29	279.94	280.67	158.08
1817	74.56	197.16	280.29	279.79	64.46	1869	133.86	153.06	280.69	280.69	138.70
1818	203.94	237.82	280.05	279.81	45.13	1870	263.25	193.73	280.45	280.71	119.37
1819	333.33	278.49	279.81	279.83	25.80	1871	32.63	234.39	280.21	280.72	100.04
1820	102.71	319.15	279.57	279.85	6.47	1872	162.02	275.05	279.97	280.74	80.72
1821	245.28	359.92	280.32	279.86	347.09	1873	304.58	315.83	280.72	280.76	61.34
1822	14.66	40.59	280.08	279.88	327.76	1874	73.96	356.49	280.48	280.77	42.01
1823	144.04	81.25	279.84	279.90	308.43	1875	203.35	37.15	280.24	280.79	22.68
1824	273.43	121.91	279.61	279.91	289.11	1876	332.73	77.81	280.01	280.81	3.35
1825	55.99	162.69	280.35	279.93	269.72	1877	115.29	118.59	280.75	280.83	343.97
1826	185.38	203.35	280.11	279.95	250.40	1878	244.68	159.25	280.51	280.84	324.64
1827	314.76	244.01	279.87	279.97	231.07	1879	14.06	199.91	280.27	280.86	305.31
1828	84.15	284.67	279.64	279.98	211.74	1880	143.45	240.58	280.04	280.88	285.98
1829	226.71	323.45	280.38	280.00	192.36	1881	286.01	281.35	280.78	280.89	266.60
1830	356.09	6.11	280.14	280.02	173.03	1882	55.39	322.01	280.54	280.91	247.28
1831	125.48	46.77	279.91	280.03	153.70	1883	184.78	2.67	280.31	280.93	227.95
1832	254.86	87.43	279.67	280.05	134.37	1884	314.16	43.34	280.07	280.95	208.62
1833	37.42	128.21	280.41	280.07	114.99	1885	96.72	84.11	280.81	280.96	189.24
1834	166.81	168.87	280.18	280.09	95.66	1886	226.11	124.77	280.57	280.98	169.91
1835	296.19	209.53	279.94	280.10	76.34	1887	355.49	165.44	280.34	281.00	150.58
1836	65.58	250.20	279.70	280.12	57.01	1888	124.88	206.10	280.10	281.01	131.25
1837	208.14	290.97	280.44	280.14	37.63	1889	267.44	246.87	280.84	281.03	111.87
1838	337.52	331.63	280.21	280.16	18.30	1890	36.82	287.54	280.61	281.05	92.54
1839	106.91	12.30	279.97	280.17	358.97	1891	166.21	328.20	280.37	281.07	73.22
1840	236.29	52.96	279.73	280.19	339.64	1892	295.59	8.86	280.13	281.08	53.89
1841	18.85	93.73	280.48	280.21	320.26	1893	78.16	49.63	280.87	281.10	34.51
1842	148.24	134.39	280.24	280.22	300.93	1894	207.54	90.30	280.64	281.12	15.18
1843	277.62	175.06	280.00	280.24	281.61	1895	336.93	130.96	280.40	281.13	355.85
1844	47.01	215.72	279.76	280.26	262.28	1896	106.31	171.62	280.16	281.15	336.52
1845	189.57	256.49	280.51	280.28	242.90	1897	248.87	212.40	290.91	281.17	317.14
1846	318.95	297.16	280.27	280.29	223.57	1898	18.26	253.06	280.67	281.19	297.81
1847	88.34	337.82	280.03	280.31	204.24	1899	147.64	293.72	280.43	281.20	278.48
1848	217.72	18.48	279.79	280.33	184.91						
1849	0.28	59.26	280.54	280.34	165.53						
1850	129.67	99.92	280.30	280.36	146.20						
1851	259.65	140.58	280.06	280.38	126.87						

Table 4.—*Mean longitude of lunar and solar elements at Jan. 1, 0 hour, Greenwich mean civil time, of each year from 1800 to 2000—Continued*

Year	s	p	h	p <sub>1</sub>	N	Year	s	p	h	p <sub>1</sub>	N
	°	°	°	°	°		°	°	°	°	°
1900	277.03	334.38	280.19	281.22	259.16	1952	323.15	290.16	279.60	282.12	333.45
1901	46.41	15.05	279.95	281.24	239.83	1953	105.72	330.94	280.35	282.13	314.07
1902	175.80	55.71	279.71	281.26	220.50	1954	235.10	11.60	280.11	282.15	294.75
1903	305.18	96.37	279.47	281.27	201.17	1955	4.49	52.26	279.87	282.17	275.42
1904	74.57	137.03	279.23	281.29	181.84	1956	133.87	92.92	279.63	282.18	256.09
1905	217.13	177.81	279.98	281.31	162.46	1957	276.43	133.70	280.38	282.20	236.71
1906	346.51	218.47	279.74	281.32	143.13	1958	45.82	174.36	280.14	282.22	217.38
1907	115.90	259.13	279.50	281.34	123.81	1959	175.20	215.02	279.90	282.24	198.05
1908	245.28	299.79	279.27	281.36	104.48	1960	304.59	255.69	279.67	282.25	178.72
1909	27.84	340.57	280.01	281.38	85.10	1961	87.15	206.46	280.41	282.27	159.34
1910	157.23	21.23	279.77	281.39	65.77	1962	216.53	337.12	280.17	282.29	140.01
1911	286.61	61.89	279.53	281.41	46.44	1963	345.92	17.78	279.93	282.30	120.69
1912	56.00	102.55	279.30	281.43	27.11	1964	115.30	58.45	279.70	282.32	101.36
1913	198.56	143.33	280.04	281.44	7.73	1965	257.86	99.22	280.44	282.34	81.98
1914	327.94	183.99	279.80	281.46	348.40	1966	27.25	139.88	280.20	282.36	62.65
1915	97.33	224.65	279.57	281.48	329.07	1967	156.63	180.54	279.97	282.37	43.32
1916	226.71	265.32	279.33	281.50	309.75	1968	286.02	221.21	279.73	282.39	23.99
1917	9.27	306.09	280.07	281.51	290.36	1969	68.58	261.98	280.47	282.41	4.61
1918	138.66	346.75	279.84	281.53	271.04	1970	197.96	302.64	280.24	282.42	345.28
1919	268.04	27.41	279.60	281.55	251.71	1971	327.35	343.31	280.00	282.44	325.95
1920	37.43	68.08	279.36	281.56	232.38	1972	96.73	23.97	279.76	282.46	306.63
1921	179.99	108.85	280.10	281.58	213.00	1973	239.29	64.74	280.50	282.48	287.24
1922	309.37	149.51	279.57	281.60	193.67	1974	8.68	105.40	280.27	282.49	267.92
1923	78.76	190.18	279.63	281.62	174.34	1975	138.06	146.07	280.03	282.51	248.59
1924	208.14	230.84	279.39	281.63	155.01	1976	267.45	186.73	279.79	282.53	229.26
1925	350.71	271.61	280.14	281.65	135.63	1977	50.01	227.50	280.54	282.54	209.88
1926	120.09	312.27	279.90	281.67	116.31	1978	179.40	268.17	280.30	282.56	190.55
1927	249.47	352.94	279.66	281.69	96.98	1979	308.78	308.83	280.06	282.58	171.22
1928	18.86	33.60	279.42	281.70	77.65	1980	78.16	349.49	279.82	282.60	151.89
1929	161.42	74.37	280.17	281.72	58.27	1981	220.73	30.26	280.57	282.61	132.51
1930	290.81	115.03	279.93	281.74	38.94	1982	350.11	70.93	280.33	282.63	113.19
1931	60.19	155.70	279.69	281.75	19.61	1983	119.50	111.59	280.09	282.65	93.86
1932	189.58	196.36	279.45	281.77	0.28	1984	248.88	152.25	279.85	282.67	74.53
1933	332.14	237.13	280.20	281.79	340.90	1985	31.44	193.02	280.60	282.68	55.15
1934	101.52	277.80	279.96	281.81	321.57	1986	160.83	233.69	280.36	282.70	35.82
1935	230.91	318.46	279.72	281.82	302.25	1987	290.21	274.35	280.12	282.72	16.49
1936	0.29	359.12	279.48	281.84	282.92	1988	59.60	315.01	279.88	282.73	357.16
1937	142.85	39.89	280.23	281.86	263.54	1989	202.16	355.79	280.63	282.75	337.78
1938	272.24	80.56	279.99	281.87	244.21	1990	331.54	36.45	280.39	282.77	318.45
1939	41.62	121.22	279.75	281.89	224.88	1991	100.93	77.11	280.15	282.79	299.13
1940	171.01	161.88	279.51	281.91	205.55	1992	230.31	117.77	279.91	282.80	279.80
1941	313.57	202.65	280.26	281.93	186.17	1993	12.87	158.55	280.66	282.82	260.42
1942	82.95	243.32	280.02	281.94	166.84	1994	142.26	199.21	280.42	282.84	241.09
1943	212.34	283.98	279.78	281.96	147.51	1995	271.64	239.87	280.18	282.85	221.76
1944	341.72	324.64	279.54	281.98	128.19	1996	41.03	280.53	279.94	282.87	202.43
1945	124.28	5.42	280.29	281.99	108.80	1997	193.59	321.31	280.69	282.89	183.05
1946	253.67	46.08	280.05	282.01	89.48	1998	312.97	1.97	280.45	282.91	163.72
1947	23.05	86.74	279.81	282.03	70.15	1999	82.36	42.63	280.21	282.92	144.39
1948	152.44	127.40	279.57	282.05	50.82	2000	211.74	83.29	279.97	282.94	125.07
1949	295.00	168.18	280.32	282.06	31.44						
1950	64.38	208.04	280.08	282.08	12.11						
1951	193.77	249.50	279.84	282.10	352.78						

Table 5.—*Differences to adapt table 4 to any month, day, and hour of Greenwich mean civil time*DIFFERENCES TO FIRST OF EACH CALENDAR MONTH OF COMMON YEARS<sup>1</sup>

Month	s	p	h	p <sub>1</sub>	N	Month	s	p	h	p <sub>1</sub>	N
Jan. 1	0.00	0.00	0.00	0.00	0.00	July 1	224.93	20.16	178.40	0.01	-9.58
Feb. 1	48.47	3.45	30.56	0.00	-1.64	Aug. 1	273.40	23.62	208.96	0.01	-11.23
Mar. 1	57.41	6.57	58.15	0.00	-3.12	Sept. 1	321.86	27.07	239.51	0.01	-12.87
Apr. 1	105.88	10.03	88.71	0.00	-4.77	Oct. 1	357.16	30.41	269.08	0.01	-14.46
May 1	141.17	13.37	118.28	0.01	-6.35	Nov. 1	45.62	33.87	299.64	0.01	-16.10
June 1	189.64	16.82	148.88	0.01	-8.00	Dec. 1	80.92	37.21	329.21	0.02	-17.69

DIFFERENCES TO BEGINNING OF EACH DAY OF MONTH FOR COMMON YEARS<sup>1</sup>

Day	s	p	h	p <sub>1</sub>	N	Day	s	p	h	p <sub>1</sub>	N
	o	o	o	o	o		o	o	o	o	o
1-----	0.00	0.00	0.00	0.00	0.00	17-----	210.82	1.78	15.77	0.00	-0.85
2-----	13.18	0.11	0.99	0.00	-0.05	18-----	224.00	1.89	16.76	0.00	-0.90
3-----	26.35	0.22	1.97	0.00	-0.11	19-----	237.18	2.01	17.74	0.00	-0.95
4-----	39.53	0.33	2.96	0.00	-0.16	20-----	250.35	2.12	18.73	0.00	-1.01
5-----	52.71	0.45	3.94	0.00	-0.21	21-----	263.53	2.23	19.71	0.00	-1.06
6-----	65.88	0.56	4.93	0.00	-0.26	22-----	276.70	2.34	20.70	0.00	-1.11
7-----	79.06	0.67	5.91	0.00	-0.32	23-----	289.88	2.45	21.68	0.00	-1.16
8-----	92.23	0.78	6.90	0.00	-0.37	24-----	303.06	2.56	22.67	0.00	-1.22
9-----	105.41	0.89	7.89	0.00	-0.42	25-----	316.23	2.67	23.66	0.00	-1.27
10-----	118.59	1.00	8.87	0.00	-0.48	26-----	329.41	2.79	24.64	0.00	-1.32
11-----	131.76	1.11	9.86	0.00	-0.53	27-----	342.59	2.90	25.63	0.00	-1.38
12-----	144.94	1.23	10.84	0.00	-0.58	28-----	355.76	3.01	26.61	0.00	-1.43
13-----	158.12	1.34	11.83	0.00	-0.64	29-----	8.94	3.12	27.60	0.00	-1.48
14-----	171.29	1.45	12.81	0.00	-0.69	30-----	22.12	3.23	28.58	0.00	-1.54
15-----	184.47	1.56	13.80	0.00	-0.74	31-----	35.29	3.34	29.57	0.00	-1.59
16-----	197.65	1.67	14.78	0.00	-0.79	32-----	48.47	3.45	30.56	0.00	-1.64

## DIFFERENCES TO BEGINNING OF EACH HOUR OF DAY, GREENWICH CIVIL TIME

Hour	s	p	h	p <sub>1</sub>	N	Hour	s	p	h	p <sub>1</sub>	N
	o	o	o	o	o		o	o	o	o	o
0-----	0.00	0.00	0.00	0.00	0.00	12-----	6.59	0.06	0.49	0.00	-0.03
1-----	0.55	0.00	0.04	0.00	0.00	13-----	7.14	0.06	0.53	0.00	-0.03
2-----	1.10	0.01	0.08	0.00	0.00	14-----	7.69	0.06	0.57	0.00	-0.03
3-----	1.65	0.01	0.12	0.00	-0.01	15-----	8.24	0.07	0.62	0.00	-0.03
4-----	2.20	0.02	0.16	0.00	-0.01	16-----	8.78	0.07	0.66	0.00	-0.04
5-----	2.75	0.02	0.21	0.00	-0.01	17-----	9.33	0.08	0.70	0.00	-0.04
6-----	3.29	0.03	0.25	0.00	-0.01	18-----	9.88	0.08	0.74	0.00	-0.04
7-----	3.84	0.03	0.29	0.00	-0.02	19-----	10.43	0.09	0.78	0.00	-0.04
8-----	4.39	0.04	0.33	0.00	-0.02	20-----	10.98	0.09	0.82	0.00	-0.04
9-----	4.94	0.04	0.37	0.00	-0.02	21-----	11.53	0.10	0.86	0.00	-0.05
10-----	5.49	0.05	0.41	0.00	-0.02	22-----	12.08	0.10	0.90	0.00	-0.05
11-----	6.04	0.05	0.45	0.00	-0.02	23-----	12.63	0.11	0.94	0.00	-0.05

<sup>1</sup> The table may also be used directly for dates between Jan. 1 and Feb. 29, inclusive, of leap years; but if the required date falls between Mar. 1 and Dec. 31, inclusive, of a leap year, the day of month should be increased by one before entering the table.

Table 6.—Values of  $I$ ,  $\nu$ ,  $\xi$ ,  $\nu'$ , and  $2\nu''$  for each degree of  $N$ .

$N$	Positive always $I$	Positive when $N$ is between 0 and $180^\circ$ ; negative when $N$ is between $180$ and $360^\circ$								$N$	
		$\nu$		$\xi$		$\nu'$		$2\nu''$			
		$^\circ$	Diff.	$^\circ$	Diff.	$^\circ$	Diff.	$^\circ$	Diff.		
0	28.60	0	0.00	19	0.00	17	0.00	13	0.00	28	360
1	28.60	0	0.19	19	0.17	17	0.13	14	0.28	29	359
2	28.60	1	0.38	18	0.34	17	0.27	13	0.57	28	358
3	28.59	0	0.56	19	0.51	16	0.40	14	0.85	29	357
4	28.59	1	0.75	19	0.67	17	0.54	13	1.14	28	356
5	28.58	0	0.94	18	0.84	17	0.67	13	1.42	28	355
6	28.58	1	1.12	19	1.01	17	0.80	14	1.70	29	354
7	28.57	1	1.31	19	1.18	17	0.94	13	1.99	28	353
8	28.56	1	1.50	18	1.35	16	1.07	13	2.27	28	352
9	28.55	2	1.68	19	1.51	17	1.20	14	2.55	28	351
10	28.53	1	1.87	18	1.68	17	1.34	13	2.83	28	350
11	28.52	2	2.05	19	1.85	17	1.47	13	3.11	28	349
12	28.50	1	2.24	18	2.02	16	1.60	13	3.39	28	348
13	28.49	2	2.42	19	2.18	17	1.73	13	3.67	28	347
14	28.47	2	2.61	18	2.35	16	1.86	13	3.95	28	346
15	28.45	2	2.79	19	2.51	17	1.99	13	4.23	28	345
16	28.43	2	2.98	18	2.68	16	2.12	13	4.51	27	344
17	28.41	2	3.16	18	2.84	17	2.25	13	4.78	28	343
18	28.39	3	3.34	18	3.01	16	2.38	13	5.06	27	342
19	28.36	2	3.52	18	3.17	17	2.51	13	5.33	27	341
20	28.34	3	3.70	18	3.34	16	2.64	13	5.60	27	340
21	28.31	2	3.88	18	3.50	16	2.77	13	5.87	27	339
22	28.29	3	4.06	18	3.66	16	2.90	13	6.14	27	338
23	28.26	3	4.24	18	3.82	16	3.03	12	6.41	27	337
24	28.23	3	4.42	18	3.98	16	3.15	13	6.68	26	336
25	28.20	4	4.60	18	4.14	16	3.28	12	6.94	27	335
26	28.16	3	4.78	17	4.30	16	3.40	13	7.21	26	334
27	28.13	4	4.95	18	4.46	16	3.53	12	7.47	26	333
28	28.09	3	5.13	17	4.62	16	3.65	13	7.73	26	332
29	28.06	4	5.30	18	4.78	16	3.78	12	7.99	26	331
30	28.02	4	5.48	17	4.94	16	3.90	12	8.25	25	330
31	27.98	4	5.65	17	5.10	15	4.02	12	8.50	25	329
32	27.94	4	5.82	17	5.25	16	4.14	12	8.75	25	328
33	27.90	4	5.99	17	5.41	15	4.26	12	9.00	25	327
34	27.86	4	6.16	17	5.56	15	4.38	12	9.25	25	326
35	27.82	5	6.33	17	5.71	15	4.50	12	9.50	25	325
36	27.77	4	6.50	16	5.86	15	4.62	12	9.74	24	324
37	27.73	5	6.66	17	6.01	15	4.74	11	9.98	24	323
38	27.68	5	6.83	16	6.16	15	4.85	12	10.22	24	322
39	27.63	5	6.99	16	6.31	15	4.97	11	10.46	23	321
40	27.58	5	7.15	16	6.46	15	5.08	11	10.69	24	320
41	27.53	5	7.31	16	6.61	14	5.19	11	10.93	23	319
42	27.48	5	7.47	16	6.75	15	5.30	11	11.16	22	318
43	27.43	5	7.63	16	6.90	14	5.41	11	11.38	22	317
44	27.38	6	7.79	15	7.04	14	5.52	11	11.60	22	316
45	27.32	6	7.94	15	7.18	14	5.63	11	11.82	22	315

Table 6.—Values of  $I$ ,  $\nu$ ,  $\xi$ ,  $\nu'$ , and  $2\nu''$  for each degree of  $N$ —Continued

$N$	Positive always		Positive when $N$ is between 0 and $180^\circ$ ; negative when $N$ is between $180$ and $360^\circ$								$N$	
			$\nu$		$\xi$		$\nu'$		$2\nu''$			
	$I$		$\circ$	$Diff.$	$\circ$	$Diff.$	$\circ$	$Diff.$	$\circ$	$Diff.$		
45.....	°	27.32	5	7.94	16	7.18	14	5.63	11	11.82	22	315
46.....	27.27	6	8.10	15	7.32	14	5.74	10	12.04	22	314	
47.....	27.21	6	8.25	15	7.46	14	5.84	11	12.26	21	313	
48.....	27.15	6	8.40	15	7.60	13	5.95	10	12.47	21	312	
49.....	27.09	6	8.55	14	7.73	14	6.05	10	12.68	20	311	
50.....	27.03	6	8.69	15	7.87	13	6.15	10	12.88	20	310	
51.....	26.97	6	8.84	14	8.00	14	6.25	10	13.08	20	309	
52.....	26.91	6	8.98	14	8.14	13	6.35	10	13.28	20	308	
53.....	26.85	7	9.12	14	8.27	13	6.45	9	13.48	19	307	
54.....	26.78	6	9.26	14	8.40	12	6.54	10	13.67	19	306	
55.....	26.72	7	9.40	14	8.52	13	6.64	9	13.86	19	305	
56.....	26.65	6	9.54	13	8.65	12	6.73	9	14.05	18	304	
57.....	26.59	7	9.67	14	8.77	13	6.82	9	14.23	17	303	
58.....	26.52	7	9.81	13	8.90	12	6.91	9	14.40	18	302	
59.....	26.45	7	9.94	13	9.02	12	7.00	9	14.58	17	301	
60.....	26.38	7	10.07	12	9.14	11	7.09	8	14.75	17	300	
61.....	26.31	7	10.19	13	9.25	12	7.17	9	14.92	16	299	
62.....	26.24	7	10.32	12	9.37	11	7.26	8	15.08	16	298	
63.....	26.17	7	10.44	12	9.48	11	7.34	8	15.24	15	297	
64.....	26.10	7	10.56	12	9.59	11	7.42	7	15.39	15	296	
65.....	26.03	8	10.68	11	9.70	11	7.49	8	15.54	15	295	
66.....	25.95	7	10.79	11	9.81	11	7.57	7	15.69	14	294	
67.....	25.88	8	10.90	11	9.92	10	7.64	8	15.83	13	293	
68.....	25.80	8	11.01	11	10.02	10	7.72	7	15.96	14	292	
69.....	25.72	7	11.12	11	10.12	10	7.79	7	16.10	13	291	
70.....	25.65	8	11.23	10	10.22	10	7.86	6	16.23	12	290	
71.....	25.57	8	11.33	10	10.32	9	7.92	7	16.35	12	289	
72.....	25.49	8	11.43	10	10.41	9	7.99	6	16.47	11	288	
73.....	25.41	8	11.53	10	10.50	9	8.05	6	16.58	11	287	
74.....	25.33	8	11.63	9	10.59	9	8.11	6	16.69	11	286	
75.....	25.25	8	11.72	9	10.68	9	8.17	6	16.80	10	285	
76.....	25.17	8	11.81	8	10.77	8	8.23	5	16.90	10	284	
77.....	25.09	8	11.89	9	10.85	8	8.28	6	17.00	9	283	
78.....	25.01	9	11.98	8	10.93	8	8.34	5	17.09	8	282	
79.....	24.92	8	12.06	8	11.01	7	8.39	5	17.17	8	281	
80.....	24.84	8	12.14	7	11.08	7	8.44	4	17.25	8	280	
81.....	24.76	9	12.21	7	11.15	7	8.48	5	17.33	7	279	
82.....	24.67	8	12.28	7	11.22	7	8.53	4	17.40	6	278	
83.....	24.59	9	12.35	7	11.29	7	8.57	4	17.46	6	277	
84.....	24.50	8	12.42	6	11.36	6	8.61	3	17.52	6	276	
85.....	24.42	9	12.48	6	11.42	5	8.64	4	17.58	5	275	
86.....	24.33	9	12.54	6	11.47	6	8.68	3	17.63	4	274	
87.....	24.24	8	12.60	5	11.53	5	8.71	3	17.67	4	273	
88.....	24.16	9	12.65	5	11.58	5	8.74	2	17.71	3	272	
89.....	24.07	9	12.70	5	11.63	5	8.76	3	17.74	3	271	
90.....	23.98	9	12.75	5	11.68	5	8.79	3	17.77	2	270	

Table 6.—Values of  $I$ ,  $\nu$ ,  $\xi$ ,  $\nu'$ , and  $2\nu''$  for each degree of  $N$ —Continued

$N$	Positive always		Positive when $N$ is between 0 and $180^\circ$ ; negative when $N$ is between $180$ and $360^\circ$								$N$
			$\nu$		$\xi$		$\nu'$		$2\nu''$		
	$I$		$\nu$	$\xi$	$\nu'$	$2\nu''$	$\nu$	$\xi$	$\nu'$	$2\nu''$	
°	°	Diff.	°	Diff.	°	Diff.	°	Diff.	°	Diff.	°
90	23.98	9	12.75	4	11.68	4	8.79	2	17.77	2	270
91	23.89	9	12.79	4	11.72	4	8.81	2	17.79	2	269
92	23.80	8	12.83	4	11.76	4	8.83	2	17.81	1	268
93	23.72	9	12.87	3	11.80	3	8.85	1	17.82	1	267
94	23.63	9	12.90	3	11.83	3	8.86	1	17.83	0	266
95	23.54	9	12.93	2	11.86	3	8.87	1	17.83	1	265
96	23.45	9	12.95	2	11.89	3	8.88	1	17.82	1	264
97	23.36	9	12.97	2	11.92	2	8.89	1	17.81	2	263
98	23.27	9	12.99	2	11.94	1	8.90	0	17.79	2	262
99	23.18	9	13.01	1	11.95	1	8.90	1	17.77	3	261
100	23.09	9	13.02	0	11.96	1	8.89	0	17.74	3	260
101	23.00	9	13.02	0	11.97	1	8.89	1	17.71	4	259
102	22.91	9	13.02	0	11.98	0	8.88	1	17.67	5	258
103	22.82	9	13.02	1	11.98	0	8.87	1	17.62	5	257
104	22.73	9	13.01	1	11.98	1	8.86	2	17.57	6	256
105	22.64	9	13.00	1	11.97	1	8.84	2	17.51	6	255
106	22.55	9	12.99	2	11.96	1	8.82	2	17.45	7	254
107	22.46	9	12.97	2	11.95	2	8.80	2	17.38	8	253
108	22.37	9	12.95	3	11.93	2	8.78	3	17.30	8	252
109	22.28	8	12.92	3	11.91	2	8.75	3	17.22	8	251
110	22.20	9	12.89	4	11.89	3	8.72	3	17.14	9	250
111	22.11	9	12.85	4	11.86	3	8.69	4	17.05	10	249
112	22.02	9	12.81	4	11.83	4	8.65	4	16.95	11	248
113	21.93	9	12.77	5	11.79	4	8.61	4	16.84	11	247
114	21.84	9	12.72	5	11.75	5	8.57	5	16.73	11	246
115	21.75	8	12.67	6	11.70	5	8.52	4	16.62	12	245
116	21.67	9	12.61	6	11.65	5	8.48	5	16.50	13	244
117	21.58	8	12.55	7	11.60	6	8.43	6	16.37	13	243
118	21.50	9	12.48	7	11.54	6	8.37	6	16.24	14	242
119	21.41	9	12.41	8	11.48	7	8.31	6	16.10	14	241
120	21.32	8	12.33	8	11.41	7	8.25	6	15.96	15	240
121	21.24	9	12.25	8	11.34	8	8.19	6	15.81	15	239
122	21.15	8	12.17	9	11.26	8	8.13	7	15.66	16	238
123	21.07	8	12.08	10	11.18	8	8.06	7	15.50	17	237
124	20.99	8	11.98	10	11.10	9	7.99	8	15.33	17	236
125	20.91	9	11.88	10	11.01	9	7.91	8	15.16	17	235
126	20.82	8	11.78	11	10.92	10	7.83	8	14.99	18	234
127	20.74	8	11.67	12	10.82	10	7.75	8	14.81	19	233
128	20.66	8	11.55	12	10.72	11	7.67	9	14.62	19	232
129	20.58	7	11.43	12	10.61	11	7.58	9	14.43	20	231
130	20.51	8	11.31	13	10.50	12	7.49	9	14.23	20	230
131	20.43	8	11.18	13	10.38	12	7.40	10	14.03	20	229
132	20.35	7	11.05	14	10.26	13	7.30	10	13.83	21	228
133	20.28	8	10.91	14	10.13	13	7.20	10	13.62	22	227
134	20.20	8	10.77	15	10.00	13	7.10	10	13.40	22	226
135	20.13	7	10.62	15	9.87	13	7.00	10	13.18	22	225

Table 6.—Values of  $I$ ,  $\nu$ ,  $\xi$ ,  $\nu'$ , and  $2\nu''$  for each degree of  $N$ —Continued

$N$	Positive always	Positive when $N$ is between 0 and $180^\circ$ ; negative when $N$ is between $180$ and $360^\circ$										$N$	
		$I$		$\nu$		$\xi$		$\nu'$		$2\nu''$			
		$^\circ$	Dif.	$^\circ$	Dif.	$^\circ$	Dif.	$^\circ$	Dif.	$^\circ$	Dif.		
135.	20.13	8	10.62	15	9.87	14	7.00	11	13.18	22	225		
136.	20.05	7	10.47	16	9.73	14	6.89	11	12.96	23	224		
137.	19.98	7	10.31	16	9.59	15	6.78	12	12.73	24	223		
138.	19.91	7	10.15	17	9.44	15	6.66	11	12.49	24	222		
139.	19.84	7	9.98	17	9.29	16	6.55	12	12.25	24	221		
140.	19.77	6	9.81	18	9.13	16	6.43	12	12.01	25	220		
141.	19.71	7	9.63	18	8.97	16	6.31	13	11.76	25	219		
142.	19.64	6	9.45	18	8.80	17	6.18	12	11.51	25	218		
143.	19.58	7	9.27	19	8.63	17	6.06	13	11.26	26	217		
144.	19.51	6	9.08	19	8.46	18	5.93	13	11.00	26	216		
145.	19.45	6	8.89	20	8.28	18	5.80	14	10.74	26	215		
146.	19.39	6	8.69	20	8.10	19	5.66	14	10.48	27	214		
147.	19.33	6	8.49	21	7.91	19	5.52	14	10.21	27	213		
148.	19.27	5	8.28	21	7.72	20	5.38	14	9.94	28	212		
149.	19.22	6	8.07	22	7.52	20	5.24	15	9.66	28	211		
150.	19.16	5	7.85	22	7.32	20	5.09	14	9.38	28	210		
151.	19.11	6	7.63	22	7.12	21	4.95	15	9.10	29	209		
152.	19.05	5	7.41	23	6.91	21	4.80	15	8.81	29	208		
153.	19.00	5	7.18	23	6.70	21	4.65	15	8.52	29	207		
154.	18.95	4	6.95	23	6.49	22	4.50	16	8.23	29	206		
155.	18.91	5	6.72	24	6.27	22	4.34	15	7.94	30	205		
156.	18.86	4	6.48	24	6.05	23	4.19	16	7.64	30	204		
157.	18.82	4	6.24	25	5.82	23	4.03	16	7.34	30	203		
158.	18.78	4	5.99	25	5.59	23	3.87	17	7.04	30	202		
159.	18.74	4	5.74	25	5.36	23	3.70	16	6.74	31	201		
160.	18.70	4	5.49	25	5.13	24	3.54	17	6.43	31	200		
161.	18.66	4	5.24	26	4.89	24	3.37	17	6.12	31	199		
162.	18.62	3	4.98	26	4.65	24	3.20	17	5.81	31	198		
163.	18.59	3	4.72	26	4.41	25	3.03	17	5.50	31	197		
164.	18.56	3	4.46	27	4.16	25	2.86	17	5.19	32	196		
165.	18.53	3	4.19	27	3.91	25	2.69	17	4.87	32	195		
166.	18.50	3	3.92	27	3.66	25	2.52	18	4.55	32	194		
167.	18.47	2	3.65	27	3.41	25	2.34	17	4.23	32	193		
168.	18.45	2	3.38	28	3.16	26	2.17	18	3.91	32	192		
169.	18.43	2	3.10	27	2.90	26	1.99	18	3.59	32	191		
170.	18.41	2	2.83	28	2.64	26	1.81	18	3.27	33	190		
171.	18.39	2	2.55	28	2.38	26	1.63	18	2.94	32	189		
172.	18.37	1	2.27	28	2.12	26	1.45	18	2.62	33	188		
173.	18.36	2	1.99	28	1.86	26	1.27	18	2.29	32	187		
174.	18.34	1	1.71	29	1.60	27	1.09	18	1.97	33	186		
175.	18.33	1	1.42	28	1.33	26	0.91	18	1.64	33	185		
176.	18.32	0	1.14	28	1.07	27	0.73	18	1.31	32	184		
177.	18.32	1	0.86	29	0.80	26	0.55	18	0.99	33	183		
178.	18.31	0	0.57	28	0.54	27	0.37	19	0.66	33	182		
179.	18.31	0	0.29	29	0.27	27	0.18	18	0.33	33	181		
180.	18.31	0	0.00	0.00	0.00	0.00	0.00	18	0.00	33	180		

Table 7.—*Log R<sub>a</sub> for amplitude of constituent L<sub>2</sub>*

<i>P</i>	<i>I</i>	18°	19°	20°	21°	22°	23°	24°	25°	26°	27°	28°	29°
°	°	0	0.0708	0.0799	0.0897	0.1002	0.1117	0.1240	0.1373	0.1517	0.1674	0.1843	0.2027
5	180	0.0695	0.0783	0.0878	0.0981	0.1092	0.1211	0.1340	0.1479	0.1628	0.1790	0.1966	0.2155
10	190	0.0654	0.0736	0.0824	0.0918	0.1019	0.1128	0.1244	0.1367	0.1500	0.1641	0.1792	0.1953
15	195	0.0590	0.0662	0.0739	0.0820	0.0906	0.0998	0.1085	0.1197	0.1305	0.1417	0.1535	0.1658
20	200	0.0506	0.0565	0.0628	0.0693	0.0762	0.0834	0.0908	0.0986	0.1065	0.1147	0.1230	0.1313
25	205	0.0407	0.0452	0.0498	0.0546	0.0596	0.0647	0.0698	0.0750	0.0802	0.0853	0.0904	0.0952
30	210	0.0297	0.0327	0.0357	0.0388	0.0418	0.0449	0.0478	0.0506	0.0533	0.0557	0.0578	0.0597
35	215	0.0182	0.0197	0.0212	0.0225	0.0238	0.0249	0.0258	0.0265	0.0269	0.0270	0.0268	0.0262
40	220	0.0065	0.0066	0.0066	0.0064	0.0060	0.0054	0.0045	0.0034	0.0019	0.0001	0.0001	0.0001
45	225	9.9951	9.9940	9.9926	9.9910	9.9891	9.9870	9.9846	9.9819	9.9789	9.9755	9.9717	9.9676
50	230	9.9843	9.9820	9.9794	9.9765	9.9734	9.9700	9.9663	9.9623	9.9580	9.9533	9.9483	9.9430
55	235	9.9773	9.9710	9.9673	9.9634	9.9592	9.9548	9.9500	9.9449	9.9395	9.9338	9.9276	9.9215
60	240	9.9653	9.9611	9.9566	9.9518	9.9467	9.9414	9.9357	9.9298	9.9235	9.9170	9.9102	9.9031
65	245	9.9575	9.9526	9.9473	9.9418	9.9360	9.9299	9.9236	9.9169	9.9100	9.9029	9.8954	9.8877
70	250	9.9510	9.9454	9.9396	9.9335	9.9271	9.9205	9.9136	9.9064	9.8990	9.8913	9.8834	9.8753
75	255	9.9458	9.9398	9.9336	9.9270	9.9202	9.9131	9.9058	9.8982	9.8904	9.8824	9.8742	9.8657
80	260	9.9421	9.9358	9.9292	9.9224	9.9152	9.9079	9.9003	9.8924	9.8844	9.8761	9.8676	9.8589
85	265	9.9399	9.9334	9.9266	9.9196	9.9123	9.9047	9.8969	9.8889	9.8807	9.8723	9.8636	9.8548
90	270	9.9391	9.9326	9.9257	9.9186	9.9113	9.9037	9.8958	9.8878	9.8795	9.8710	9.8623	9.8535

Table 8.—*Values of R for argument of constituent L<sub>2</sub>*

[Tabular values are positive when  $P$  is in first or third quadrants, negative when  $P$  is in second or fourth quadrants]

Table 9.—*Log Q<sub>a</sub> for amplitude of constituent M<sub>1</sub>*

<i>P</i>	<i>Log Q<sub>a</sub></i>	Diff.	<i>P</i>	<i>P</i>	<i>Log Q<sub>a</sub></i>	Diff.	<i>P</i>
° 0	° 180	9.7133	° 0	° 180	360	45	° 135
1	181	9.7133	2	179	359	46	134
2	182	9.7135	2	178	358	47	133
3	183	9.7137	4	177	357	48	132
4	184	9.7141	4	176	356	49	131
5	185	9.7145	6	175	355	50	130
6	186	9.7151	7	174	354	51	129
7	187	9.7158	7	173	353	52	128
8	188	9.7165	9	172	352	53	127
9	189	9.7174	10	171	351	54	126
10	190	9.7184	10	170	350	55	125
11	191	9.7194	12	169	349	56	124
12	192	9.7206	13	168	348	57	123
13	193	9.7219	13	167	347	58	122
14	194	9.7232	15	166	346	59	121
15	195	9.7247	16	165	345	60	120
16	196	9.7263	17	164	344	61	119
17	197	9.7280	18	163	343	62	118
18	198	9.7298	19	162	342	63	117
19	199	9.7317	20	161	341	64	116
20	200	9.7337	21	160	340	65	115
21	201	9.7358	22	159	339	66	114
22	202	9.7380	23	158	338	67	113
23	203	9.7403	24	157	337	68	112
24	204	9.7427	25	156	336	69	111
25	205	9.7452	27	155	335	70	110
26	206	9.7479	27	154	334	71	109
27	207	9.7506	28	153	333	72	108
28	208	9.7534	30	152	332	73	107
29	209	9.7564	31	151	331	74	106
30	210	9.7595	31	150	330	75	105
31	211	9.7626	33	149	329	76	104
32	212	9.7659	34	148	328	77	103
33	213	9.7693	35	147	327	78	102
34	214	9.7728	36	146	326	79	101
35	215	9.7764	37	145	325	80	100
36	216	9.7801	38	144	324	81	99
37	217	9.7839	39	143	323	82	98
38	218	9.7878	40	142	322	83	97
39	219	9.7918	42	141	321	84	96
40	220	9.7960	42	140	320	85	95
41	221	9.8002	43	139	319	86	94
42	222	9.8045	45	138	318	87	93
43	223	9.8090	46	137	317	88	92
44	224	9.8136	46	136	316	89	91
45	225	9.8182	45	135	315	90	90

Table 10.—Values of  $Q$  for argument of constituent  $M_1$ 

$P$	$Q$	Diff.	$P$	$Q$	Diff.	$P$	$Q$	Diff.	$P$	$Q$	Diff.
0	0.0	0.5	45	25.8	0.8	90	90.0	2.1	135	154.2	0.8
1	0.5	0.5	46	26.6	0.8	91	92.1	2.0	136	155.0	0.8
2	1.0	0.5	47	27.4	0.8	92	94.1	2.1	137	155.8	0.7
3	1.5	0.4	48	28.2	0.9	93	96.2	2.0	138	156.5	0.7
4	1.9	0.5	49	29.1	0.8	94	98.2	2.1	139	157.2	0.7
5	2.4	0.5	50	29.9	0.9	95	100.3	2.0	140	157.9	0.7
6	2.9	0.5	51	30.8	0.9	96	102.3	2.0	141	158.6	0.7
7	3.4	0.5	52	31.7	1.0	97	104.3	1.9	142	159.3	0.7
8	3.9	0.5	53	32.7	0.9	98	106.2	2.0	143	160.0	0.7
9	4.4	0.5	54	33.6	1.0	99	108.2	1.9	144	160.7	0.6
10	4.9	0.5	55	34.6	1.0	100	110.1	1.8	145	161.3	
11	5.4	0.5	56	35.6	1.0	101	111.9	1.9	146	162.0	0.7
12	5.9	0.5	57	36.6	1.1	102	113.8	1.7	147	162.6	0.6
13	6.4	0.5	58	37.7	1.1	103	115.5	1.8	148	163.2	0.6
14	6.9	0.5	59	38.8	1.1	104	117.3	1.7	149	163.8	0.6
15	7.4	0.5	60	39.9	1.2	105	119.0	1.7	150	164.4	0.6
16	7.9	0.5	61	41.1	1.2	106	120.7	1.6	151	165.0	
17	8.4	0.5	62	42.3	1.2	107	122.3	1.6	152	165.6	0.6
18	8.9	0.5	63	43.5	1.2	108	123.9	1.6	153	166.2	0.5
19	9.4	0.6	64	44.7	1.3	109	125.5	1.5	154	166.7	
20	10.0	0.5	65	46.0	1.3	110	127.0	1.5	155	167.3	0.6
21	10.5	0.5	66	47.3	1.4	111	128.5	1.4	156	167.9	0.5
22	11.0	0.6	67	48.7	1.4	112	129.9	1.4	157	168.4	
23	11.6	0.5	68	50.1	1.4	113	131.3	1.4	158	169.0	0.5
24	12.1	0.6	69	51.5	1.5	114	132.7	1.3	159	169.5	0.5
25	12.7	0.6	70	53.0	1.5	115	134.0	1.3	160	170.0	
26	13.3	0.5	71	54.5	1.6	116	135.3	1.2	161	170.6	0.5
27	13.8	0.6	72	56.1	1.6	117	136.5	1.2	162	171.1	0.5
28	14.4	0.6	73	57.7	1.6	118	137.7	1.2	163	171.6	0.5
29	15.0	0.6	74	59.3	1.7	119	138.9	1.2	164	172.1	0.5
30	15.6	0.6	75	61.0	1.7	120	140.1	1.1	165	172.6	0.5
31	16.2	0.6	76	62.7	1.8	121	141.2	1.1	166	173.1	
32	16.8	0.6	77	64.5	1.7	122	142.3	1.1	167	173.6	0.5
33	17.4	0.6	78	66.2	1.9	123	143.4	1.0	168	174.1	0.5
34	18.0	0.7	79	68.1	1.8	124	144.4	1.0	169	174.6	
35	18.7	0.6	80	69.9	1.9	125	145.4	1.0	170	175.1	0.5
36	19.3	0.7	81	71.8	2.0	126	146.4	0.9	171	175.6	0.5
37	20.0	0.7	82	73.8	1.9	127	147.3	1.0	172	176.1	
38	20.7	0.7	83	75.7	2.0	128	148.3	0.9	173	176.6	0.5
39	21.4	0.7	84	77.7	2.0	129	149.2	0.9	174	177.1	0.5
40	22.1	0.7	85	79.7	2.1	130	150.1	0.8	175	177.6	
41	22.8	0.7	86	81.8	2.0	131	150.9	0.9	176	178.1	0.4
42	23.5	0.7	87	83.8	2.1	132	151.8	0.8	177	178.5	0.5
43	24.2	0.8	88	85.9	2.0	133	152.6	0.8	178	179.0	
44	25.0	0.8	89	87.9	2.1	134	153.4	0.8	179	179.5	0.5
45	25.8	0.8	90	90.0	2.1	135	154.2	0.8	180	180.0	

Table 10.—Values of  $Q$  for argument of constituent  $M_1$ —Continued

P	Q	Diff.									
◦	◦		◦	◦		◦	◦		◦	◦	
180	180.0	0.5	225	205.8	0.8	270	270.0	2.1	315	334.2	0.8
181	180.5	0.5	226	206.6	0.8	271	272.1	2.0	316	335.0	0.8
182	181.0	0.5	227	207.4	0.8	272	274.1	2.1	317	335.8	0.7
183	181.5	0.4	228	208.2	0.9	273	276.2	2.0	318	336.5	0.7
184	181.9	0.5	229	209.1	0.8	274	278.2		319	337.2	
185	182.4	0.5	230	209.9	0.9	275	280.3	2.1	320	337.9	0.7
186	182.9	0.5	231	210.8	0.9	276	282.3	2.0	321	338.6	0.7
187	183.4	0.5	232	211.7	1.0	277	284.3		322	339.3	
188	183.9	0.5	233	212.7	1.0	278	286.2	1.9	323	340.0	0.7
189	184.4	0.5	234	213.6	0.9	279	288.2	2.0	324	340.7	0.6
190	184.9	0.5	235	214.6	1.0	280	290.1	1.8	325	341.3	0.7
191	185.4	0.5	236	215.6	1.0	281	291.9	1.9	326	342.0	0.6
192	185.9	0.5	237	216.6	1.1	282	293.8	1.7	327	342.6	0.6
193	186.4	0.5	238	217.7	1.1	283	295.5		328	343.2	0.6
194	186.9	0.5	239	218.8	1.1	284	297.3	1.8	329	343.8	0.6
195	187.4	0.5	240	219.9	1.2	285	299.0	1.7	330	344.4	0.6
196	187.9	0.5	241	221.1	1.2	286	300.7	1.6	331	345.0	0.6
197	188.4	0.5	242	222.3	1.2	287	302.3	1.6	332	345.6	0.6
198	188.9	0.5	243	223.5	1.2	288	303.9	1.6	333	346.2	0.5
199	189.4	0.6	244	224.7		289	305.5		334	346.7	
200	190.0	0.5	245	226.0	1.3	290	307.0	1.5	335	347.3	0.6
201	190.5	0.5	246	227.3	1.4	291	308.5	1.4	336	347.9	0.5
202	191.0	0.6	247	228.7	1.4	292	309.9		337	348.4	0.6
203	191.6	0.5	248	230.1	1.4	293	311.3	1.4	338	349.0	0.5
204	192.1	0.6	249	231.5	1.5	294	312.7	1.3	339	349.5	0.5
205	192.7	0.6	250	233.0	1.5	295	314.0		340	350.0	0.6
206	193.3	0.5	251	234.5	1.6	296	315.3	1.2	341	350.6	0.5
207	193.8	0.6	252	236.1	1.6	297	316.5	1.2	342	351.1	0.5
208	194.4	0.6	253	237.7	1.6	298	317.7		343	351.6	0.5
209	195.0	0.6	254	239.3	1.7	299	318.9	1.2	344	352.1	0.5
210	195.6	0.6	255	241.0	1.7	300	320.1	1.1	345	352.6	0.5
211	196.2	0.6	256	242.7	1.8	301	321.2	1.1	346	353.1	0.5
212	196.8	0.6	257	244.5	1.7	302	322.3	1.1	347	353.6	0.5
213	197.4	0.6	258	246.2	1.9	303	323.4	1.0	348	354.1	0.5
214	198.0	0.7	259	248.1	1.8	304	324.4	1.0	349	354.6	0.5
215	198.7	0.6	260	249.9	1.9	305	325.4	1.0	350	355.1	0.5
216	199.3	0.7	261	251.8	2.0	306	326.4	0.9	351	355.6	0.5
217	200.0	0.7	262	253.8	1.9	307	327.3	1.0	352	356.1	0.5
218	200.7	0.7	263	255.7	2.0	308	328.3	0.9	353	356.6	0.5
219	201.4	0.7	264	257.7	2.0	309	329.2	0.9	354	357.1	0.5
220	202.1	0.7	265	259.7	2.1	310	330.1	0.8	355	357.6	
221	202.8	0.7	266	261.8	2.0	311	330.9	0.9	356	358.1	0.4
222	203.5	0.7	267	263.8	2.1	312	331.8	0.8	357	358.5	0.5
223	204.2	0.8	268	265.9	2.0	313	332.6	0.8	358	359.0	
224	205.0	0.8	269	267.9	2.1	314	333.4	0.8	359	359.5	0.5
225	205.8	0.8	270	270.0	2.1	315	334.2	0.8	360	360.0	

Table 11.—Values of  $u$  for equilibrium arguments[Use sign at head of column when  $N$  is between 0 and  $180^\circ$ , reverse sign when  $N$  is between  $180$  and  $360^\circ$ ]

$N$	$J_1$	$K_1$	$K_2$	$M_2, N_2$ $2N, MS$ $\lambda, \mu, \nu$	$M_3$	$M_4, MN$	$M_5$	$M_8$	$O_1, Q_1$ $2Q, \rho$	$OO$	$MK$	$2MK$	$Mf$	$N$
o	o	o	o	o	o	o	o	o	o	o	o	o	o	o
0	—	—	—	—	0.00	0.00	0.00	0.00	—	—	—	—	—	360
0	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	360
1	0.19	0.13	0.28	0.04	0.05	0.08	0.11	0.15	0.15	0.53	0.17	0.06	0.34	359
2	0.38	0.27	0.57	0.08	0.11	0.15	0.23	0.30	0.30	1.05	0.34	0.12	0.67	358
3	0.56	0.40	0.85	0.11	0.17	0.23	0.34	0.45	0.45	1.57	0.52	0.17	1.01	357
4	0.75	0.54	1.14	0.15	0.23	0.30	0.45	0.60	0.60	2.10	0.69	0.23	1.35	356
5	0.94	0.67	1.42	0.19	0.28	0.38	0.56	0.75	0.75	2.62	0.86	0.29	1.68	355
6	1.12	0.80	1.70	0.23	0.34	0.45	0.68	0.90	0.90	3.14	1.03	0.35	2.02	354
7	1.31	0.94	1.99	0.26	0.40	0.53	0.79	1.05	1.05	3.67	1.20	0.41	2.36	353
8	1.50	1.07	2.27	0.30	0.45	0.60	0.90	1.20	1.20	4.19	1.37	0.47	2.69	352
9	1.68	1.20	2.55	0.34	0.51	0.68	1.01	1.35	1.35	4.71	1.54	0.53	3.03	351
10	1.87	1.34	2.83	0.37	0.56	0.75	1.12	1.49	1.49	5.23	1.71	0.59	3.36	350
11	2.05	1.47	3.11	0.41	0.62	0.82	1.24	1.64	1.64	5.75	1.88	0.64	3.70	349
12	2.24	1.60	3.39	0.45	0.67	0.90	1.34	1.79	1.79	6.27	2.05	0.70	4.03	348
13	2.42	1.73	3.67	0.48	0.73	0.97	1.45	1.94	1.94	6.79	2.21	0.76	4.36	347
14	2.61	1.86	3.95	0.52	0.78	1.04	1.56	2.09	2.09	7.31	2.38	0.82	4.70	346
15	2.79	1.99	4.23	0.56	0.84	1.12	1.67	2.23	2.23	7.82	2.55	0.88	5.03	345
16	2.98	2.12	4.51	0.60	0.89	1.19	1.79	2.38	2.38	8.34	2.72	0.93	5.36	344
17	3.16	2.25	4.78	0.63	0.95	1.26	1.90	2.53	2.53	8.85	2.89	0.99	5.69	343
18	3.34	2.38	5.06	0.67	1.00	1.34	2.00	2.67	2.68	9.36	3.05	1.05	6.02	342
19	3.52	2.51	5.33	0.70	1.06	1.41	2.11	2.81	2.82	9.87	3.22	1.11	6.35	341
20	3.71	2.64	5.60	0.74	1.11	1.48	2.21	2.95	2.97	10.38	3.38	1.17	6.67	340
21	3.89	2.77	5.87	0.77	1.16	1.55	2.32	3.09	3.11	10.89	3.54	1.23	7.00	339
22	4.07	2.90	6.14	0.81	1.21	1.62	2.42	3.23	3.26	11.39	3.71	1.28	7.33	338
23	4.25	3.03	6.41	0.84	1.26	1.69	2.53	3.37	3.40	11.89	3.87	1.34	7.65	337
24	4.42	3.15	6.68	0.88	1.31	1.75	2.63	3.51	3.55	12.39	4.03	1.40	7.97	336
25	4.60	3.28	6.94	0.91	1.37	1.82	2.73	3.64	3.69	12.89	4.19	1.46	8.29	335
26	4.78	3.40	7.21	0.94	1.42	1.89	2.83	3.78	3.83	13.39	4.35	1.52	8.61	334
27	4.96	3.53	7.47	0.98	1.47	1.96	2.94	3.92	3.98	13.89	4.51	1.57	8.93	333
28	5.13	3.65	7.73	1.01	1.52	2.02	3.04	4.05	4.12	14.38	4.67	1.63	9.25	332
29	5.30	3.78	7.99	1.04	1.57	2.09	3.13	4.18	4.26	14.87	4.82	1.69	9.57	331
30	5.48	3.90	8.24	1.08	1.62	2.16	3.23	4.31	4.40	15.36	4.98	1.75	9.88	330
31	5.65	4.02	8.50	1.11	1.67	2.22	3.33	4.45	4.54	15.84	5.13	1.80	10.19	329
32	5.82	4.14	8.75	1.14	1.72	2.29	3.43	4.58	4.68	16.32	5.29	1.86	10.50	328
33	5.99	4.26	9.00	1.17	1.76	2.35	3.52	4.70	4.82	16.80	5.44	1.92	10.81	327
34	6.16	4.38	9.25	1.20	1.81	2.41	3.61	4.82	4.96	17.28	5.59	1.97	11.12	326
35	6.33	4.50	9.50	1.24	1.85	2.47	3.71	4.94	5.10	17.76	5.74	2.03	11.43	325
36	6.50	4.62	9.74	1.27	1.90	2.53	3.80	5.06	5.23	18.23	5.89	2.09	11.73	324
37	6.66	4.74	9.98	1.30	1.94	2.59	3.89	5.18	5.37	18.69	6.03	2.15	12.03	323
38	6.83	4.85	10.22	1.33	1.99	2.65	3.98	5.30	5.50	19.16	6.18	2.20	12.33	322
39	6.99	4.97	10.46	1.36	2.03	2.71	4.07	5.42	5.64	19.62	6.32	2.26	12.63	321
40	7.15	5.08	10.69	1.38	2.08	2.77	4.15	5.54	5.77	20.08	6.46	2.31	12.92	320
41	7.31	5.19	10.93	1.41	2.12	2.82	4.24	5.65	5.90	20.53	6.60	2.37	13.22	319
42	7.47	5.30	11.16	1.44	2.16	2.88	4.32	5.76	6.03	20.98	6.74	2.42	13.51	318
43	7.63	5.41	11.38	1.47	2.20	2.94	4.40	5.87	6.16	21.43	6.88	2.48	13.80	317
44	7.79	5.52	11.60	1.50	2.24	2.99	4.49	5.98	6.29	21.87	7.02	2.53	14.08	316
45	7.94	5.63	11.82	1.52	2.28	3.04	4.57	6.09	6.42	22.31	7.15	2.59	14.37	315

NOTE.—For  $L_2$  and  $M_1$  see Table 13; for  $2SM$  and  $MSf$ , take  $u$  of  $M_2$  with sign reversed; for  $P_1$ ,  $R_2$ ,  $S_1$ ,  $S_2$ ,  $S_3$ ,  $S_4$ ,  $T_2$ ,  $Mm$ ,  $Sa$ , and  $Ssa$ , take  $u=0$ .

Table 11.—Values of  $u$  for equilibrium arguments—Continued[Use sign at head of column when  $N$  is between 0 and  $180^\circ$ , reverse sign when  $N$  is between  $180$  and  $360^\circ$ ]

$N$	$J_1$	$K_1$	$K_2$	$M_2, N_2$ $2N_1, M_S$ $\lambda, \mu, \nu$	$M_3$	$M_4, MN$	$M_6$	$M_8$	$O_1, Q_1$ $2Q_1, \rho_1$	$OO$	$MK$	$2MK$	$Mf$	$N$
o	o	o	o	o	o	o	o	o	o	o	o	o	o	o
45	7.94	5.63	11.82	1.52	2.28	3.04	4.57	6.09	6.42	22.31	7.15	2.59	14.37	315
46	8.10	5.74	12.04	1.55	2.32	3.10	4.64	6.19	6.55	22.75	7.28	2.64	14.65	314
47	8.25	5.84	12.26	1.57	2.36	3.15	4.72	6.30	6.68	23.18	7.41	2.69	14.93	313
48	8.40	5.95	12.47	1.60	2.40	3.20	4.80	6.40	6.80	23.60	7.54	2.75	15.20	312
49	8.55	6.05	12.68	1.62	2.44	3.25	4.87	6.50	6.92	24.02	7.67	2.80	15.47	311
50	8.70	6.15	12.88	1.65	2.47	3.30	4.94	6.59	7.05	24.44	7.80	2.85	15.74	310
51	8.84	6.25	13.08	1.67	2.51	3.34	5.01	6.68	7.17	24.85	7.92	2.91	16.01	309
52	8.99	6.35	13.28	1.69	2.54	3.39	5.08	6.78	7.29	25.26	8.04	2.96	16.28	308
53	9.13	6.45	13.48	1.72	2.58	3.44	5.15	6.87	7.41	25.66	8.17	3.01	16.54	307
54	9.27	6.54	13.67	1.74	2.61	3.48	5.22	6.96	7.53	26.06	8.28	3.06	16.80	306
55	9.41	6.64	13.86	1.76	2.64	3.52	5.28	7.04	7.65	26.46	8.40	3.12	17.05	305
56	9.54	6.73	14.05	1.78	2.67	3.56	5.34	7.12	7.76	26.85	8.51	3.17	17.30	304
57	9.68	6.82	14.23	1.80	2.70	3.60	5.40	7.20	7.88	27.23	8.62	3.22	17.55	303
58	9.81	6.91	14.40	1.82	2.73	3.64	5.46	7.28	7.99	27.61	8.73	3.27	17.80	302
59	9.94	7.00	14.58	1.84	2.76	3.68	5.52	7.36	8.10	27.98	8.84	3.32	18.04	301
60	10.07	7.09	14.75	1.86	2.79	3.72	5.58	7.44	8.21	28.34	8.95	3.37	18.28	300
61	10.19	7.17	14.92	1.88	2.82	3.76	5.63	7.51	8.32	28.70	9.05	3.42	18.51	299
62	10.32	7.26	15.08	1.90	2.84	3.79	5.69	7.58	8.42	29.06	9.15	3.46	18.74	298
63	10.44	7.34	15.24	1.91	2.87	3.82	5.74	7.65	8.53	29.41	9.25	3.51	18.97	297
64	10.56	7.42	15.39	1.93	2.89	3.86	5.78	7.71	8.63	29.75	9.35	3.56	19.19	296
65	10.68	7.49	15.54	1.94	2.92	3.89	5.83	7.78	8.73	30.09	9.44	3.61	19.41	295
66	10.79	7.57	15.69	1.96	2.94	3.92	5.88	7.84	8.83	30.42	9.53	3.65	19.63	294
67	10.91	7.64	15.83	1.98	2.96	3.95	5.93	7.90	8.93	30.74	9.62	3.69	19.84	293
68	11.02	7.72	15.96	1.99	2.98	3.98	5.97	7.96	9.03	31.06	9.71	3.74	20.04	292
69	11.12	7.79	16.10	2.00	3.00	4.00	6.01	8.01	9.12	31.37	9.79	3.78	20.25	291
70	11.23	7.86	16.23	2.02	3.02	4.03	6.05	8.06	9.22	31.68	9.87	3.83	20.45	290
71	11.33	7.92	16.35	2.03	3.04	4.06	6.08	8.11	9.31	31.98	9.95	3.87	20.64	289
72	11.43	7.99	16.47	2.04	3.06	4.08	6.11	8.15	9.40	32.27	10.03	3.91	20.83	288
73	11.53	8.05	16.58	2.05	3.08	4.10	6.15	8.20	9.48	32.55	10.10	3.95	21.01	287
74	11.63	8.11	16.69	2.06	3.09	4.12	6.18	8.24	9.57	32.82	10.17	3.99	21.20	286
75	11.72	8.17	16.80	2.07	3.10	4.14	6.21	8.28	9.65	33.09	10.24	4.03	21.37	285
76	11.81	8.23	16.90	2.08	3.12	4.16	6.24	8.32	9.73	33.35	10.31	4.07	21.54	284
77	11.90	8.28	17.00	2.09	3.13	4.18	6.26	8.35	9.81	33.60	10.37	4.11	21.71	283
78	11.98	8.34	17.09	2.10	3.14	4.19	6.29	8.38	9.88	33.85	10.43	4.15	21.87	282
79	12.06	8.39	17.17	2.10	3.15	4.20	6.31	8.41	9.96	34.09	10.49	4.18	22.02	281
80	12.14	8.44	17.25	2.11	3.16	4.22	6.32	8.43	10.03	34.31	10.54	4.22	22.17	280
81	12.22	8.48	17.33	2.11	3.17	4.23	6.34	8.46	10.10	34.53	10.60	4.25	22.32	279
82	12.29	8.53	17.40	2.12	3.18	4.24	6.36	8.48	10.17	34.74	10.65	4.29	22.46	278
83	12.36	8.57	17.46	2.12	3.19	4.25	6.37	8.50	10.23	34.95	10.69	4.32	22.59	277
84	12.42	8.61	17.52	2.13	3.19	4.26	6.38	8.51	10.30	35.14	10.73	4.35	22.72	276
85	12.49	8.64	17.58	2.13	3.20	4.26	6.39	8.52	10.36	35.33	10.77	4.38	22.84	275
86	12.55	8.68	17.63	2.13	3.20	4.27	6.40	8.53	10.41	35.50	10.81	4.41	22.96	274
87	12.60	8.71	17.67	2.14	3.20	4.27	6.41	8.54	10.47	35.67	10.84	4.44	23.07	273
88	12.65	8.74	17.71	2.14	3.20	4.27	6.41	8.54	10.52	35.83	10.87	4.47	23.17	272
89	12.70	8.76	17.74	2.14	3.20	4.27	6.41	8.54	10.57	35.98	10.90	4.49	23.27	271
90	12.75	8.79	17.77	2.14	3.20	4.27	6.41	8.54	10.62	36.12	10.93	4.52	23.37	270

NOTE.—For  $L_1$  and  $M_1$  see Table 13; for  $2SM$  and  $MSf$ , take  $u$  of  $M_2$  with sign reversed; for  $P_1$ ,  $R_1$ ,  $S_1$ ,  $S_2$ ,  $S_3$ ,  $S_4$ ,  $T_2$ ,  $Mm$ ,  $Sa$ , and  $Ssa$ , take  $u=0$ .

Table 11.—Values of  $u$  for equilibrium arguments—Continued[Use sign at head of column when  $N$  is between 0 and  $180^\circ$ , reverse sign when  $N$  is between  $180$  and  $360^\circ$ ]

$N$	$J_1$	$K_1$	$K_2$	$M_2, N_2$ $2N, MS$ $\lambda, \mu, \nu$	$M_3$	$M_4, MN$	$M_5$	$M_8$	$O_1, Q_1$ $2Q, \rho_1$	$OO$	$MK$	$2MK$	$Mf$	$N$
◦	◦	◦	◦	◦	◦	◦	◦	◦	◦	◦	◦	◦	◦	◦
90	12.75	8.79	17.77	2.14	3.20	4.27	6.41	8.54	10.62	36.12	10.93	4.52	23.37	270
91	12.79	8.81	17.79	2.14	3.20	4.27	6.41	8.54	10.66	36.25	10.95	4.54	23.46	269
92	12.83	8.83	17.81	2.13	3.20	4.27	6.40	8.54	10.70	36.37	10.96	4.56	23.54	268
93	12.87	8.85	17.82	2.13	3.20	4.26	6.40	8.53	10.74	36.48	10.98	4.58	23.61	267
94	12.90	8.86	17.83	2.13	3.20	4.26	6.39	8.52	10.77	36.58	10.99	4.60	23.67	266
95	12.93	8.87	17.83	2.13	3.19	4.26	6.38	8.51	10.80	36.67	11.00	4.62	23.73	265
96	12.96	8.88	17.82	2.12	3.19	4.25	6.37	8.49	10.83	36.75	11.01	4.64	23.79	264
97	12.98	8.89	17.81	2.12	3.18	4.24	6.35	8.47	10.86	36.82	11.01	4.65	23.84	263
98	13.00	8.90	17.79	2.11	3.17	4.22	6.34	8.45	10.88	36.87	11.01	4.67	23.88	262
99	13.01	8.90	17.77	2.11	3.16	4.21	6.32	8.43	10.90	36.92	11.00	4.68	23.91	261
100	13.02	8.89	17.74	2.10	3.15	4.20	6.30	8.41	10.92	36.95	11.00	4.69	23.93	260
101	13.03	8.89	17.71	2.09	3.14	4.19	6.28	8.38	10.93	36.98	10.99	4.70	23.95	259
102	13.03	8.88	17.67	2.09	3.13	4.17	6.26	8.34	10.94	36.99	10.97	4.71	23.96	258
103	13.02	8.87	17.62	2.08	3.11	4.15	6.23	8.30	10.95	37.00	10.95	4.72	23.97	257
104	13.02	8.86	17.57	2.07	3.10	4.14	6.20	8.27	10.95	36.99	10.93	4.72	23.97	256
105	13.01	8.84	17.51	2.06	3.09	4.12	6.17	8.23	10.95	36.96	10.90	4.73	23.96	255
106	12.99	8.82	17.45	2.05	3.07	4.10	6.14	8.19	10.94	36.93	10.87	4.73	23.94	254
107	12.97	8.80	17.38	2.04	3.06	4.08	6.11	8.15	10.94	36.89	10.84	4.73	23.91	253
108	12.95	8.78	17.30	2.03	3.04	4.06	6.08	8.11	10.93	36.83	10.81	4.72	23.88	252
109	12.93	8.75	17.22	2.02	3.02	4.03	6.05	8.06	10.91	36.76	10.77	4.72	23.84	251
110	12.90	8.72	17.14	2.00	3.00	4.00	6.01	8.01	10.89	36.67	10.72	4.72	23.79	250
111	12.86	8.69	17.05	1.99	2.98	3.98	5.97	7.96	10.87	36.58	10.68	4.71	23.73	249
112	12.82	8.65	16.95	1.98	2.96	3.95	5.93	7.90	10.84	36.48	10.63	4.70	23.66	248
113	12.77	8.61	16.84	1.96	2.94	3.92	5.88	7.84	10.81	36.36	10.57	4.69	23.59	247
114	12.72	8.57	16.73	1.94	2.92	3.89	5.83	7.78	10.78	36.23	10.51	4.68	23.50	246
115	12.67	8.52	16.62	1.93	2.89	3.86	5.78	7.71	10.74	36.09	10.45	4.67	23.41	245
116	12.61	8.48	16.50	1.91	2.87	3.82	5.74	7.65	10.70	35.93	10.39	4.65	23.31	244
117	12.55	8.43	16.37	1.90	2.84	3.79	5.69	7.58	10.65	35.76	10.32	4.63	23.21	243
118	12.48	8.37	16.24	1.88	2.82	3.76	5.64	7.52	10.60	35.57	10.25	4.61	23.09	242
119	12.41	8.31	16.10	1.86	2.79	3.72	5.59	7.45	10.55	35.37	10.18	4.59	22.96	241
120	12.34	8.25	15.96	1.84	2.77	3.69	5.53	7.38	10.49	35.16	10.10	4.57	22.83	240
121	12.26	8.19	15.81	1.82	2.74	3.65	5.47	7.30	10.43	34.94	10.02	4.54	22.69	239
122	12.17	8.13	15.66	1.80	2.71	3.61	5.41	7.22	10.37	34.70	9.93	4.52	22.54	238
123	12.08	8.06	15.50	1.78	2.68	3.57	5.35	7.14	10.30	34.49	9.84	4.49	22.37	237
124	11.98	7.99	15.33	1.76	2.64	3.52	5.29	7.05	10.22	34.19	9.75	4.46	22.20	236
125	11.88	7.91	15.16	1.74	2.61	3.48	5.22	6.96	10.14	33.91	9.65	4.43	22.03	235
126	11.78	7.83	14.99	1.72	2.58	3.44	5.15	6.87	10.06	33.62	9.55	4.40	21.84	234
127	11.67	7.75	14.81	1.70	2.54	3.39	5.09	6.78	9.97	33.31	9.45	4.36	21.64	233
128	11.56	7.67	14.62	1.67	2.51	3.34	5.02	6.69	9.88	32.99	9.34	4.32	21.44	232
129	11.44	7.58	14.43	1.65	2.48	3.30	4.95	6.60	9.79	32.66	9.23	4.28	21.23	231
130	11.31	7.49	14.23	1.63	2.44	3.26	4.88	6.51	9.69	32.31	9.12	4.23	21.00	230
131	11.18	7.40	14.03	1.60	2.41	3.21	4.81	6.42	9.58	31.95	9.00	4.19	20.76	229
132	11.05	7.30	13.83	1.58	2.37	3.16	4.74	6.32	9.47	31.58	8.88	4.14	20.52	228
133	10.91	7.20	13.62	1.55	2.33	3.11	4.66	6.22	9.36	31.19	8.76	4.09	20.27	227
134	10.77	7.10	13.40	1.53	2.29	3.06	4.58	6.11	9.24	30.79	8.63	4.04	20.01	226
135	10.62	7.00	13.18	1.50	2.25	3.00	4.51	6.01	9.12	30.37	8.50	3.99	19.75	225

NOTE.—For  $L_2$  and  $M_1$  see table 13; for  $2SM$  and  $MSf$ , take  $u$  of  $M_2$  with sign reversed; for  $P_1, R_2, S_1, S_2, S_3, S_4, T_2, M_m, S_m, Ssa$ , take  $u=0$ .

Table 11.—Values of  $u$  for equilibrium arguments—Continued[Use sign at head of column when  $N$  is between 0 and  $180^\circ$ , reverse sign when  $N$  is between  $180$  and  $360^\circ$ ]

$N$	$J_1$	$K_1$	$K_2$	$M_{21}, N_{21}$ $2N, M_S$ $\lambda, \mu, \nu$	$M_3$	$M_4, MN$	$M_5$	$M_8$	$O_1, Q_1$ $2Q, \rho$	$OO$	$MK$	$2MK$	$MF$	$N$
o	o	o	o	o	o	o	o	o	o	o	o	o	o	o
135	10.62	7.00	13.18	1.50	2.25	3.00	4.51	6.01	9.12	30.37	8.50	3.99	19.75	225
136	10.47	6.89	12.96	1.48	2.21	2.95	4.43	5.90	9.00	29.94	8.36	3.94	19.47	224
137	10.31	6.78	12.73	1.45	2.17	2.90	4.34	5.79	8.87	29.49	8.22	3.88	19.18	223
138	10.15	6.66	12.50	1.42	2.13	2.84	4.26	5.68	8.73	29.04	8.08	3.82	18.88	222
139	9.98	6.55	12.26	1.39	2.09	2.78	4.18	5.57	8.59	28.56	7.94	3.76	18.58	221
140	9.81	6.43	12.02	1.36	2.05	2.73	4.09	5.46	8.45	28.08	7.79	3.70	18.26	220
141	9.64	6.31	11.77	1.34	2.00	2.67	4.01	5.34	8.30	27.58	7.64	3.64	17.94	219
142	9.46	6.18	11.52	1.31	1.96	2.61	3.92	5.22	8.15	27.07	7.49	3.57	17.61	218
143	9.27	6.06	11.27	1.28	1.91	2.55	3.83	5.10	8.00	26.54	7.33	3.50	17.27	217
144	9.08	5.93	11.01	1.25	1.87	2.49	3.74	4.98	7.84	26.00	7.17	3.43	16.92	216
145	8.89	5.80	10.74	1.22	1.82	2.43	3.65	4.86	7.67	25.45	7.01	3.36	16.56	215
146	8.69	5.66	10.48	1.19	1.78	2.37	3.56	4.74	7.50	24.89	6.84	3.29	16.20	214
147	8.49	5.52	10.21	1.16	1.73	2.31	3.47	4.62	7.33	24.31	6.68	3.21	15.82	213
148	8.28	5.38	9.94	1.12	1.69	2.25	3.37	4.50	7.16	23.72	6.51	3.14	15.44	212
149	8.07	5.24	9.66	1.09	1.64	2.18	3.28	4.37	6.98	23.12	6.33	3.06	15.05	211
150	7.85	5.10	9.38	1.06	1.59	2.12	3.18	4.24	6.80	22.51	6.16	2.98	14.65	210
151	7.63	4.95	9.10	1.03	1.54	2.06	3.08	4.11	6.61	21.88	5.98	2.90	14.24	209
152	7.41	4.80	8.81	1.00	1.49	1.99	2.99	3.98	6.42	21.24	5.80	2.81	13.83	208
153	7.18	4.65	8.52	0.96	1.44	1.92	2.89	3.85	6.22	20.59	5.61	2.73	13.41	207
154	6.95	4.50	8.23	0.93	1.39	1.86	3.78	3.71	6.03	19.93	5.43	2.64	12.98	206
155	6.72	4.34	7.94	0.90	1.34	1.79	2.69	3.58	5.82	19.26	5.24	2.55	12.54	205
156	6.48	4.19	7.64	0.86	1.29	1.72	2.59	3.45	5.62	18.58	5.05	2.46	12.10	204
157	6.24	4.03	7.34	0.83	1.24	1.66	2.48	3.31	5.41	17.89	4.85	2.37	11.65	203
158	5.99	3.87	7.04	0.79	1.19	1.59	2.38	3.18	5.20	17.19	4.66	2.28	11.19	202
159	5.74	3.70	6.74	0.76	1.14	1.52	2.28	3.04	4.99	16.48	4.46	2.18	10.73	201
160	5.49	3.54	6.43	0.72	1.09	1.45	2.17	2.90	4.77	15.75	4.26	2.09	10.26	200
161	5.24	3.37	6.12	0.69	1.04	1.38	2.07	2.76	4.55	15.02	4.06	1.99	9.79	199
162	4.98	3.20	5.81	0.66	0.98	1.31	1.97	2.62	4.33	14.29	3.86	1.89	9.31	198
163	4.72	3.03	5.50	0.62	0.93	1.24	1.86	2.48	4.10	13.54	3.66	1.79	8.82	197
164	4.46	2.86	5.19	0.58	0.88	1.17	1.75	2.34	3.87	12.78	3.45	1.70	8.33	196
165	4.19	2.69	4.87	0.55	0.82	1.10	1.64	2.19	3.64	12.02	3.24	1.60	7.83	195
166	3.92	2.52	4.55	0.51	0.77	1.02	1.54	2.05	3.41	11.25	3.03	1.49	7.33	194
167	3.65	2.34	4.23	0.48	0.71	0.95	1.43	1.90	3.18	10.48	2.82	1.39	6.83	193
168	3.38	2.17	3.91	0.44	0.66	0.88	1.32	1.76	2.94	9.70	2.61	1.29	6.32	192
169	3.10	1.99	3.59	0.40	0.61	0.81	1.21	1.62	2.70	8.91	2.39	1.18	5.80	191
170	2.83	1.81	3.27	0.37	0.55	0.74	1.10	1.47	2.46	8.12	2.18	1.08	5.29	190
171	2.55	1.63	2.94	0.33	0.50	0.66	1.00	1.33	2.22	7.32	1.97	0.97	4.77	189
172	2.27	1.45	2.62	0.30	0.44	0.59	0.89	1.18	1.97	6.51	1.75	0.86	4.24	188
173	1.99	1.27	2.29	0.26	0.39	0.51	0.77	1.03	1.73	5.71	1.53	0.76	3.72	187
174	1.71	1.09	1.97	0.22	0.33	0.44	0.66	0.88	1.49	4.90	1.31	0.65	3.19	186
175	1.42	0.91	1.64	0.18	0.28	0.37	0.55	0.74	1.24	4.09	1.10	0.54	2.66	185
176	1.14	0.73	1.31	0.15	0.22	0.30	0.44	0.59	0.99	3.27	0.88	0.43	2.13	184
177	0.86	0.55	0.99	0.11	0.17	0.22	0.33	0.44	0.75	2.46	0.66	0.33	1.60	183
178	0.57	0.37	0.66	0.07	0.11	0.14	0.22	0.29	0.50	1.64	0.44	0.22	1.07	182
179	0.29	0.18	0.33	0.04	0.05	0.07	0.11	0.14	0.25	0.82	0.22	0.11	0.53	181
180	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	180

NOTE.—For  $L_2$  and  $M_1$  see Table 13; for  $2SM$  and  $MSf$ , take  $u$  of  $M_1$  with sign reversed; for  $P_1$ ,  $R_2$ ,  $S_1$ ,  $S_2$ ,  $S_3$ ,  $S_4$ ,  $T_2$ ,  $Mm$ ,  $Sa$ , and  $Ssa$ , take  $u=0$ .

Table 12.—*Log factor F corresponding to every tenth of a degree of I*

<i>I</i> Constituent	18.3°	Diff.	18.4°	Diff.	18.5°	Diff.	18.6°	Diff.	18.7°	Diff.	18.8°	Diff.
J <sub>1</sub> -----	0.0827	-21	0.0806	-20	0.0786	-20	0.0766	-20	0.0746	-19	0.0727	-20
K <sub>1</sub> -----	0.0547	-12	0.0535	-12	0.0523	-11	0.0512	-12	0.0500	-12	0.0488	-11
K <sub>2</sub> -----	0.1263	-22	0.1241	-21	0.1220	-21	0.1199	-22	0.1177	-22	0.1155	-21
M <sub>2</sub> , N <sub>2</sub> , 2N-----	9.9839	+2	9.9841	+3	9.9844	+2	9.9846	+3	9.9849	+2	9.9851	+3
M <sub>3</sub> -----	9.9758	+4	9.9762	+4	9.9766	+3	9.9769	+4	9.9773	+4	9.9777	+3
M <sub>4</sub> , MN-----	9.9678	+4	9.9682	+5	9.9687	+5	9.9692	+5	9.9697	+5	9.9702	+5
M <sub>8</sub> -----	9.9516	+8	9.9524	+7	9.9531	+7	9.9538	+8	9.9546	+7	9.9553	+8
M <sub>8</sub> -----	9.9355	+10	9.9365	+10	9.9375	+10	9.9385	+10	9.9395	+10	9.9405	+10
O <sub>1</sub> , Q <sub>1</sub> 2Q, $\rho_1$ -----	0.0939	-22	0.0917	-21	0.0896	-22	0.0874	-21	0.0853	-21	0.0832	-21
OO-----	0.3139	-69	0.3070	-70	0.3000	-69	0.2931	-68	0.2863	-69	0.2794	-68
MK-----	0.0386	-10	0.0376	-9	0.0367	-9	0.0358	-9	0.0349	-10	0.0339	-9
2MK-----	0.0224	-7	0.0217	-6	0.0211	-7	0.0204	-7	0.0197	-7	0.0190	-6
Mf-----	0.2039	-46	0.1993	-45	0.1948	-45	0.1903	-45	0.1858	-45	0.1813	-44
Mm-----	9.9465	+8	9.9473	+8	9.9481	+8	9.9489	+8	9.9497	+8	9.9505	+9
<i>I</i> Constituent	18.9°	Diff.	19.0°	Diff.	19.1°	Diff.	19.2°	Diff.	19.3°	Diff.	19.4°	Diff.
J <sub>1</sub> -----	0.0707	-20	0.0687	-19	0.0668	-19	0.0649	-19	0.0630	-19	0.0611	-19
K <sub>1</sub> -----	0.0477	-12	0.0465	-12	0.0453	-11	0.0442	-11	0.0431	-12	0.0419	-11
K <sub>2</sub> -----	0.1134	-22	0.1112	-22	0.1090	-22	0.1068	-23	0.1045	-22	0.1023	-22
M <sub>2</sub> , N <sub>2</sub> , 2N-----	9.9854	+2	9.9856	+3	9.9859	+2	9.9861	+3	9.9864	+2	9.9866	+3
M <sub>3</sub> -----	9.9780	+4	9.9784	+4	9.9788	+4	9.9792	+4	9.9796	+4	9.9800	+4
M <sub>4</sub> , MN-----	9.9707	+5	9.9712	+5	9.9717	+6	9.9723	+5	9.9728	+5	9.9733	+5
M <sub>8</sub> -----	9.9561	+8	9.9569	+7	9.9576	+8	9.9584	+8	9.9592	+7	9.9599	+8
M <sub>8</sub> -----	9.9415	+10	9.9425	+10	9.9435	+10	9.9445	+10	9.9455	+11	9.9466	+10
O <sub>1</sub> , Q <sub>1</sub> , 2Q, $\rho_1$ -----	0.0811	-21	0.0790	-20	0.0770	-21	0.0749	-20	0.0729	-20	0.0709	-21
OO-----	0.2726	-67	0.2659	-67	0.2592	-67	0.2525	-66	0.2459	-66	0.2393	-66
MK-----	0.0330	-9	0.0321	-9	0.0312	-9	0.0303	-9	0.0294	-9	0.0285	-8
2MK-----	0.0184	-7	0.0177	-6	0.0171	-7	0.0164	-6	0.0158	-6	0.0152	-6
Mf-----	0.1769	-44	0.1725	-44	0.1681	-44	0.1637	-43	0.1594	-43	0.1551	-43
Mm-----	9.9514	+8	9.9522	+8	9.9530	+9	9.9539	+8	9.9547	+9	9.9556	+8
<i>I</i> Constituent	19.5°	Diff.	19.6°	Diff.	19.7°	Diff.	19.8°	Diff.	19.9°	Diff.	20.0°	Diff.
J <sub>1</sub> -----	0.0592	-18	0.0574	-19	0.0555	-18	0.0537	-19	0.0518	-18	0.0500	-18
K <sub>1</sub> -----	0.0408	-12	0.0396	-11	0.0385	-11	0.0374	-12	0.0362	-11	0.0351	-11
K <sub>2</sub> -----	0.1001	-23	0.0978	-22	0.0956	-23	0.0933	-22	0.0911	-23	0.0888	-24
M <sub>2</sub> , N <sub>2</sub> , 2N-----	9.9869	+3	9.9872	+2	9.9874	+3	9.9877	+3	9.9880	+2	9.9882	+3
M <sub>3</sub> -----	9.9804	+3	9.9807	+4	9.9811	+4	9.9815	+4	9.9819	+4	9.9823	+4
M <sub>4</sub> , MN-----	9.9738	+5	9.9743	+6	9.9749	+5	9.9754	+5	9.9759	+5	9.9764	+6
M <sub>8</sub> -----	9.9607	+8	9.9615	+8	9.9623	+8	9.9631	+8	9.9639	+8	9.9647	+8
M <sub>8</sub> -----	9.9476	+11	9.9487	+10	9.9497	+11	9.9508	+10	9.9518	+11	9.9529	+11
O <sub>1</sub> , Q <sub>1</sub> , 2Q, $\rho_1$ -----	0.0688	-20	0.0668	-20	0.0648	-19	0.0629	-20	0.0609	-20	0.0589	-19
OO-----	0.2327	-65	0.2262	-65	0.2197	-65	0.2132	-64	0.2068	-64	0.2004	-64
MK-----	0.0277	-9	0.0268	-9	0.0259	-9	0.0250	-8	0.0242	-9	0.0233	-8
2MK-----	0.0146	-6	0.0140	-7	0.0133	-6	0.0127	-6	0.0121	-6	0.0115	-6
Mf-----	0.1508	-43	0.1465	-42	0.1423	-43	0.1380	-42	0.1338	-41	0.1297	-42
Mm-----	9.9564	+9	9.9573	+8	9.9581	+9	9.9590	+9	9.9599	+9	9.9608	+9

\*Log  $F$  of  $\lambda_2$ ,  $\mu_2$ ,  $\nu_2$ , MS, 2SM, and MSf are each equal to log  $F$  of M<sub>2</sub>.Log  $F$  of P<sub>1</sub>, R<sub>2</sub>, S<sub>1</sub>, S<sub>2</sub>, S<sub>4</sub>, S<sub>6</sub>, T<sub>2</sub>, Sa, and Ssa are each zero.For log  $F$  of L<sub>2</sub> and M<sub>1</sub> see Table 13.

Table 12.—*Log factor F corresponding to every tenth of a degree of I—Con.*

<i>I</i> Constituent	20.1°	Diff.	20.2°	Diff.	20.3°	Diff.	20.4°	Diff.	20.5°	Diff.	20.6°	Diff.
J <sub>1</sub> -----	0.0482	-18	0.0464	-17	0.0447	-18	0.0429	-18	0.0411	-17	0.0394	-17
K <sub>1</sub> -----	0.0340	-11	0.0329	-11	0.0318	-11	0.0307	-11	0.0296	-11	0.0285	-11
K <sub>2</sub> -----	0.0864	-23	0.0841	-23	0.0818	-23	0.0795	-24	0.0771	-23	0.0748	-23
M <sub>2</sub> * <sub>1</sub> , N <sub>2</sub> , 2N <sub>1</sub> -----	9.9885	+3	9.9888	+2	9.9890	+3	9.9893	+3	9.9896	+3	9.9899	+2
M <sub>3</sub> -----	9.9827	+4	9.9831	+4	9.9835	+5	9.9840	+4	9.9844	+4	9.9848	+4
M <sub>4</sub> , MN-----	9.9770	+5	9.9775	+6	9.9781	+5	9.9786	+6	9.9792	+5	9.9797	+6
M <sub>6</sub> -----	9.9655	+8	9.9663	+8	9.9671	+8	9.9679	+8	9.9687	+9	9.9696	+8
M <sub>8</sub> -----	9.9540	+10	9.9550	+11	9.9561	+11	9.9572	+11	9.9583	+11	9.9594	+11
O <sub>1</sub> , Q <sub>1</sub> , 2Q <sub>1</sub> , p <sub>1</sub> -----	0.0570	-19	0.0551	-20	0.0531	-19	0.0512	-19	0.0493	-18	0.0475	-19
OO-----	0.1940	-63	0.1877	-63	0.1814	-63	0.1751	-62	0.1689	-62	0.1627	-62
MK-----	0.0225	-8	0.0217	-9	0.0208	-8	0.0200	-9	0.0191	-8	0.0183	-8
2MK-----	0.0109	-5	0.0104	-6	0.0098	-5	0.0093	-6	0.0087	-5	0.0082	-6
Mf-----	0.1255	-41	0.1214	-41	0.1173	-41	0.1132	-41	0.1091	-40	0.1051	-41
Mm-----	9.9617	+9	9.9626	+9	9.9635	+9	9.9644	+9	9.9653	+9	9.9662	+9
<i>I</i> Constituent	20.7°	Diff.	20.8°	Diff.	20.9°	Diff.	21.0°	Diff.	21.1°	Diff.	21.2°	Diff.
J <sub>1</sub> -----	0.0377	-17	0.0360	-17	0.0343	-17	0.0326	-17	0.0309	-17	0.0292	-16
K <sub>1</sub> -----	0.0274	-11	0.0263	-11	0.0252	-11	0.0241	-11	0.0230	-11	0.0219	-10
K <sub>2</sub> -----	0.0725	-24	0.0701	-23	0.0678	-24	0.0654	-24	0.0630	-23	0.0607	-24
M <sub>2</sub> * <sub>1</sub> , N <sub>2</sub> , 2N <sub>1</sub> -----	9.9901	+3	9.9904	+3	9.9907	+3	9.9910	+2	9.9912	+3	9.9915	+3
M <sub>3</sub> -----	9.9852	+4	9.9856	+4	9.9860	+4	9.9864	+5	9.9869	+4	9.9873	+4
M <sub>4</sub> , MN-----	9.9803	+5	9.9808	+6	9.9814	+5	9.9819	+6	9.9825	+6	9.9831	+5
M <sub>6</sub> -----	9.9704	+8	9.9712	+8	9.9720	+9	9.9729	+8	9.9737	+9	9.9746	+8
M <sub>8</sub> -----	9.9605	+11	9.9616	+11	9.9627	+12	9.9639	+11	9.9650	+11	9.9661	+12
O <sub>1</sub> , Q <sub>1</sub> , 2Q <sub>1</sub> , p <sub>1</sub> -----	0.0456	-19	0.0437	-18	0.0419	-19	0.0400	-18	0.0382	-18	0.0364	-18
OO-----	0.1565	-61	0.1504	-61	0.1443	-61	0.1382	-61	0.1321	-60	0.1261	-60
MK-----	0.0175	-8	0.0167	-8	0.0159	-8	0.0151	-8	0.0143	-8	0.0135	-8
2MK-----	0.0076	-5	0.0071	-6	0.0065	-5	0.0060	-5	0.0055	-5	0.0050	-5
Mf-----	0.1010	-40	0.0970	-39	0.0931	-40	0.0891	-39	0.0852	-40	0.0812	-39
Mm-----	9.9671	+9	9.9680	+10	9.9690	+9	9.9699	+10	9.9709	+9	9.9718	+10
<i>I</i> Constituent	21.3°	Diff.	21.4°	Diff.	21.5°	Diff.	21.6°	Diff.	21.7°	Diff.	21.8°	Diff.
J <sub>1</sub> -----	0.0276	-17	0.0259	-16	0.0243	-16	0.0227	-16	0.0211	-16	0.0195	-16
K <sub>1</sub> -----	0.0209	-11	0.0198	-11	0.0187	-10	0.0177	-11	0.0166	-10	0.0156	-11
K <sub>2</sub> -----	0.0583	-24	0.0559	-25	0.0534	-24	0.0510	-24	0.0486	-24	0.0462	-24
M <sub>2</sub> * <sub>1</sub> , N <sub>2</sub> , 2N <sub>1</sub> -----	9.9918	+3	9.9921	+3	9.9924	+3	9.9927	+3	9.9930	+3	9.9933	+3
M <sub>3</sub> -----	9.9877	+5	9.9882	+4	9.9886	+4	9.9890	+4	9.9894	+5	9.9899	+4
M <sub>4</sub> , MN-----	9.9836	+6	9.9842	+6	9.9848	+6	9.9854	+5	9.9859	+6	9.9865	+6
M <sub>6</sub> -----	9.9754	+9	9.9763	+9	9.9772	+8	9.9780	+9	9.9789	+9	9.9798	+8
M <sub>8</sub> -----	9.9673	+11	9.9684	+12	9.9696	+11	9.9707	+12	9.9719	+11	9.9730	+12
O <sub>1</sub> , Q <sub>1</sub> , 2Q <sub>1</sub> , p <sub>1</sub> -----	0.0346	-18	0.0328	-18	0.0310	-18	0.0292	-17	0.0275	-18	0.0257	-17
OO-----	0.1201	-60	0.1141	-59	0.1082	-59	0.1023	-59	0.0964	-58	0.0906	-58
MK-----	0.0127	-8	0.0119	-8	0.0111	-8	0.0103	-7	0.0096	-8	0.0088	-7
2MK-----	0.0045	-5	0.0040	-5	0.0035	-5	0.0030	-5	0.0025	-4	0.0021	-5
Mf-----	0.0773	-38	0.0735	-39	0.0696	-38	0.0658	-39	0.0619	-38	0.0581	-37
Mm-----	9.9728	+9	9.9737	+10	9.9747	+10	9.9757	+10	9.9767	+9	9.9776	+10

\* Log *F* of  $\lambda_2$ ,  $\mu_2$ ,  $\nu_2$ , SM, 2SM, and MSF are each equal to log *F* of M<sub>2</sub>.Log *F* of P<sub>1</sub>, R<sub>2</sub>, S<sub>1</sub>, S<sub>2</sub>, S<sub>4</sub>, S<sub>6</sub>, T<sub>2</sub>, Sa, and Ssa are each zero.For log *F* of L<sub>2</sub> and M<sub>1</sub> see Table 13.

Table 12.—*Log factor F corresponding to every tenth of a degree of I—Con.*

<i>I</i> Constituent	21.9°	Diff.	22.0°	Diff.	22.1°	Diff.	22.2°	Diff.	22.3°	Diff.	22.4°	Diff.
J <sub>1</sub> -----	0.0179	-16	0.0163	-15	0.0148	-16	0.0132	-15	0.0117	-16	0.0101	-15
K <sub>1</sub> -----	0.0145	-10	0.0135	-11	0.0124	-10	0.0114	-11	0.0103	-10	0.0093	-10
K <sub>2</sub> -----	0.0438	-24	0.0414	-24	0.0390	-25	0.0365	-24	0.0341	-24	0.0317	-24
M <sub>2</sub> * , N <sub>2</sub> , 2N-----	9.9936	+2	9.9938	+3	9.9941	+3	9.9944	+3	9.9947	+3	9.9950	+3
M <sub>3</sub> -----	9.9903	+5	9.9908	+4	9.9912	+5	9.9917	+4	9.9921	+5	9.9926	+4
M <sub>4</sub> MN-----	9.9871	+6	9.9877	+6	9.9883	+6	9.9889	+6	9.9895	+6	9.9901	+6
M <sub>6</sub> -----	9.9806	+9	9.9815	+9	9.9824	+9	9.9833	+9	9.9842	+9	9.9851	+9
M <sub>8</sub> -----	9.9742	+12	9.9754	+12	9.9766	+11	9.9777	+12	9.9789	+12	9.9801	+12
O <sub>1</sub> , Q <sub>1</sub> , 2Q, p <sub>1</sub> -----	0.0240	-18	0.0222	-17	0.0205	-17	0.0188	-17	0.0171	-17	0.0154	-17
OO-----	0.0848	-58	0.0790	-58	0.0732	-57	0.0675	-57	0.0618	-57	0.0561	-57
MK-----	0.0081	-8	0.0073	-8	0.0065	-7	0.0058	-7	0.0051	-8	0.0043	-7
2MK-----	0.0016	-5	0.0011	-4	0.0007	-5	0.0002	-4	9.9998	-4	9.9994	-4
Mf-----	0.0544	-38	0.0506	-37	0.0469	-38	0.0431	-37	0.0394	-37	0.0357	-36
Mm-----	9.9786	+10	9.9796	+10	9.9806	+10	9.9816	+11	9.9827	+10	9.9837	+10
<i>I</i> Constituent	22.5°	Diff.	22.6°	Diff.	22.7°	Diff.	22.8°	Diff.	22.9°	Diff.	23.0°	Diff.
J <sub>1</sub> -----	0.0086	-15	0.0071	-15	0.0056	-15	0.0041	-15	0.0026	-14	0.0012	-15
K <sub>1</sub> -----	0.0083	-10	0.0073	-10	0.0063	-11	0.0052	-10	0.0042	-10	0.0032	-10
K <sub>2</sub> -----	0.0293	-25	0.0268	-24	0.0244	-25	0.0219	-25	0.0194	-24	0.0170	-25
M <sub>2</sub> * , N <sub>2</sub> , 2N-----	9.9953	+3	9.9956	+3	9.9959	+3	9.9962	+4	9.9966	+3	9.9969	+3
M <sub>3</sub> -----	9.9930	+5	9.9935	+4	9.9939	+5	9.9944	+4	9.9948	+5	9.9953	+5
M <sub>4</sub> , MN-----	9.9907	+6	9.9913	+6	9.9919	+6	9.9925	+6	9.9931	+6	9.9937	+6
M <sub>6</sub> -----	9.9860	+9	9.9869	+9	9.9878	+9	9.9887	+10	9.9897	+9	9.9906	+9
M <sub>8</sub> -----	9.9813	+12	9.9825	+13	9.9838	+12	9.9850	+12	9.9862	+12	9.9874	+13
O <sub>1</sub> , Q <sub>1</sub> , 2Q, p <sub>1</sub> -----	0.0137	-17	0.0120	-16	0.0104	-17	0.0087	-16	0.0071	-17	0.0054	-16
OO-----	0.0504	-56	0.0448	-56	0.0392	-56	0.0336	-55	0.0281	-55	0.0226	-55
MK-----	0.0036	-7	0.0029	-7	0.0022	-7	0.0015	-7	0.0008	-7	0.0001	-7
2MK-----	9.9990	-5	9.9985	-4	9.9981	-4	9.9977	-4	9.9973	-4	9.9969	-3
Mf-----	0.0321	-37	0.0284	-36	0.0248	-36	0.0212	-36	0.0176	-36	0.0140	-36
Mm-----	9.9847	+10	9.9857	+11	9.9868	+10	9.9878	+11	9.9889	+10	9.9899	+11
<i>I</i> Constituent	23.1°	Diff.	23.2°	Diff.	23.3°	Diff.	23.4°	Diff.	23.5°	Diff.	23.6°	Diff.
J <sub>1</sub> -----	9.9997	-15	9.9982	-14	9.9968	-14	9.9954	-14	9.9940	-14	9.9926	-14
K <sub>1</sub> -----	0.0022	-10	0.0012	-10	0.0002	-10	9.9992	-10	9.9982	-9	9.9973	-10
K <sub>2</sub> -----	0.0145	-24	0.0121	-25	0.0096	-24	0.0072	-25	0.0047	-25	0.0022	-24
M <sub>2</sub> * , N <sub>2</sub> , 2N-----	9.9972	+3	9.9975	+3	9.9978	+3	9.9981	+3	9.9984	+3	9.9987	+4
M <sub>3</sub> -----	9.9958	+5	9.9963	+4	9.9967	+5	9.9972	+4	9.9976	+5	9.9981	+5
M <sub>4</sub> , MN-----	9.9943	+7	9.9950	+6	9.9956	+6	9.9962	+6	9.9968	+7	9.9975	+6
M <sub>6</sub> -----	9.9915	+9	9.9924	+10	9.9934	+9	9.9943	+10	9.9953	+9	9.9962	+10
M <sub>8</sub> -----	9.9887	+12	9.9899	+13	9.9912	+12	9.9924	+13	9.9937	+12	9.9949	+13
O <sub>1</sub> , Q <sub>1</sub> , 2Q, p <sub>1</sub> -----	0.0038	-16	0.0022	-16	0.0006	-16	9.9990	-16	9.9974	-16	9.9958	-16
OO-----	0.0171	-55	0.0116	-54	0.0062	-55	0.0007	-54	9.9953	-53	9.9900	-54
MK-----	9.9994	-7	9.9987	-7	9.9980	-6	9.9974	-7	9.9967	-7	9.9960	-6
2MK-----	9.9966	-4	9.9962	-4	9.9958	-3	9.9955	-4	9.9951	-3	9.9948	-4
Mf-----	0.0104	-35	0.0069	-35	0.0034	-35	9.9999	-35	9.9964	-35	9.9929	-35
Mm-----	9.9910	+11	9.9921	+10	9.9931	+11	9.9942	+11	9.9953	+11	9.9964	+11

\*Log *F* of  $\lambda_2 \mu_2, v_2, MS, 2SM$  and  $MSf$  are each equal to log *F* of  $M_2$ .Log *F* of  $P_1, R_2 S_1, S_2, S_4, S_6, T_2 S_4$  and  $S_8$  are each zero.For log *F* of  $L_2$  and  $M_1$  see Table 13.

Table 12.—*Log factor F corresponding to every tenth of a degree of I—Con.*

<i>I</i> Constituent	23.7°	Diff.	23.8°	Diff.	23.9°	Diff.	24.0°	Diff.	24.1°	Diff.	24.2°	Diff.
J <sub>1</sub> -----	9.9912	-14	9.9898	-14	9.9884	-14	9.9870	-13	9.9857	-14	9.9843	-13
K <sub>1</sub> -----	9.9963	-9	9.9954	-10	9.9944	-10	9.9934	-10	9.9924	-9	9.9915	-10
K <sub>2</sub> -----	9.9998	-25	9.9973	-25	9.9948	-24	9.9924	-25	9.9899	-25	9.9874	-24
M <sub>2</sub> * <sup>a</sup> , N <sub>2</sub> , 2N-----	9.9991	+3	9.9994	+3	9.9997	+3	0.0000	+3	0.0003	+4	0.0007	+3
M <sub>3</sub> -----	9.9986	+5	9.9991	+4	9.9995	+5	0.0000	+5	0.0005	+5	0.0010	+5
M <sub>4</sub> , MN-----	9.9981	+6	9.9987	+7	9.9994	+6	0.0000	+7	0.0007	+6	0.0013	+7
M <sub>6</sub> -----	9.9972	+9	9.9981	+10	9.9991	+10	0.0001	+9	0.0010	+10	0.0020	+10
M <sub>8</sub> -----	9.9962	+13	9.9975	+13	9.9988	+13	0.0001	+12	0.0013	+13	0.0026	+13
O <sub>1</sub> , Q <sub>1</sub> , 2Q, p <sub>1</sub> -----	9.9942	-15	9.9927	-16	9.9911	-15	9.9896	-16	9.9880	-15	9.9865	-15
OO-----	9.9846	-53	9.9793	-53	9.9740	-53	9.9687	-53	9.9634	-52	9.9582	-52
MK-----	9.9954	-7	9.9947	-6	9.9941	-7	9.9934	-7	9.9927	-6	9.9921	-6
2MK-----	9.9944	-3	9.9941	-4	9.9937	-3	9.9934	-3	9.9931	-3	9.9928	-3
Mf-----	9.9894	-34	9.9860	-35	9.9825	-34	9.9791	-34	9.9757	-33	9.9724	-34
Mm-----	9.9975	+11	9.9986	+11	9.9997	+12	0.0009	+11	0.0020	+11	0.0031	+12
<i>I</i> Cotstituent	24.3°	Diff.	24.4°	Diff.	24.5°	Diff.	24.6°	Diff.	24.7°	Diff.	24.8°	Diff.
J <sub>1</sub> -----	9.9830	-14	9.9816	-13	9.9803	-13	9.9790	-13	9.9777	-13	9.9764	-13
K <sub>1</sub> -----	9.9905	-9	9.9896	-9	9.9887	-10	9.9877	-9	9.9868	-10	9.9858	-9
K <sub>2</sub> -----	9.9850	-25	9.9825	-25	9.9800	-24	9.9776	-25	9.9751	-25	9.9726	-25
M <sub>2</sub> * <sup>a</sup> , N <sub>2</sub> , 2N-----	0.0010	+3	0.0013	+3	0.0016	+4	0.0020	+3	0.0023	+3	0.0026	+4
M <sub>3</sub> -----	0.0015	+5	0.0020	+5	0.0025	+5	0.0030	+5	0.0035	+5	0.0040	+5
M <sub>4</sub> , MN-----	0.0020	+6	0.0026	+7	0.0033	+6	0.0039	+7	0.0046	+7	0.0053	+6
M <sub>6</sub> -----	0.0030	+9	0.0039	+10	0.0049	+10	0.0059	+10	0.0069	+10	0.0079	+10
M <sub>8</sub> -----	0.0039	+14	0.0053	+13	0.0066	+13	0.0079	+13	0.0092	+13	0.0105	+14
O <sub>1</sub> , Q <sub>1</sub> , 2Q, p <sub>1</sub> -----	9.9850	-15	9.9835	-15	9.9820	-15	9.9805	-15	9.9790	-15	9.9775	-15
OO-----	9.9530	-52	9.9478	-52	9.9426	-51	9.9375	-51	9.9324	-51	9.9273	-51
MK-----	9.9915	-6	9.9909	-6	9.9903	-6	9.9897	-6	9.9891	-6	9.9885	-6
2MK-----	9.9925	-3	9.9922	-3	9.9919	-3	9.9916	-2	9.9914	-3	9.9911	-2
Mf-----	9.9690	-34	9.9656	-33	9.9623	-33	9.9590	-33	9.9557	-33	9.9524	-33
Mm-----	0.0043	+11	0.0054	+12	0.0066	+11	0.0077	+12	0.0089	+12	0.0101	+11
<i>I</i> Constituent	24.9°	Diff.	25.0°	Diff.	25.1°	Diff.	25.2°	Diff.	25.3°	Diff.	25.4°	Diff.
J <sub>1</sub> -----	9.9751	-13	9.9738	-12	9.9726	-13	9.9713	-12	9.9701	-13	9.9688	-12
K <sub>1</sub> -----	9.9849	-9	9.9840	-9	9.9831	-9	9.9822	-10	9.9812	-9	9.9803	-9
K <sub>2</sub> -----	9.9701	-24	9.9677	-25	9.9652	-24	9.9628	-25	9.9603	-24	9.9579	-25
M <sub>2</sub> * <sup>a</sup> , N <sub>2</sub> , 2N-----	0.0030	+3	0.0033	+3	0.0036	+4	0.0040	+3	0.0043	+4	0.0047	+3
M <sub>3</sub> -----	0.0045	+5	0.0050	+5	0.0055	+5	0.0060	+5	0.0065	+5	0.0070	+5
M <sub>4</sub> , MN-----	0.0059	+7	0.0066	+7	0.0073	+7	0.0080	+6	0.0086	+7	0.0093	+7
M <sub>6</sub> -----	0.0089	+10	0.0099	+10	0.0109	+10	0.0119	+11	0.0130	+10	0.0140	+10
M <sub>8</sub> -----	0.0119	+13	0.0132	+14	0.0146	+13	0.0159	+14	0.0173	+13	0.0186	+14
O <sub>1</sub> , Q <sub>1</sub> , 2Q, p <sub>1</sub> -----	9.9760	-14	9.9746	-15	9.9731	-14	9.9717	-15	9.9702	-14	9.9688	-14
OO-----	9.9222	-51	9.9171	-50	9.9121	-50	9.9071	-50	9.9021	-50	9.8971	-49
MK-----	9.9879	-6	9.9873	-6	9.9867	-6	9.9861	-5	9.9856	-6	9.9850	-6
2MK-----	9.9909	-3	9.9906	-2	9.9904	-3	9.9901	-2	9.9899	-2	9.9897	-3
Mf-----	9.9491	-32	9.9459	-33	9.9426	-32	9.9394	-32	9.9362	-32	9.9330	-32
Mm-----	0.0112	+12	0.0124	+12	0.0136	+12	0.0148	+12	0.0160	+12	0.0172	+13

<sup>a</sup>Log F of  $\lambda_2$ ,  $\mu_2$ ,  $\nu_2$ , MS, 2SM, and MSf are each equal to log F of M<sub>2</sub>.Log F of P<sub>1</sub>, R<sub>2</sub>, S<sub>1</sub>, S<sub>2</sub>, S<sub>4</sub>, S<sub>5</sub>, T<sub>2</sub>, Sa, and Ssa are each zero.For log F of L<sub>2</sub> and M<sub>1</sub> see Table 13.

Table 12.—*Log factor F corresponding to every tenth of a degree of I—Con.*

<i>I</i> Constituent	25.5°	Diff.	25.6°	Diff.	25.7°	Diff.	25.8°	Diff.	25.9°	Diff.	26.0°	Diff.
J <sub>1</sub> -----	9.9676	-12	9.9664	-12	9.9652	-13	9.9639	-12	9.9627	-11	9.9616	-12
K <sub>1</sub> -----	9.9794	-9	9.9785	-9	9.9776	-8	9.9768	-9	9.9759	-9	9.9750	-9
K <sub>2</sub> -----	9.9554	-25	9.9529	-25	9.9504	-24	9.9480	-25	9.9455	-24	9.9431	-25
M <sub>2</sub> * N <sub>2</sub> 2N-----	0.0050	+3	0.0053	+4	0.0057	+3	0.0060	+4	0.0064	+3	0.0067	+4
M <sub>3</sub> -----	0.0075	+5	0.0080	+5	0.0085	+6	0.0091	+5	0.0096	+5	0.0101	+5
M <sub>4</sub> MN-----	0.0100	+7	0.0107	+7	0.0114	+7	0.0121	+7	0.0128	+7	0.0135	+7
M <sub>6</sub> -----	0.0150	+10	0.0160	+11	0.0171	+10	0.0181	+11	0.0192	+10	0.0202	+11
M <sub>8</sub> -----	0.0200	+14	0.0214	+14	0.0228	+13	0.0241	+14	0.0255	+14	0.0269	+14
O <sub>1</sub> Q <sub>1</sub> 2Q <sub>1</sub> p <sub>1</sub> -----	9.9674	-14	9.9660	-14	9.9646	-14	9.9632	-14	9.9618	-14	9.9604	-14
OO-----	9.8922	-49	9.8873	-49	9.8824	-49	9.8775	-49	9.8726	-49	9.8677	-48
MK-----	9.9844	-5	9.9839	-6	9.9833	-5	9.9828	-5	9.9823	-6	9.9817	-5
2MK-----	9.9894	-2	9.9892	-2	9.9890	-2	9.9888	-2	9.9886	-1	9.9885	-2
Mf-----	9.9298	-32	9.9266	-31	9.9235	-32	9.9203	-31	9.9172	-31	9.9141	-31
Mm-----	0.0185	+12	0.0197	+12	0.0209	+13	0.0222	+12	0.0234	+13	0.0247	+12
<i>I</i> Constituent	26.1°	Diff.	26.2°	Diff.	26.3°	Diff.	26.4°	Diff.	26.5°	Diff.	26.6°	Diff.
J <sub>1</sub> -----	9.9604	-12	9.9592	-12	9.9580	-11	9.9569	-12	9.9557	-11	9.9546	-11
K <sub>1</sub> -----	9.9741	-9	9.9732	-8	9.9724	-9	9.9715	-9	9.9706	-8	9.9698	-9
K <sub>2</sub> -----	9.9406	-24	9.9382	-25	9.9357	-24	9.9333	-25	9.9308	-24	9.9284	-24
M <sub>2</sub> * N <sub>2</sub> 2N-----	0.0071	+3	0.0074	+4	0.0078	+3	0.0081	+4	0.0085	+4	0.0089	+3
M <sub>3</sub> -----	0.0106	+6	0.0112	+5	0.0117	+5	0.0122	+6	0.0128	+5	0.0133	+5
M <sub>4</sub> MN-----	0.0142	+7	0.0149	+7	0.0156	+7	0.0163	+7	0.0170	+7	0.0177	+7
M <sub>6</sub> -----	0.0213	+10	0.0223	+11	0.0234	+10	0.0244	+11	0.0255	+11	0.0266	+11
M <sub>8</sub> -----	0.0283	+15	0.0298	+14	0.0312	+14	0.0326	+14	0.0340	+14	0.0354	+15
O <sub>1</sub> Q <sub>1</sub> 2Q <sub>1</sub> p <sub>1</sub> -----	9.9590	-13	9.9577	-14	9.9563	-14	9.9549	-13	9.9536	-13	9.9523	-14
OO-----	9.8629	-48	9.8581	-48	9.8533	-47	9.8486	-48	9.8438	-47	9.8391	-47
MK-----	9.9812	-5	9.9807	-5	9.9802	-5	9.9797	-5	9.9792	-5	9.9787	-5
2MK-----	9.9883	-2	9.9881	-1	9.9880	-2	9.9878	-1	9.9877	-2	9.9875	-1
Mf-----	9.9110	-31	9.9079	-31	9.9048	-31	9.9017	-30	9.8987	-30	9.8957	-31
Mm-----	0.0259	+13	0.0272	+13	0.0285	+13	0.0298	+12	0.0310	+13	0.0323	+13
<i>I</i> Constituent	26.7°	Diff.	26.8°	Diff.	26.9°	Diff.	27.0°	Diff.	27.1°	Diff.	27.2°	Diff.
J <sub>1</sub> -----	9.9535	-11	9.9524	-12	9.9512	-11	9.9501	-11	9.9490	-11	9.9479	-10
K <sub>1</sub> -----	9.9689	-8	9.9681	-9	9.9672	-8	9.9664	-8	9.9656	-9	9.9647	-8
K <sub>2</sub> -----	9.9260	-25	9.9235	-24	9.9211	-24	9.9187	-25	9.9162	-24	9.9138	-24
M <sub>2</sub> * N <sub>2</sub> 2N-----	0.0092	+4	0.0096	+3	0.0099	+4	0.0103	+4	0.0107	+3	0.0110	+4
M <sub>3</sub> -----	0.0138	+6	0.0144	+5	0.0149	+6	0.0155	+5	0.0160	+6	0.0166	+5
M <sub>4</sub> MN-----	0.0184	+8	0.0192	+7	0.0199	+7	0.0206	+7	0.0213	+8	0.0221	+7
M <sub>6</sub> -----	0.0277	+10	0.0287	+11	0.0298	+11	0.0309	+11	0.0320	+11	0.0331	+11
M <sub>8</sub> -----	0.0369	+14	0.0383	+15	0.0398	+14	0.0412	+15	0.0427	+14	0.0441	+15
O <sub>1</sub> Q <sub>1</sub> 2Q <sub>1</sub> p <sub>1</sub> -----	9.9509	-13	9.9496	-13	9.9483	-13	9.9470	-13	9.9457	-13	9.9444	-13
OO-----	9.8344	-47	9.8297	-47	9.8250	-47	9.8203	-46	9.8157	-46	9.8111	-46
MK-----	9.9782	-5	9.9777	-5	9.9772	-5	9.9767	-5	9.9762	-4	9.9758	-5
2MK-----	9.9874	-2	9.9872	-1	9.9871	-1	9.9870	-1	9.9869	-1	9.9868	-1
Mf-----	9.8926	-30	9.8896	-30	9.8866	-29	9.8837	-30	9.8807	-30	9.8777	-29
Mm-----	0.0336	+14	0.0350	+13	0.0363	+13	0.0376	+13	0.0389	+14	0.0403	+13

\*Log *F* of  $\lambda_2$ ,  $\mu_2$ ,  $\nu_2$ , MS, 2SM, and MSf are each equal to log *F* of M<sub>2</sub>.Log *F* of P<sub>1</sub>, R<sub>2</sub>, S<sub>1</sub>, S<sub>2</sub>, S<sub>4</sub>, S<sub>6</sub>, T<sub>2</sub>, Sa, and Ssa are each zero.For log *F* of L<sub>2</sub> and M<sub>1</sub> see Table 13.

Table 12.—*Log factor F corresponding to every tenth of a degree of I—Con.*

<i>I</i>	27.3°	Diff.	27.4°	Diff.	27.5°	Diff.	27.6°	Diff.	27.7°	Diff.
Constituent										
J <sub>1</sub> .....	9.9469	-11	9.9458	-11	9.9447	-10	9.9437	-11	9.9426	-10
K <sub>1</sub> .....	9.9639	-8	9.9631	-8	9.9623	-8	9.9615	-8	9.9607	-8
K <sub>2</sub> .....	9.9114	-24	9.9090	-24	9.9066	-24	9.9042	-24	9.9018	-24
M <sub>2</sub> * , N <sub>2</sub> , 2N.....	0.0114	+4	0.0118	+3	0.0121	+4	0.0125	+4	0.0129	+4
M <sub>3</sub> .....	0.0171	+6	0.0177	+5	0.0182	+6	0.0188	+5	0.0193	+6
M <sub>4</sub> , MN.....	0.0228	+7	0.0235	+8	0.0243	+7	0.0250	+8	0.0258	+7
M <sub>5</sub> .....	0.0342	+11	0.0353	+11	0.0364	+11	0.0375	+12	0.0387	+11
M <sub>6</sub> .....	0.0456	+15	0.0471	+15	0.0486	+15	0.0501	+14	0.0515	+15
O <sub>1</sub> , Q <sub>1</sub> , 2Q, p <sub>1</sub> .....	9.9431	-13	9.9418	-13	9.9405	-12	9.9393	-13	9.9380	-12
OO.....	9.8065	-46	9.8019	-46	9.7973	-45	9.7928	-45	9.7883	-45
MK.....	9.9753	-4	9.9749	-5	9.9744	-4	9.9740	-5	9.9735	-4
2MK.....	9.9867	-1	9.9866	-1	9.9865	0	9.9865	-1	9.9864	0
Mf.....	9.8748	-29	9.8719	-30	9.8689	-29	9.8660	-29	9.8631	-28
Mm.....	0.0416	+14	0.0430	+14	0.0444	+13	0.0457	+14	0.0471	+14

<i>I</i>	27.8°	Diff.	27.9°	Diff.	28.0°	Diff.	28.1°	Diff.	28.2°	Diff.
Constituent										
J <sub>1</sub> .....	9.9416	-11	9.9405	-10	9.9395	-10	9.9385	-10	9.9375	-10
K <sub>1</sub> .....	9.9599	-9	9.9590	-8	9.9582	-8	9.9574	-7	9.9567	-8
K <sub>2</sub> .....	9.8994	-24	9.8970	-24	9.8946	-24	9.8922	-24	9.8898	-24
M <sub>2</sub> * , N <sub>2</sub> , 2N.....	0.0133	+3	0.0136	+4	0.0140	+4	0.0144	+4	0.0148	+4
M <sub>3</sub> .....	0.0199	+6	0.0205	+5	0.0210	+6	0.0216	+6	0.0222	+5
M <sub>4</sub> , MN.....	0.0265	+8	0.0273	+7	0.0280	+8	0.0288	+7	0.0295	+8
M <sub>5</sub> .....	0.0398	+11	0.0409	+11	0.0420	+12	0.0432	+11	0.0443	+12
M <sub>6</sub> .....	0.0530	+15	0.0545	+16	0.0561	+15	0.0576	+15	0.0591	+15
O <sub>1</sub> , Q <sub>1</sub> , 2Q, p <sub>1</sub> .....	9.9368	-13	9.9355	-12	9.9343	-13	9.9330	-12	9.9318	-12
OO.....	9.7538	-45	9.7793	-45	9.7748	-45	9.7703	-44	9.7659	-44
MK.....	9.9731	-4	9.9727	-4	9.9723	-5	9.9718	-4	9.9714	-4
2MK.....	9.9864	-1	9.9863	0	9.9863	-1	9.9862	0	9.9862	0
Mf.....	9.8603	-29	9.8574	-29	9.8545	-28	9.8517	-28	9.8489	-29
Mm.....	0.0485	+14	0.0499	+14	0.0513	+14	0.0527	+15	0.0542	+14

<i>I</i>	28.3°	Diff.	28.4°	Diff.	28.5°	Diff.	28.6°
Constituent							
J <sub>1</sub> .....	9.9365	-10	9.9355	-10	9.9345	-10	9.9335
K <sub>1</sub> .....	9.9559	-8	9.9551	-8	9.9543	-8	9.9535
K <sub>2</sub> .....	9.8874	-24	9.8850	-24	9.8826	-23	9.8803
M <sub>2</sub> * , N <sub>2</sub> , 2N.....	0.0152	+3	0.0155	+4	0.0159	+4	0.0163
M <sub>3</sub> .....	0.0227	+6	0.0233	+6	0.0239	+6	0.0245
M <sub>4</sub> , MN.....	0.0303	+8	0.0311	+7	0.0318	+8	0.0326
M <sub>5</sub> .....	0.0455	+11	0.0466	+12	0.0478	+11	0.0489
M <sub>6</sub> .....	0.0606	+15	0.0621	+16	0.0637	+15	0.0652
O <sub>1</sub> , Q <sub>1</sub> , 2Q, p <sub>1</sub> .....	9.9306	-12	9.9294	-12	9.9282	-12	9.9270
OO.....	9.7615	-44	9.7571	-44	9.7527	-44	9.7483
MK.....	9.9710	-4	9.9706	-4	9.9702	-4	9.9698
2MK.....	9.9862	0	9.9862	0	9.9862	0	9.9862
Mf.....	9.8460	-28	9.8432	-28	9.8404	-28	9.8376
Mm.....	0.0556	+14	0.0570	+14	0.0584	+15	0.0599

\*Log *F* of λ<sub>2</sub>, μ<sub>2</sub> ν<sub>2</sub>, MS, and MSf are each equal to log *F* of M<sub>2</sub>.Log *F* of P<sub>1</sub>, R<sub>2</sub>, S<sub>1</sub>, S<sub>2</sub>, S<sub>6</sub>, T<sub>2</sub>, Sa, and Ssa are each zero.For log *F* of L<sub>2</sub> and M<sub>1</sub> see Table 13.

Table 13.—Values of  $u$  and  $\log F$  of  $L_2$  and  $M_1$  for years 1900 to 2000

Year	$N$	$u$ of $L_2$	Diff.	$u$ of $M_1$	Diff.	$\log F(L_2)$	Diff.	$\log F(M_1)$	Diff.	$N$
1899	260	° +11.4	° 6.3	° 353.5	° 5.2	0.0964	161	9.7295	35	° 260
1900	255	+5.1	6.4	358.7	5.0	0.1125	73	9.7260	78	255
	250	-1.3	4.9	3.7	5.1	0.1052	259	9.7338	189	250
	245	-6.2	2.9	8.8	5.4	0.0793	348	9.7527	297	245
	240	-9.1	0.9	14.2	6.2	0.0445	353	9.7824	404	240
1901	235	-10.0	0.6	20.4	7.2	0.0092	313	9.8228	506	235
	230	-9.4	1.6	27.6	8.9	0.9779	250	9.8734	596	230
	225	-7.8	2.3	36.5	11.4	0.9529	182	9.9330	647	225
1902	220	-5.5	2.5	47.9	14.6	0.9347	114	9.9977	609	220
	215	-3.0	2.6	62.5	18.0	0.9233	50	0.0586	412	215
	210	-0.4	2.6	80.5	9.183	0.9183	10	0.0998	58	210
	205	+2.2	2.3	99.9	17.7	0.9193	61	0.1056	276	205
1903	200	+4.5	2.0	117.6	14.4	0.9254	115	0.0780	497	200
	195	+6.5	1.5	132.0	11.0	0.9369	154	0.0233	523	195
	190	+8.0	0.8	143.0	8.6	0.9523	190	0.9760	485	190
	185	+8.8	0.1	151.6	7.0	0.9713	213	0.9275	414	185
1904	180	+8.9	0.7	158.6	5.8	0.9926	222	9.8861	336	180
	175	+8.2	1.8	164.4	5.1	0.0148	209	9.8525	256	175
	170	+6.4	2.8	169.5	4.6	0.0357	166	9.8269	178	170
	165	+3.6	3.5	174.1	4.5	0.0523	93	9.8091	100	165
1905	160	+0.1	3.8	178.6	4.5	0.0616	5	9.7991	21	160
	155	-3.7	3.6	183.1	4.6	0.0611	109	9.7970	63	155
	150	-7.3	2.8	187.7	5.1	0.0502	195	9.8033	152	150
	145	-10.1	1.7	192.8	5.8	0.0307	249	9.8185	248	145
1906	140	-11.8	0.3	198.6	7.0	0.0058	208	9.8433	349	140
	135	-12.1	0.8	205.6	8.8	0.9790	254	9.8782	135	
	130	-11.3	1.9	214.4	11.5	0.9536	216	9.9228	446	130
	125	-9.4	2.7	225.9	15.4	0.9320	161	9.9744	494	125
1907	120	-6.7	3.3	241.3	19.4	0.9159	90	0.0238	283	120
	115	-3.4	3.7	260.7	21.1	0.9069	11	0.0521	124	115
	110	+0.3	3.9	251.8	19.0	0.9058	79	0.0397	514	110
	105	+4.2	3.8	300.8	15.0	0.9137	177	0.9883	688	105
1908	100	+8.0	3.2	315.8	11.3	0.9314	282	0.9195	683	100
	95	+11.2	2.2	327.1	8.7	0.9596	390	0.8512	591	95
	90	+13.4	0.4	335.8	7.2	0.9986	487	0.7921	472	90
1909	85	+13.8	2.5	343.0	6.1	0.0473	536	9.7449	346	85
	80	+11.3	6.0	319.1	5.5	0.1009	463	9.7103	220	80
	75	+5.3	9.2	354.6	5.4	0.1472	491	9.6883	75	
	70	-3.9	9.3	0.0	5.5	0.1663	209	9.6789	33	70
1910	65	-13.2	6.2	5.5	5.8	0.1454	500	9.6822	164	65
	60	-19.4	2.1	11.3	6.7	0.0945	602	0.6986	299	60
	55	-21.5	1.1	18.0	8.2	0.0343	556	9.7285	437	55
	50	-20.4	3.4	26.2	10.5	0.9787	449	9.7722	561	50
1911	45	-17.0	4.8	36.7	14.1	0.9338	324	0.8286	635	45
	40	-12.2	5.8	50.8	19.0	0.9014	191	0.8921	540	40
	35	-6.4	1.1	69.8	22.8	0.9883	59	0.9461	165	35
	30	-0.2	6.2	92.6	22.2	0.9764	77	0.9626	343	30
1912	25	+6.1	6.0	114.8	17.7	0.8841	219	0.9283	25	
	20	+12.1	5.3	132.5	13.1	0.9060	368	0.8640	681	20
	15	+17.4	3.9	145.6	9.9	0.9428	526	0.7959	585	15
	10	+21.3	1.4	155.5	7.7	0.9954	680	0.7374	450	10
1913	5	+22.7	2.8	163.2	6.6	0.0634	780	0.6924	308	5
	0	+19.9	8.9	169.8	5.9	0.1414	675	0.6616	0	
	355	+11.0	13.6	175.7	5.6	0.2089	187	0.6447	169	355
	350	-2.6	11.9	181.3	5.7	0.2276	450	0.6414	33	350
1914	345	-14.5	6.0	187.0	6.1	0.1826	754	0.6515	240	345
	340	-20.5	0.7	193.1	6.9	0.1072	738	0.6755	378	340
	335	-21.2	2.5	200.0	8.5	0.0334	604	0.7133	335	
	330	-18.7	2.5	208.5	9.9730			0.7651	518	330

Table 13.—Values of  $u$  and  $\log F$  of  $L_2$  and  $M_1$  for years 1900 to 2000—Con.

Year	$N$	$u$ of $L_2$	Diff.	$u$ of $M_1$	Diff.	$\log F(L_2)$	Diff.	$\log F(M_1)$	Diff.	$N$
1914	330	-18.7	4.4	208.5	10.9	9.9730	447	9.7651	643	330
1915	325	-14.3	5.5	219.4	14.6	9.9283	295	9.8294	699	325
	320	-8.8	5.9	234.0	19.5	9.8988	153	9.8993	574	320
	315	-2.9	5.9	253.5	22.8	9.8835	20	9.9567	174	315
	310	+3.0	5.6	276.3	21.6	9.8815	108	9.9741	316	310
1916	305	+8.6	5.0	297.9	17.0	9.8923	230	9.9425	581	305
	300	+13.6	3.8	314.9	12.5	9.9153	346	9.8844	600	300
	295	+17.4	2.2	327.4	9.4	9.9499	449	9.8244	506	295
1917	290	+19.6	0.2	336.8	7.5	9.9948	510	9.7738	370	290
	285	+19.4	3.2	344.3	6.3	0.0458	496	9.7368	239	285
	280	+16.1	6.5	350.6	5.6	0.0954	351	9.7129	111	280
	275	+9.6	7.9	356.2	5.2	0.1305	63	9.7018	15	275
1918	270	+1.7	7.1	1.4	5.2	0.1368	227	9.7033	270	270
	265	-5.4	4.4	6.6	5.5	0.1141	401	9.7167	134	265
	260	-9.8	1.8	12.1	6.0	0.0740	440	9.7419	252	260
	255	-11.6	0.4	18.1	7.1	0.0300	398	9.7759	486	255
1919	250	-11.2	1.8	25.2	8.6	9.9902	320	9.8275	250	250
	245	-9.4	2.6	33.8	11.2	9.9582	234	9.8869	594	245
	240	-6.8	3.1	45.0	14.5	9.9348	148	9.9539	664	240
	235	-3.7	3.2	59.5	18.2	9.9200	67	0.0203	493	235
1920	230	-0.5	3.1	77.7	20.2	9.9133	6	0.0696	230	230
	225	+2.6	2.8	97.9	18.8	9.9139	71	0.0830	134	225
	220	+5.4	2.2	116.7	15.1	9.9210	128	0.0580	250	220
	215	+7.6	1.6	131.8	11.5	9.938	174	0.0113	510	215
1921	210	+9.2	0.9	143.3	8.9	9.9512	209	9.9603	210	210
	205	+10.1	0.0	152.2	7.0	9.9721	227	9.9141	462	205
	200	+10.1	1.1	159.2	5.9	9.9948	224	9.8760	381	200
	195	+9.0	2.0	165.1	5.0	0.0172	196	9.8466	294	195
1922	190	+7.0	2.9	170.1	4.7	0.0368	141	9.8257	190	190
	185	+4.1	3.5	174.8	4.4	0.0509	63	9.8127	185	185
	180	+0.6	3.5	179.2	4.3	0.0572	27	9.8073	54	180
	175	-2.9	3.2	183.5	4.5	0.0545	113	9.8095	22	175
1923	170	-6.1	2.4	188.0	4.9	0.0432	179	9.8192	176	170
	165	-8.5	1.4	192.9	5.6	0.0253	218	9.8368	259	165
	160	-9.9	0.4	198.5	6.5	0.0035	231	9.8627	347	160
1924	155	-10.3	0.5	205.0	8.2	9.9804	220	9.8974	155	155
	150	-9.8	1.4	213.2	10.4	9.9584	191	9.9407	433	150
	145	-8.4	2.1	223.6	13.9	9.9393	148	9.9906	498	145
	140	-6.3	2.6	237.5	17.6	9.9245	94	0.0404	351	140
1925	135	-3.7	3.0	255.1	20.2	9.9151	30	0.0755	9	135
	130	-0.7	3.3	275.3	19.3	9.9121	42	0.0764	130	130
	125	+2.6	3.2	294.6	15.9	9.9163	121	0.0377	631	125
	120	+5.8	3.2	310.5	12.2	9.9284	207	9.9746	683	120
1926	115	+8.7	2.2	322.7	9.4	9.9491	296	9.9063	626	115
	110	+10.9	0.9	332.1	7.6	9.9787	380	9.8437	525	110
	105	+11.8	0.9	339.7	6.3	0.0167	440	9.7912	409	105
	100	+10.9	3.6	346.0	5.7	0.0607	434	9.7503	290	100
1927	95	+7.3	6.4	351.7	5.3	0.1041	306	9.7213	172	95
	90	+0.9	8.1	357.0	5.2	0.1347	34	9.7041	52	90
	85	-7.2	7.3	2.2	5.4	0.1381	276	9.6989	71	85
	80	-14.5	4.4	7.6	6.0	0.1105	475	9.7060	200	80
1928	75	-18.9	1.1	13.6	7.1	0.0630	525	9.7260	332	75
	70	-20.0	1.6	20.7	8.7	0.0105	477	9.7592	466	70
	65	-18.4	3.4	29.4	9.6	9.9628	383	9.8058	579	65
	60	-15.0	4.7	40.8	15.5	9.9245	268	9.8637	609	60
1929	55	-10.3	5.5	56.3	20.4	9.8977	146	9.9246	431	55
	50	-4.8	5.9	76.7	23.0	9.8831	20	9.9677	13	50
	45	+1.1	6.0	99.7	20.9	9.8811	112	9.9664	45	45
	40	+7.1		120.6		9.8923		9.9184	480	40

Table 13.—Values of  $u$  and  $\log F$  of  $L_2$  and  $M_1$  for years 1900 to 2000—Con.

Year	$N$	$u$ of $L_2$	Diff.	$u$ of $M_1$	Diff.	$\log F(L_2)$	Diff.	$\log F(M_1)$	Diff.	$N$
1929	40	○ +7.1	5.7	120.6	16.1	9.8923	252	9.9184	686	○ 40
1930	35	+12.8	4.8	136.7	11.9	9.9175	401	9.8498	670	35
	30	+17.6	3.3	148.6	9.0	9.9576	558	9.7828	558	30
	25	+20.9	0.4	157.6	7.4	0.0134	706	9.7270	420	25
	20	+21.3	4.2	165.0	6.3	0.0840	774	9.6850	281	20
1931	15	+17.1	10.4	171.3	5.7	0.1614	588	9.6569	141	15
	10	+6.7	13.9	177.0	5.6	0.2202	16	9.6428	7	10
	5	-7.2	10.6	182.6	5.8	0.2218	572	9.6421	129	5
1932	0	-17.8	4.5	188.4	6.3	0.1646	781	9.6550	0	0
	355	-22.3	0.3	194.7	7.3	0.0865	720	9.6817	267	355
	350	-22.0	3.3	202.0	9.1	0.0145	576	9.7225	546	350
	345	-18.7	4.9	211.1	12.0	9.9569	412	9.7771	662	345
1933	340	-13.8	5.9	223.1	16.2	9.9157	262	9.8433	340	340
	335	-7.9	6.3	239.3	21.0	9.8895	119	9.9110	677	335
	330	-1.6	6.2	260.3	23.2	9.8776	18	9.9578	468	330
	325	+4.6	5.9	283.5	20.4	9.8794	151	9.9572	449	325
1934	320	+10.5	5.1	303.9	15.6	9.8945	282	9.9123	633	320
	315	+15.6	3.9	319.5	11.4	9.9227	410	9.8490	594	315
	310	+19.5	1.9	330.9	8.7	9.9637	523	9.7896	478	310
	305	+21.4	1.1	339.6	7.1	0.0160	592	9.7418	340	305
1935	300	+20.3	4.9	346.7	6.1	0.0752	550	9.7078	204	300
	295	+15.4	8.5	352.8	5.5	0.1302	316	9.6874	71	295
	290	+6.9	9.4	358.3	5.4	0.1618	70	9.6803	55	290
	285	-2.5	7.1	37	5.4	0.1548	393	9.6858	181	285
1936	280	-9.6	9.1			0.1155	530	9.7039	310	280
	275	-13.2	3.6	15.1	6.0	0.0625	496	9.7349	428	275
	270	-13.5	0.3	21.7	6.6	0.0129	418	9.7777	270	270
	265	-11.9	1.6	29.8	8.1	9.9711	313	9.8331	658	265
1937	260	-9.0	3.6	40.2	13.6	9.9398	208	9.8989	701	260
	255	-5.4	3.8	53.8	17.7	9.9190	103	9.9690	595	255
	250	-1.6	3.8	71.5	20.7	9.9087	24	0.0285	270	250
	245	+2.2	3.4	92.2	20.3	9.9063	63	0.0555	160	245
1938	240	+5.6	2.9	112.5	16.8	9.9126	136	0.0395	446	240
	235	+8.5	2.1	129.3	12.7	9.9262	196	9.9949	522	235
	230	+10.6	1.2	142.0	9.6	9.9458	240	9.9427	475	230
	225	+11.8	0.0	151.6	7.4	9.9698	262	9.8952	383	225
1939	220	+11.8	1.1	159.0	6.2	9.9960	256	9.8569	284	220
	215	+10.7	2.4	165.2	5.3	0.0216	216	9.8285	215	215
	210	+8.3	3.3	170.5	4.7	0.0432	141	9.8099	186	210
1940	205	+5.0	3.8	175.2	4.5	0.0573	44	9.8002	13	205
	200	+1.2	3.6	179.7	4.5	0.0617	56	9.7989	68	200
	195	-2.4	3.1	184.2	4.5	0.0561	139	9.8057	145	195
	190	-5.5	2.1	188.7	4.9	0.0422	194	9.8202	223	190
1941	185	-7.6	1.2	193.6	5.5	0.0228	218	9.8425	301	185
	180	-8.8	0.3	199.1	6.5	0.0010	219	9.8726	381	180
	175	-9.1	0.6	205.6	7.9	9.9791	200	9.9107	457	175
	170	-8.5	1.3	213.5	10.1	9.9591	171	9.9564	513	170
1942	165	-7.2	1.8	223.6	13.0	9.9420	130	0.0077	509	165
	160	-5.4	2.2	236.6	16.6	9.9290	83	0.0586	380	160
	155	-3.2	2.6	253.2	19.2	9.9207	31	0.0966	78	155
	150	-0.6	2.7	272.4	18.9	9.9176	28	0.1044	299	150
1943	145	+2.1	2.6	291.3	16.0	9.9204	91	0.0745	563	145
	140	+4.7	2.4	307.3	12.5	9.9295	160	0.0182	652	140
	135	+7.1	1.9	319.8	9.8	9.9455	230	9.9530	624	135
	130	+9.0	1.0	329.6	7.7	9.9685	297	9.8906	542	130
1944	125	+10.0	0.3	337.3	6.5	9.9982	349	9.8364	441	125
	120	+9.7	2.1	343.8	5.6	0.0331	365	9.7923	334	120
	115	+7.6	4.4	349.4	5.2	0.0696	306	9.7589	225	115
	110	+3.2		354.6		0.1002	9.7364		110	

Table 13.—Values of  $u$  and  $\log F$  of  $L_2$  and  $M_1$  for years 1900 to 2000—Con.

Year	$N$	$u$ of $L_2$	Diff.	$u$ of $M_1$	Diff.	$\log F(L_2)$	Diff.	$\log F(M_1)$	Diff.	$N$
1944	110	+3.2	6.2	354.6	5.1	0.1002	147	9.7364	115	110
1945	105	-3.0	6.6	359.7	5.1	0.1149	88	9.7249	3	105
	100	-9.6	5.3	4.8	5.5	0.1061	304	9.7246	117	100
	95	-14.9	2.8	10.3	6.2	0.0757	426	9.7363	240	95
	90	-17.7	0.2	16.5	7.5	0.0331	446	9.7603	370	90
1946	85	-17.9	1.8	24.0	9.5	0.9885	397	9.7973	494	85
	80	-16.1	3.4	33.5	12.6	0.9488	325	9.8467	583	80
	75	-12.7	4.5	46.1	17.0	0.9163	200	9.9050	553	75
1947	70	-8.2	5.2	63.1	21.5	0.8963	99	9.9603	279	70
	65	-3.0	5.6	84.6	9.8864	20	9.9882	205	65	
	60	+2.6	5.5	107.2	22.6	0.8884	146	9.9677	589	60
	55	+8.1	5.1	126.2	19.0	0.9030	282	9.9088	706	55
1948	50	+13.2	4.2	140.7	10.7	0.9812	428	9.8382	648	50
	45	+17.4	2.4	151.4	8.4	0.9740	579	9.7734	526	45
	40	+19.8	0.8	159.8	6.9	0.0319	706	9.7208	388	40
	35	+19.0	5.6	166.7	6.0	0.1025	724	9.6820	250	35
1949	30	+13.4	11.4	172.7	5.7	0.1749	447	9.6570	115	30
	25	+2.0	13.2	178.4	5.6	0.2196	150	9.6455	20	25
	20	-11.2	13.2	184.0	5.8	0.2046	639	9.6475	154	20
	15	-20.1	8.9	189.8	6.6	0.1407	764	9.6629	295	15
1950	10	-23.1	1.1	196.4	7.7	0.0643	680	9.6924	437	10
	5	-22.0	3.8	204.1	9.8	0.9963	532	9.7361	572	5
	0	-18.2	5.3	213.9	9.431	0.9431	377	9.7933	670	0
	355	-12.9	6.1	227.0	13.1	0.9054	227	9.8603	634	355
1951	350	-6.8	6.5	244.7	22.3	0.8827	84	9.9237	335	350
	345	-0.3	6.4	267.0	22.9	0.98743	54	9.9572	345	345
	340	+6.1	6.0	289.9	19.1	0.8979	193	9.9398	174	340
	335	+12.1	5.2	309.0	14.2	0.8990	332	9.8843	652	335
1952	330	+17.3	3.8	323.2	10.6	0.9322	471	9.8191	579	330
	325	+21.1	1.5	333.8	8.2	0.9793	597	9.7612	450	325
	320	+22.6	2.1	342.0	6.7	0.0390	666	9.7162	310	320
	315	+20.5	6.7	348.7	6.0	0.1056	583	9.6852	172	315
1953	310	+13.8	10.6	354.7	5.5	0.1639	242	9.6680	38	310
	305	+3.2	0.2	0.2	5.5	0.1881	238	9.6642	91	305
	300	-7.2	10.4	5.7	5.7	0.1643	558	9.6733	300	300
	295	-13.6	6.4	11.4	6.3	0.1085	600	9.6958	225	295
1954	290	-15.9	0.9	17.7	7.4	0.0485	537	9.7307	484	290
	285	-15.0	2.8	25.1	9.3	0.9948	420	9.7791	608	285
	280	-12.2	3.8	34.4	12.2	0.9528	298	9.8599	703	280
1955	275	-8.4	4.4	46.6	16.1	0.9230	175	9.9102	674	275
	270	-4.0	4.5	62.7	20.2	0.9055	68	9.9776	449	270
	265	+0.5	8.2	89.9	21.8	0.8987	33	0.0225	12	265
	260	+4.8	4.3	104.7	19.0	0.9020	124	0.0237	379	260
1956	255	+8.5	3.0	123.7	14.6	0.9144	204	9.9858	538	255
	250	+11.5	1.9	138.3	10.9	0.9348	267	9.9320	516	250
	245	+13.4	0.6	149.2	8.4	0.9615	308	9.8804	424	245
	240	+14.0	0.9	157.6	6.7	0.9923	315	9.8380	312	240
1957	235	+13.1	2.5	164.3	5.6	0.0238	273	9.8068	202	235
	230	+10.6	3.9	169.9	5.1	0.0511	182	9.7866	97	230
	225	+6.7	4.4	175.0	4.7	0.0693	53	9.7769	1	225
	220	+2.3	4.3	179.7	4.6	0.0746	77	9.7770	92	220
1958	215	-2.0	3.4	184.3	4.7	0.0669	177	9.7862	180	215
	210	-5.4	2.1	189.0	5.1	0.0492	232	9.8042	266	210
	205	-7.5	1.0	194.1	5.6	0.0260	246	9.8308	350	205
	200	-8.5	0.0	199.7	6.6	0.0014	233	9.8658	432	200
1959	195	-8.5	0.9	206.3	8.1	0.9781	202	9.9090	506	195
	190	-7.6	1.3	214.4	10.2	0.9579	162	9.9596	555	190
	185	-6.3	1.9	224.6	13.1	0.9417	118	0.0151	185	185
	180	-4.4	2.7	237.7	0.0	0.9299	0.0692	0.0692	541	180

Table 13.—Values of  $u$  and  $\log F$  of  $L_2$  and  $M_1$  for years 1900 to 2000—Con.

Year	$N$	$u$ of $L_2$	Diff.	$u$ of $M_1$	Diff.	$\log F(L_2)$	Diff.	$\log F(M_1)$	Diff.	$N$
1959	180	−4.4	2.1	237.7	16.4	9.9299	71	0.0692	404	180
1960	175	−2.3	2.3	254.1	18.6	9.9228	21	0.1096	107	175
	170	0.0	2.3	272.7	18.3	9.9207	30	0.1203	252	170
	165	+2.3	2.2	291.0	15.6	9.9237	83	0.0951	507	165
	160	+4.5	2.0	306.6	12.3	9.9320	137	0.0444	603	160
1961	155	+6.5	1.6	318.9	9.6	9.9457	191	0.9841	589	155
	150	+8.1	0.8	328.5	7.6	9.9648	242	0.9252	523	150
	145	+8.9	0.2	336.1	6.4	9.9890	280	0.8729	435	145
1962	140	+8.7	1.5	342.5	5.6	0.0170	295	0.8294	344	140
	135	+7.2	3.1	348.1	5.1	0.0465	263	0.7950	247	135
	130	+4.1	4.7	353.2	4.8	0.0728	166	0.7703	149	130
	125	−0.6	5.4	358.0	4.9	0.0894	8	0.7554	47	125
1963	120	−6.0	2.9	2.9	5.1	0.0902	169	0.7507	58	120
	115	−11.0	5.0	8.0	5.6	0.0733	305	0.7565	170	115
	110	−14.5	3.5	13.6	6.6	0.0428	370	0.7735	290	110
	105	−16.0	1.5	20.2	8.1	0.0058	368	0.8025	412	105
1964	100	−15.5	2.1	28.3	10.5	0.9690	320	0.8437	510	100
	95	−13.4	3.3	38.8	14.1	0.9370	245	0.8956	567	95
	90	−10.1	4.2	52.9	18.7	0.9125	156	0.9523	465	90
	85	−5.9	4.8	71.6	22.1	0.8969	48	0.9988	61	85
1965	80	−1.1	5.1	93.7	21.2	0.8921	60	0.0049	392	80
	75	+4.0	5.0	114.9	17.0	0.8981	180	0.9657	666	75
	70	+9.0	4.3	131.9	12.8	0.9161	309	0.8991	702	70
	65	+13.3	3.2	144.7	9.6	0.9470	447	0.8289	615	65
1966	60	+16.5	1.4	154.3	7.7	0.9917	581	0.7674	489	60
	55	+17.9	2.0	162.0	6.5	0.0498	675	0.7185	353	55
	50	+15.9	7.0	168.5	5.8	0.1173	622	0.6832	218	50
	45	+8.9	11.5	174.3	5.5	0.1795	274	0.6614	85	45
1967	40	−2.6	11.6	179.8	5.6	0.2069	283	0.6529	48	40
	35	−14.2	185.4	6.0	0.1786	651	0.6577	182	35	
	30	−21.2	7.0	191.4	6.8	0.1135	717	0.6759	322	30
	25	−23.1	1.9	198.2	8.2	0.0418	624	0.7081	464	25
1968	20	−21.3	4.1	206.4	10.6	0.9794	484	0.7545	592	20
	15	−17.2	217.0	14.3	0.9310	335	0.8137	666	15	
	10	−11.8	231.3	19.2	0.8975	190	0.8803	567	10	
	5	−5.6	250.5	23.0	0.8785	50	0.9370	185	5	
1969	0	+0.9	6.5	273.5	22.2	0.8735	91	0.9555	0	0
	355	+7.4	6.1	295.7	17.7	0.8826	232	0.9227	624	355
	350	+13.5	5.1	313.4	13.0	0.9058	379	0.8603	657	350
1970	345	+18.6	3.6	326.4	9.8	0.9437	528	0.7946	345	345
	340	+22.2	0.9	336.2	7.7	0.9965	663	0.7388	558	340
	335	+23.1	3.3	343.9	6.5	0.0628	727	0.6965	423	335
	330	+19.8	8.7	350.4	5.9	0.1355	583	0.6683	282	330
1971	325	+11.1	12.2	356.3	5.5	0.1938	121	0.6540	9	325
	320	−1.1	1.8	5.5	0.2059	416	0.6531	125	320	
	315	−11.7	10.6	7.4	0.1643	669	0.6656	256	315	
	310	−17.1	5.4	13.4	6.0	0.0974	661	0.6912	393	310
1972	305	−18.0	2.1	20.2	8.2	0.0313	546	0.7305	528	305
	300	−15.9	28.4	10.4	0.9767	408	0.7833	649	300	
	295	−12.2	37.7	38.8	0.9359	271	0.8482	709	295	
	290	−7.5	4.7	52.8	14.0	0.9088	143	0.9191	611	290
1973	285	−2.4	5.1	71.3	22.0	0.8945	26	0.9802	285	285
	280	+2.7	4.6	93.3	21.6	0.8919	86	0.0058	224	280
	275	+7.3	4.0	114.9	17.1	0.9005	183	0.9834	508	275
	270	+11.3	3.1	132.0	12.8	0.9188	274	0.9326	566	270
1974	265	+14.4	1.6	144.8	9.7	0.9462	343	0.8760	491	265
	260	+16.0	0.1	154.5	7.6	0.9805	380	0.8269	374	260
	255	+15.9	2.2	162.1	6.2	0.0185	364	0.7895	255	255
	250	+13.7	168.3	0.0549	0.97642	0.97642	253	0.7642	250	250

Table 13.—Values of  $u$  and  $\log F$  of  $L_2$  and  $M_1$  for years 1900 to 2000—Con.

Year	$N$	$u$ of $L_2$	Diff.	$u$ of $M_1$	Diff.	$\log F(L_2)$	Diff.	$\log F(M_1)$	Diff.	$N$
1974	250	° +13.7	4.1	° 168.3	5.5	0.0549	274	9.7642	134	° 250
1975	245	+9.6	5.4	173.8	5.0	0.0823	116	9.7508	22	245
	240	+4.2	5.3	178.8	4.8	0.0939	68	9.7486	22	240
	235	-1.1	4.2	183.6	4.9	0.0871	211	9.7568	82	235
	230	-5.3	2.7	188.5	5.3	0.0660	286	9.7751	183	230
1976	225	-8.0	1.0	193.8	5.8	0.0374	300	9.8033	378	225
	220	-9.0	0.2	199.6	6.8	0.0074	273	9.8411	471	220
	215	-8.8	1.1	206.4	8.3	9.9801	228	9.8882	554	215
	210	-7.7	1.8	214.7	10.5	9.9573	174	9.9436	606	210
1977	205	-5.9	2.2	225.2	13.5	9.9399	118	0.0042	589	205
	200	-3.7	2.3	238.7	16.8	9.9281	63	0.0631	439	200
	195	-1.4	2.4	255.5	18.7	9.9218	10	0.1070	129	195
1978	190	+1.0	2.2	274.2	18.1	9.9208	40	0.1199	228	190
	185	+3.2	2.0	292.3	15.3	9.9248	89	0.0971	469	185
	180	+5.2	1.7	307.6	11.9	9.9337	135	0.0502	553	180
	175	+6.9	1.2	319.5	9.3	9.9472	177	9.9949	536	175
1979	170	+8.1	0.5	328.8	7.4	9.9649	213	9.9413	477	170
	165	+8.6	0.3	336.2	6.2	9.9862	237	9.8936	398	165
	160	+8.3	1.4	342.4	5.4	0.0099	241	9.8538	317	160
	155	+6.9	2.6	347.8	4.9	0.0340	213	9.8221	232	155
1980	150	+4.3	3.7	352.7	4.6	0.0553	143	9.7989	147	150
	145	+0.6	4.3	357.3	4.6	0.0696	33	9.7842	59	145
	140	-3.7	4.3	1.9	0.0729	99	9.7783	33	140	
	135	-8.0	3.4	6.7	4.8	0.0630	214	9.7816	131	135
1981	130	-11.4	2.0	11.9	6.0	0.0416	287	9.7947	237	130
	125	-13.4	0.4	17.9	7.1	0.0129	312	9.8184	349	125
	120	-13.8	1.0	25.0	9.0	9.9817	293	9.8533	457	120
	115	-12.8	2.3	34.0	12.0	9.9524	246	9.8990	533	115
1982	110	-10.5	3.2	46.0	16.0	9.9278	177	9.9523	508	110
	105	-7.3	3.9	62.0	20.1	9.9101	95	0.0031	276	105
	100	-3.4	4.3	82.1	21.7	9.9006	2	0.0307	161	100
	95	+0.9	4.4	103.8	19.1	9.9004	100	0.0146	549	95
1983	90	+5.3	4.2	122.9	14.7	9.9104	211	9.9597	703	90
	85	+9.5	3.5	137.6	11.1	9.9315	330	9.8894	676	85
	80	+13.0	2.3	148.7	8.6	9.9645	454	9.8218	572	80
	75	+15.3	0.1	157.3	7.0	0.0099	560	9.7646	445	75
1984	70	+15.4	3.4	164.3	6.1	0.0659	604	9.7201	70	70
	65	+12.0	7.8	170.4	5.6	0.1263	478	9.6889	312	65
	60	+4.2	10.8	176.0	5.4	0.1741	98	9.6706	183	60
1985	55	-6.6	9.6	181.4	5.6	0.1839	365	9.6652	77	55
	50	-16.2	5.3	187.0	6.2	0.1474	625	9.6729	212	50
	45	-21.5	5.9	193.2	7.1	0.0849	650	9.6941	350	45
	40	-22.4	2.3	200.3	8.7	0.0199	560	9.7291	490	40
1986	35	-20.1	4.2	209.0	11.4	9.9639	429	9.7781	614	35
	30	-15.9	5.5	220.4	15.7	9.9210	290	9.8395	640	30
	25	-10.4	6.2	236.1	20.5	9.8920	152	9.9035	471	25
	20	-4.2	6.5	256.6	23.4	9.8768	14	9.9506	18	20
1987	15	+2.3	6.3	280.0	21.0	9.8754	126	9.9524	15	15
	10	+8.6	5.9	301.0	16.3	9.8880	270	9.9068	668	10
	5	+14.5	4.9	317.3	11.9	9.9150	422	9.8400	5	5
	0	+19.4	3.2	329.2	9.1	9.9572	579	9.7750	0	0
1988	355	+22.6	0.1	338.3	7.3	0.0151	717	9.7215	395	355
	350	+22.7	4.6	345.6	6.3	0.0868	761	9.6820	350	350
	345	+18.1	10.7	351.9	5.8	0.1629	536	9.6565	345	345
	340	+7.4	13.4	357.7	5.6	0.2165	39	9.6451	114	340
1989	335	-6.0	9.8	3.3	5.7	0.2126	573	9.6469	154	335
	330	-15.8	4.0	9.0	6.2	0.1553	744	9.6623	289	330
	325	-19.8	0.4	15.2	7.3	0.0809	677	9.6912	325	325
	320	-19.4	22.5	0.132	0.0132	9.7341	429	9.7341	320	320

Table 13.—Values of  $u$  and  $\log F$  of  $L_2$  and  $M_1$  for years 1900 to 2000—Con.

Year	$N$	$u$ of $L_2$	Diff.	$u$ of $M_1$	Diff.	$\log F(L_2)$	Diff.	$\log F(M_1)$	Diff.	$N$
1989	320	—19.4	3.0	22.5	9.0	0.0132	537	9.7341	565	320
1990	315	—16.4	4.6	31.5	11.7	9.9595	386	9.7906	676	315
	310	—11.8	5.3	43.2	15.8	9.9209	242	9.8582	694	310
	305	—6.5	5.6	59.0	9.8967	108	9.9276	500	305	
	300	—0.9	5.5	79.5	22.7	9.8859	17	9.9776	51	300
1991	295	+4.6	5.1	102.2	20.4	9.8876	135	9.9827	394	295
	290	+9.7	4.3	122.6	15.6	9.9011	246	9.9433	585	290
	285	+14.0	3.0	138.2	11.5	9.9257	344	9.8848	565	285
	280	+17.0	1.3	149.7	8.9	9.9601	428	9.8283	464	280
1992	275	+18.3	1.0	158.6	6.9	0.0029	450	9.7819	326	275
	270	+17.3	3.7	165.5	6.0	0.0479	408	9.7493	203	270
	265	+13.6	5.9	171.5	5.4	0.0887	250	9.7290	79	265
1993	260	+7.7	6.9	176.9	5.1	0.1137	10	9.7211	38	260
	255	+0.8	5.8	182.0	5.0	0.1147	214	9.7249	255	
	250	—5.0	3.6	187.0	5.4	0.0933	342	9.7400	151	250
	245	—8.6	1.6	192.4	5.9	0.0591	373	9.7662	262	245
1994	240	—10.2	0.3	198.3	6.9	0.0218	340	9.8033	477	240
	235	—9.9	1.4	205.2	8.3	9.9878	280	9.8510	576	235
	230	—8.5	2.1	213.5	10.6	9.9598	209	9.9086	230	
	225	—6.4	2.6	224.1	13.8	9.9389	137	9.9731	645	225
1995	220	—3.8	2.7	237.9	17.2	9.9252	69	0.0372	496	220
	215	—1.1	2.7	255.1	19.4	9.9183	6	0.0868	173	215
	210	+1.6	2.5	274.5	9.177	51	0.1041	202	210	
	205	+4.1	2.1	293.1	15.4	9.9228	102	0.0839	444	205
1996	200	+6.2	1.7	308.5	11.9	9.9330	147	0.0395	519	200
	195	+7.9	1.0	320.4	9.3	9.9477	184	9.9876	195	
	190	+8.9	0.3	329.7	7.3	9.9661	209	9.9382	427	190
	185	+9.2	0.6	337.0	6.0	9.9870	221	9.8955	348	185
1997	180	+8.6	1.6	343.0	5.3	0.0091	211	9.8607	268	180
	175	+7.0	2.5	348.3	4.7	0.0302	175	9.8339	175	
	170	+4.5	3.3	353.0	4.5	0.0477	109	9.8148	191	170
	165	+1.2	3.7	357.5	4.4	0.0586	19	9.8035	113	165
1998	160	—2.5	3.6	1.9	4.5	0.0605	81	9.7999	45	160
	155	—6.1	3.0	6.4	4.9	0.0524	169	9.8044	155	
	150	—9.1	2.0	11.3	5.6	0.0355	231	9.8174	130	150
	145	—11.1	0.7	16.9	6.5	0.0124	258	9.8395	221	145
1999	140	—11.8	0.4	23.4	8.1	9.9866	253	9.8714	416	140
	135	—11.4	3.1	31.5	10.5	9.9613	224	9.9130	497	135
	130	—9.9	2.4	42.0	13.9	9.9389	174	9.9627	514	130
2000	125	—7.5	3.1	55.9	18.0	9.9215	112	0.0141	375	125
	120	—4.4	3.5	73.9	20.9	9.9103	37	0.0516	24	120
	115	—0.9	3.7	94.8	19.9	9.9066	45	0.0540	398	115
	110	+2.8	114.7	9.9111	0.0142	110		110		

Table 14.—*Node factor f for middle of each year, 1850 to 1999*

Constituent	1850	1851	1852	1853	1854	1855	1856	1857	1858	1859
J <sub>1</sub>	0.892	0.948	1.007	1.061	1.105	1.138	1.158	1.165	1.160	1.141
K <sub>1</sub>	0.922	0.959	0.999	1.037	1.069	1.092	1.107	1.113	1.108	1.095
K <sub>2</sub>	0.816	0.887	0.977	1.075	1.168	1.246	1.298	1.317	1.302	1.254
L <sub>2</sub>	1.163	0.905	0.725	1.055	1.263	0.944	0.469	0.962	1.283	1.001
M <sub>1</sub>	1.023	1.675	1.974	1.559	1.118	1.860	2.348	1.872	1.177	1.776
M <sub>2*</sub> , N <sub>2</sub> , 2N, λ <sub>2</sub> , μ <sub>2</sub> , ν <sub>2</sub>	1.027	1.017	1.005	0.993	0.981	0.972	0.966	0.963	0.965	0.971
M <sub>3</sub>	1.042	1.026	1.008	0.989	0.972	0.958	0.949	0.945	0.948	0.957
M <sub>4</sub> , MN	1.056	1.035	1.011	0.986	0.963	0.944	0.932	0.928	0.931	0.942
M <sub>6</sub>	1.085	1.053	1.016	0.978	0.944	0.918	0.900	0.894	0.899	0.915
M <sub>8</sub>	1.114	1.071	1.021	0.971	0.927	0.892	0.869	0.861	0.867	0.888
O <sub>1</sub> , Q <sub>1</sub> , 2Q, p <sub>1</sub>	0.874	0.933	0.998	1.059	1.110	1.150	1.174	1.183	1.176	1.153
OO	0.631	0.786	0.983	1.204	1.422	1.608	1.735	1.783	1.745	1.627
MK	0.948	0.976	1.004	1.029	1.048	1.062	1.069	1.072	1.070	1.063
2MK	0.974	0.993	1.010	1.022	1.029	1.032	1.032	1.032	1.032	1.032
Mf	0.743	0.856	0.990	1.129	1.257	1.360	1.427	1.452	1.432	1.370
Mm	1.094	1.059	1.016	0.973	0.933	0.900	0.879	0.871	0.878	0.897
Constituent	1860	1861	1862	1863	1864	1865	1866	1867	1868	1869
J <sub>1</sub>	1.110	1.066	1.013	0.955	0.898	0.852	0.829	0.832	0.863	0.912
K <sub>1</sub>	1.072	1.041	1.004	0.964	0.926	0.898	0.883	0.885	0.904	0.936
K <sub>2</sub>	1.179	1.086	0.988	0.897	0.823	0.773	0.749	0.753	0.784	0.840
L <sub>2</sub>	0.568	0.924	1.225	1.117	0.865	0.879	1.082	1.190	1.091	0.840
M <sub>1</sub>	2.227	1.792	1.046	1.260	1.680	1.609	1.164	0.812	1.189	1.731
M <sub>2*</sub> , N <sub>2</sub> , 2N, λ <sub>2</sub> , μ <sub>2</sub> , ν <sub>2</sub>	0.980	0.991	1.004	1.016	1.026	1.034	1.038	1.037	1.032	1.024
M <sub>3</sub>	0.970	0.987	1.006	1.024	1.040	1.651	1.057	1.056	1.049	1.036
M <sub>4</sub> , MN	0.960	0.983	1.008	1.032	1.054	1.069	1.076	1.075	1.066	1.048
M <sub>6</sub>	0.941	0.974	1.011	1.049	1.081	1.105	1.117	1.115	1.100	1.073
M <sub>8</sub>	0.922	0.966	1.015	1.065	1.110	1.143	1.159	1.156	1.135	1.099
O <sub>1</sub> , Q <sub>1</sub> , 2Q, p <sub>1</sub>	1.116	1.065	1.005	0.941	0.880	0.832	0.808	0.812	0.843	0.896
OO	1.447	1.230	1.008	0.807	0.647	0.540	0.489	0.497	0.563	0.685
MK	1.050	1.032	1.007	0.979	0.951	0.928	0.916	0.918	0.933	0.958
2MK	1.029	1.023	1.011	0.995	0.976	0.960	0.950	0.952	0.964	0.981
Mf	1.270	1.145	1.007	0.872	0.755	0.670	0.629	0.635	0.689	0.783
Mm	0.928	0.968	1.011	1.054	1.091	1.117	1.130	1.128	1.111	1.082
Constituent	1870	1871	1872	1873	1874	1875	1876	1877	1878	1879
J <sub>1</sub>	0.971	1.028	1.079	1.120	1.147	1.162	1.164	1.154	1.130	1.094
K <sub>1</sub>	0.974	1.014	1.050	1.079	1.099	1.111	1.112	1.104	1.087	1.061
K <sub>2</sub>	0.920	1.014	1.112	1.201	1.269	1.309	1.315	1.287	1.227	1.144
L <sub>2</sub>	0.828	1.148	1.224	0.816	0.545	1.087	1.270	0.858	0.543	1.036
M <sub>1</sub>	1.811	1.300	1.185	2.004	2.286	1.656	1.227	1.998	2.269	1.645
M <sub>2*</sub> , N <sub>2</sub> , 2N, λ <sub>2</sub> , μ <sub>2</sub> , ν <sub>2</sub>	1.013	1.000	0.988	0.977	0.969	0.964	0.963	0.967	0.974	0.984
M <sub>3</sub>	1.019	1.001	0.982	0.966	0.954	0.947	0.946	0.951	0.961	0.976
M <sub>4</sub> , MN	1.026	1.001	0.976	0.955	0.939	0.930	0.928	0.935	0.949	0.968
M <sub>6</sub>	1.039	1.001	0.965	0.933	0.910	0.896	0.894	0.904	0.924	0.953
M <sub>8</sub>	1.052	1.002	0.953	0.912	0.881	0.864	0.862	0.874	0.900	0.938
O <sub>1</sub> , Q <sub>1</sub> , 2Q, p <sub>1</sub>	0.958	1.022	1.080	1.127	1.161	1.179	1.182	1.169	1.140	1.098
OO	0.858	1.067	1.290	1.500	1.666	1.764	1.779	1.708	1.563	1.366
MK	0.987	1.014	1.037	1.054	1.065	1.071	1.072	1.068	1.059	1.044
2MK	1.000	1.015	1.025	1.030	1.032	1.032	1.032	1.032	1.031	1.027
Mf	0.907	1.044	1.181	1.300	1.390	1.442	1.450	1.413	1.335	1.224
Mm	1.043	1.000	0.957	0.919	0.891	0.874	0.872	0.884	0.908	0.943

\*Factor *f* of MS, 2SM, and MSf are each equal to factor *f* of M<sub>2</sub>. Factor *f* of P<sub>1</sub>, R<sub>2</sub>, S<sub>1</sub>, S<sub>2</sub>, S<sub>4</sub>, S<sub>6</sub>, T<sub>2</sub>, Sa, and Ssa are each unity.

Table 14.—*Node factor f for middle of each year, 1850 to 1999—Continued*

Constituent	1880	1881	1882	1883	1884	1885	1886	1887	1888	1889
J <sub>1</sub> .....	1.047	0.991	0.932	0.878	0.840	0.827	0.841	0.880	0.934	0.994
K <sub>1</sub> .....	1.027	0.988	0.949	0.914	0.890	0.882	0.891	0.915	0.950	0.990
K <sub>2</sub> .....	1.048	0.951	0.866	0.800	0.760	0.748	0.762	0.803	0.869	0.955
L <sub>2</sub> .....	1.246	1.020	0.786	0.944	1.152	1.171	1.006	0.824	0.945	1.205
M <sub>1</sub> .....	1.046	1.528	1.824	1.529	0.970	0.877	1.364	1.721	1.593	1.075
M <sub>2</sub> * N <sub>2</sub> , 2N, λ <sub>2</sub> , μ <sub>2</sub> , ν <sub>2</sub> .....	0.996	1.009	1.020	1.030	1.036	1.038	1.036	1.029	1.020	1.008
M <sub>3</sub> .....	0.994	1.013	1.031	1.045	1.054	1.057	1.054	1.044	1.030	1.012
M <sub>4</sub> , MN.....	0.992	1.017	1.041	1.060	1.073	1.077	1.072	1.060	1.040	1.016
M <sub>6</sub> .....	0.988	1.026	1.062	1.092	1.111	1.118	1.111	1.091	1.061	1.024
M <sub>8</sub> .....	0.984	1.035	1.084	1.124	1.151	1.160	1.150	1.123	1.082	1.033
O <sub>1</sub> , Q <sub>1</sub> , 2Q, p <sub>1</sub> .....	1.043	0.980	0.916	0.860	0.820	0.806	0.821	0.862	0.919	0.983
OO.....	1.144	0.926	0.739	0.599	0.513	0.486	0.516	0.604	0.746	0.936
MK.....	1.023	0.997	0.968	0.941	0.922	0.915	0.922	0.942	0.969	0.998
2MK.....	1.019	1.005	0.988	0.969	0.955	0.950	0.955	0.970	0.988	1.006
Mf.....	1.092	0.953	0.823	0.717	0.649	0.626	0.651	0.721	0.828	0.959
Mm.....	0.984	1.028	1.069	1.102	1.124	1.131	1.123	1.101	1.067	1.026
Constituent	1890	1891	1892	1893	1894	1895	1896	1897	1898	1899
J <sub>1</sub> .....	1.049	1.096	1.132	1.155	1.165	1.162	1.146	1.118	1.077	1.026
K <sub>1</sub> .....	1.028	1.062	1.088	1.105	1.112	1.110	1.099	1.078	1.054	1.012
K <sub>2</sub> .....	1.052	1.148	1.230	1.289	1.316	1.308	1.267	1.197	1.108	1.010
L <sub>2</sub> .....	1.153	0.709	0.683	1.185	1.219	0.704	0.607	1.141	1.229	0.897
M <sub>1</sub> .....	1.323	2.091	2.158	1.434	1.369	2.176	2.240	1.471	1.166	1.781
M <sub>2</sub> * N <sub>2</sub> , 2N, λ <sub>2</sub> , μ <sub>2</sub> , ν <sub>2</sub> .....	0.996	0.984	0.974	0.967	0.963	0.964	0.969	0.978	0.989	1.001
M <sub>3</sub> .....	0.993	0.976	0.961	0.950	0.946	0.947	0.954	0.967	0.983	1.002
M <sub>4</sub> , MN.....	0.991	0.968	0.948	0.934	0.928	0.930	0.939	0.956	0.977	1.002
M <sub>6</sub> .....	0.987	0.952	0.923	0.903	0.894	0.896	0.910	0.934	0.966	1.003
M <sub>8</sub> .....	0.982	0.936	0.898	0.873	0.861	0.864	0.882	0.913	0.955	1.004
O <sub>1</sub> , Q <sub>1</sub> , 2Q, p <sub>1</sub> .....	1.046	1.100	1.142	1.170	1.182	1.179	1.160	1.125	1.078	1.020
OO.....	1.153	1.375	1.571	1.713	1.780	1.761	1.660	1.491	1.281	1.058
MK.....	1.024	1.045	1.059	1.068	1.072	1.071	1.065	1.054	1.036	1.013
2MK.....	1.019	1.028	1.031	1.032	1.032	1.032	1.032	1.030	1.025	1.014
Mf.....	1.098	1.230	1.339	1.416	1.451	1.441	1.387	1.296	1.175	1.038
Mm.....	0.983	0.941	0.907	0.883	0.872	0.875	0.892	0.921	0.958	1.001
Constituent	1900	1901	1902	1903	1904	1905	1906	1907	1908	1909
J <sub>1</sub> .....	0.968	0.910	0.861	0.832	0.829	0.854	0.900	0.957	1.016	1.069
K <sub>1</sub> .....	0.973	0.934	0.903	0.885	0.883	0.899	0.928	0.965	1.005	1.042
K <sub>2</sub> .....	0.916	0.838	0.782	0.752	0.750	0.774	0.825	0.900	0.992	1.090
L <sub>2</sub> .....	0.753	1.030	1.193	1.117	0.925	0.858	1.051	1.221	1.062	0.653
M <sub>1</sub> .....	1.902	1.399	0.858	1.069	1.507	1.643	1.340	0.946	1.479	2.112
M <sub>2</sub> * N <sub>2</sub> , 2N, λ <sub>2</sub> , μ <sub>2</sub> , ν <sub>2</sub> .....	1.013	1.024	1.032	1.037	1.038	1.034	1.026	1.016	1.003	0.991
M <sub>3</sub> .....	1.020	1.036	1.049	1.056	1.057	1.051	1.039	1.023	1.005	0.986
M <sub>4</sub> , MN.....	1.027	1.049	1.066	1.076	1.076	1.068	1.053	1.031	1.007	0.982
M <sub>6</sub> .....	1.040	1.074	1.101	1.115	1.117	1.104	1.080	1.047	1.010	0.973
M <sub>8</sub> .....	1.054	1.100	1.136	1.157	1.159	1.142	1.108	1.063	1.013	0.964
O <sub>1</sub> , Q <sub>1</sub> , 2Q, p <sub>1</sub> .....	0.956	0.893	0.812	0.811	0.808	0.834	0.882	0.944	1.008	1.068
OO.....	0.850	0.679	0.559	0.496	0.490	0.543	0.652	0.814	1.017	1.240
MK.....	0.986	0.957	0.933	0.918	0.916	0.929	0.952	0.980	1.008	1.033
2MK.....	0.999	0.980	0.963	0.952	0.951	0.960	0.977	0.996	1.012	1.023
Mf.....	0.901	0.779	0.686	0.634	0.630	0.673	0.759	0.877	1.012	1.151
Mm.....	1.045	1.083	1.112	1.128	1.130	1.116	1.089	1.052	1.010	0.966

\*Factor f of MS, 2SM, and MSf are each equal to factor f of M<sub>2</sub>. Factor f of P<sub>1</sub>, R<sub>2</sub>, S<sub>1</sub>, S<sub>2</sub>, S<sub>3</sub>, S<sub>6</sub>, T<sub>2</sub>, Sa, and Ssa are each unity.

Table 14.—*Node factor f for middle of each year, 1850 to 1999—Continued*

Constituent	1910	1911	1912	1913	1914	1915	1916	1917	1918	1919
J <sub>1</sub> -----	1.111	1.142	1.160	1.165	1.157	1.137	1.104	1.059	1.004	0.945
K <sub>1</sub> -----	1.073	1.096	1.109	1.113	1.107	1.092	1.067	1.035	0.997	0.958
K <sub>2</sub> -----	1.182	1.256	1.303	1.317	1.296	1.243	1.165	1.071	0.973	0.884
L <sub>2</sub> -----	0.834	1.246	1.135	0.561	0.729	1.221	1.172	0.761	0.780	1.118
M <sub>1</sub> -----	1.972	1.248	1.557	2.297	2.146	1.310	1.371	1.993	1.909	1.243
M <sub>2*</sub> , N <sub>2</sub> , 2N, λ <sub>2</sub> , μ <sub>2</sub> , ν <sub>2</sub> -----	0.980	0.970	0.965	0.963	0.966	0.972	0.982	0.993	1.006	1.018
M <sub>3</sub> -----	0.969	0.956	0.948	0.945	0.949	0.958	0.972	0.990	1.009	1.027
M <sub>4</sub> , MN-----	0.959	0.942	0.931	0.928	0.933	0.945	0.964	0.987	1.012	1.036
M <sub>6</sub> -----	0.940	0.914	0.898	0.894	0.901	0.918	0.946	0.980	1.017	1.054
M <sub>8</sub> -----	0.920	0.887	0.867	0.861	0.870	0.893	0.928	0.973	1.023	1.073
O <sub>1</sub> , Q <sub>1</sub> , 2Q, ρ <sub>1</sub> -----	1.118	1.154	1.176	1.183	1.173	1.148	1.109	1.056	0.995	0.931
OO-----	1.455	1.633	1.748	1.783	1.732	1.602	1.414	1.195	0.974	0.779
MK-----	1.051	1.063	1.070	1.072	1.069	1.061	1.048	1.028	1.003	0.975
2MK-----	1.029	1.032	1.032	1.032	1.032	1.032	1.028	1.021	1.009	0.992
Mf-----	1.275	1.373	1.434	1.452	1.425	1.356	1.252	1.124	0.985	0.851
Mm-----	0.927	0.896	0.877	0.871	0.880	0.902	0.934	0.975	1.018	1.060
Constituent	1920	1921	1922	1923	1924	1925	1926	1927	1928	1929
J <sub>1</sub> -----	0.890	0.847	0.827	0.836	0.870	0.921	0.980	1.037	1.086	1.125
K <sub>1</sub> -----	0.921	0.894	0.882	0.887	0.909	0.942	0.981	1.020	1.055	1.083
K <sub>2</sub> -----	0.813	0.767	0.748	0.756	0.791	0.852	0.934	1.030	1.127	1.214
L <sub>2</sub> -----	1.198	1.034	0.870	0.932	1.133	1.199	0.963	0.669	0.975	1.270
M <sub>1</sub> -----	0.896	1.308	1.597	1.503	1.082	0.954	1.619	2.063	1.739	1.138
M <sub>2*</sub> , N <sub>2</sub> , 2N, λ <sub>2</sub> , μ <sub>2</sub> , ν <sub>2</sub> -----	1.028	1.035	1.038	1.036	1.031	1.022	1.011	0.998	0.986	0.976
M <sub>3</sub> -----	1.042	1.052	1.057	1.055	1.047	1.034	1.016	0.998	0.979	0.964
M <sub>4</sub> , MN-----	1.056	1.071	1.077	1.074	1.063	1.045	1.022	0.997	0.973	0.952
M <sub>6</sub> -----	1.086	1.108	1.118	1.114	1.096	1.068	1.033	0.995	0.959	0.929
M <sub>8</sub> -----	1.116	1.146	1.160	1.154	1.130	1.092	1.044	0.994	0.946	0.906
O <sub>1</sub> , Q <sub>1</sub> , 2Q, ρ <sub>1</sub> -----	0.871	0.827	0.806	0.815	0.850	0.905	0.968	1.032	1.088	1.134
OO-----	0.626	0.528	0.487	0.504	0.579	0.710	0.889	1.102	1.325	1.530
MK-----	0.947	0.925	0.915	0.920	0.937	0.963	0.992	1.018	1.040	1.056
2MK-----	0.973	0.958	0.950	0.953	0.966	0.984	1.002	1.017	1.026	1.031
Mf-----	0.730	0.660	0.626	0.641	0.701	0.801	0.928	1.066	1.201	1.317
Mm-----	1.096	1.120	1.131	1.126	1.107	1.076	1.036	0.993	0.950	0.914
Constituent	1930	1931	1932	1933	1934	1935	1936	1937	1938	1939
J <sub>1</sub> -----	1.150	1.163	1.164	1.151	1.126	1.088	1.038	0.982	0.923	0.871
K <sub>1</sub> -----	1.102	1.112	1.112	1.102	1.083	1.056	1.021	0.982	0.943	0.909
K <sub>2</sub> -----	1.278	1.312	1.313	1.279	1.216	1.130	1.033	0.937	0.854	0.792
L <sub>2</sub> -----	1.022	0.471	0.873	1.270	1.078	0.636	0.859	1.190	1.162	0.933
M <sub>1</sub> -----	1.312	2.353	1.992	1.197	1.614	2.148	1.850	1.098	1.079	1.534
M <sub>2*</sub> , N <sub>2</sub> , 2N, λ <sub>2</sub> , μ <sub>2</sub> , ν <sub>2</sub> -----	0.968	0.964	0.964	0.968	0.975	0.986	0.998	1.011	1.022	1.031
M <sub>3</sub> -----	0.952	0.946	0.946	0.952	0.963	0.979	0.997	1.016	1.033	1.047
M <sub>4</sub> , MN-----	0.937	0.929	0.929	0.936	0.951	0.972	0.996	1.021	1.044	1.063
M <sub>6</sub> -----	0.907	0.895	0.895	0.906	0.928	0.958	0.994	1.032	1.067	1.096
M <sub>8</sub> -----	0.878	0.863	0.862	0.877	0.905	0.945	0.992	1.043	1.091	1.130
O <sub>1</sub> , Q <sub>1</sub> , 2Q, ρ <sub>1</sub> -----	1.165	1.181	1.181	1.165	1.134	1.090	1.034	0.970	0.907	0.852
OO-----	1.686	1.772	1.773	1.690	1.535	1.332	1.108	0.894	0.714	0.581
MK-----	1.066	1.071	1.071	1.067	1.057	1.041	1.019	0.992	0.964	0.938
2MK-----	1.032	1.032	1.032	1.032	1.031	1.026	1.017	1.003	0.985	0.966
Mf-----	1.402	1.446	1.447	1.403	1.320	1.205	1.070	0.931	0.804	0.704
Mm-----	0.887	0.873	0.873	0.887	0.913	0.949	0.991	1.035	1.075	1.107

\* Factor *f* of MS, 2SM, and MSf are each equal to factor *f* of M<sub>2</sub>.  
Factor *f* of P<sub>1</sub>, R<sub>2</sub>, S<sub>1</sub>, S<sub>2</sub>, S<sub>4</sub>, S<sub>8</sub>, T<sub>2</sub>, Sa, and Ssa are each unity.

Table 14.—*Node factor f for middle of each year, 1850 to 1999—Continued*

Constituent	1940	1941	1942	1943	1944	1945	1946	1947	1948	1949
J <sub>1</sub> .....	0.836	0.827	0.846	0.888	0.944	1.003	1.057	1.103	1.136	1.157
K <sub>1</sub> .....	0.888	0.882	0.894	0.920	0.956	0.996	1.034	1.067	1.091	1.107
K <sub>2</sub> .....	0.757	0.748	0.766	0.812	0.882	0.970	1.068	1.162	1.242	1.295
L <sub>2</sub> .....	0.860	1.021	1.180	1.144	0.876	0.748	1.091	1.255	0.894	0.482
M <sub>1</sub> .....	1.623	1.313	0.879	1.076	1.714	1.944	1.480	1.138	1.927	2.339
M <sub>2</sub> * <sup>a</sup> , N <sub>2</sub> , 2N, λ <sub>2</sub> , μ <sub>2</sub> , ν <sub>2</sub> ..	1.036	1.038	1.035	1.028	1.018	1.006	0.994	0.982	0.972	0.966
M <sub>3</sub> .....	1.055	1.057	1.053	1.042	1.027	1.009	0.990	0.973	0.959	0.949
M <sub>4</sub> , MN.....	1.074	1.077	1.071	1.057	1.036	1.012	0.987	0.964	0.945	0.933
M <sub>6</sub> .....	1.113	1.118	1.108	1.086	1.055	1.018	0.981	0.947	0.919	0.901
M <sub>8</sub> .....	1.154	1.160	1.147	1.117	1.074	1.025	0.975	0.929	0.894	0.870
O <sub>1</sub> , Q <sub>1</sub> , 2Q, p <sub>1</sub> .....	0.816	0.806	0.826	0.870	0.929	0.994	1.055	1.107	1.147	1.173
OO.....	0.505	0.486	0.526	0.623	0.774	0.969	1.189	1.408	1.598	1.729
MK.....	0.920	0.915	0.925	0.946	0.974	1.002	1.028	1.047	1.061	1.069
2MK.....	0.953	0.950	0.957	0.973	0.991	1.008	1.021	1.028	1.032	1.032
Mf.....	0.642	0.626	0.659	0.736	0.848	0.981	1.120	1.249	1.354	1.424
Mm.....	1.126	1.131	1.121	1.096	1.061	1.019	0.976	0.935	0.902	0.880
Constituent	1950	1951	1952	1953	1954	1955	1956	1957	1958	1959
J <sub>1</sub> .....	1.165	1.160	1.143	1.112	1.070	1.002	0.959	0.901	0.855	0.829
K <sub>1</sub> .....	1.113	1.109	1.096	1.074	1.043	1.001	0.966	0.929	0.900	0.883
K <sub>2</sub> .....	1.317	1.303	1.257	1.184	1.092	0.995	0.903	0.827	0.776	0.749
L <sub>2</sub> .....	1.074	1.330	1.014	0.653	1.001	1.260	1.112	0.867	0.915	1.115
M <sub>1</sub> .....	1.717	1.120	1.778	2.161	1.664	0.964	1.276	1.656	1.527	1.053
M <sub>2</sub> * <sup>a</sup> , N <sub>2</sub> , 2N, λ <sub>2</sub> , μ <sub>2</sub> , ν <sub>2</sub> ..	0.963	0.965	0.970	0.979	0.990	1.003	1.015	1.026	1.033	1.038
M <sub>3</sub> .....	0.945	0.948	0.956	0.969	0.986	1.004	1.023	1.039	1.051	1.057
M <sub>4</sub> , MN.....	0.928	0.931	0.941	0.959	0.981	1.006	1.031	1.052	1.068	1.076
M <sub>6</sub> .....	0.894	0.898	0.914	0.939	0.972	1.009	1.046	1.079	1.104	1.116
M <sub>8</sub> .....	0.861	0.867	0.887	0.920	0.962	1.012	1.062	1.107	1.141	1.158
O <sub>1</sub> , Q <sub>1</sub> , 2Q, p <sub>1</sub> .....	1.183	1.177	1.155	1.119	1.069	1.010	0.945	0.884	0.835	0.808
OO.....	1.784	1.750	1.637	1.459	1.246	1.023	0.819	0.656	0.546	0.491
MK.....	1.072	1.070	1.063	1.051	1.033	1.009	0.981	0.953	0.930	0.916
2MK.....	1.032	1.032	1.032	1.029	1.023	1.012	0.996	0.977	0.960	0.951
Mf.....	1.452	1.435	1.375	1.278	1.154	1.016	0.880	0.761	0.675	0.630
Mm.....	0.872	0.877	0.896	0.926	0.965	1.008	1.051	1.088	1.116	1.130
Constituent	1960	1961	1962	1963	1964	1965	1966	1967	1968	1969
J <sub>1</sub> .....	0.831	0.860	0.909	0.766	1.025	1.076	1.117	1.146	1.161	1.165
K <sub>1</sub> .....	0.885	0.903	0.934	0.972	1.011	1.048	1.077	1.098	1.110	1.113
K <sub>2</sub> .....	0.752	0.781	0.836	0.914	1.008	1.106	1.195	1.265	1.307	1.316
L <sub>2</sub> .....	1.199	1.081	0.849	0.893	1.200	1.237	0.838	0.690	1.185	1.310
M <sub>1</sub> .....	0.767	1.197	1.690	1.699	1.166	1.175	1.976	2.175	1.503	1.197
M <sub>2</sub> * <sup>a</sup> , N <sub>2</sub> , 2N, λ <sub>2</sub> , μ <sub>2</sub> , ν <sub>2</sub> ..	1.037	1.033	1.024	1.014	1.001	0.989	0.978	0.969	0.964	0.963
M <sub>3</sub> .....	1.056	1.049	1.037	1.020	1.002	0.983	0.967	0.954	0.947	0.945
M <sub>4</sub> , MN.....	1.076	1.066	1.050	1.027	1.003	0.978	0.956	0.940	0.930	0.928
M <sub>6</sub> .....	1.116	1.111	1.075	1.041	1.004	0.967	0.935	0.911	0.897	0.894
M <sub>8</sub> .....	1.157	1.137	1.102	1.055	1.005	0.956	0.914	0.883	0.865	0.861
O <sub>1</sub> , Q <sub>1</sub> , 2Q, p <sub>1</sub> .....	0.810	0.840	0.891	0.954	1.018	1.076	1.124	1.159	1.178	1.182
OO.....	0.495	0.557	0.675	0.845	1.053	1.276	1.487	1.655	1.758	1.782
MK.....	0.917	0.932	0.956	0.985	1.013	1.036	1.053	1.064	1.071	1.072
2MK.....	0.952	0.962	0.980	0.998	1.014	1.024	1.030	1.032	1.032	1.032
Mf.....	0.633	0.684	0.776	0.898	1.035	1.172	1.293	1.385	1.439	1.451
Mm.....	1.128	1.113	1.084	1.046	1.003	0.959	0.922	0.893	0.876	0.872

\* Factor *f* of M<sub>8</sub>, 2SM, and MSf are each equal to factor *f* of M<sub>2</sub>.  
Factor *f* of P<sub>1</sub>, R<sub>2</sub>, S<sub>1</sub>, S<sub>2</sub>, S<sub>3</sub>, T<sub>2</sub>, Sa, and Ssa are each unity.

Table 14.—*Node factor f for middle of each year, 1850 to 1999—Continued*

Constituent	1970	1971	1972	1973	1974	1975	1976	1977	1978	1979
J <sub>1</sub> .....	1.155	1.132	1.097	1.051	0.995	0.936	0.881	0.842	0.827	0.839
K <sub>1</sub> .....	1.105	1.088	1.063	1.029	0.991	0.951	0.916	0.891	0.882	0.890
K <sub>2</sub> .....	1.289	1.232	1.150	1.055	0.957	0.871	0.804	0.763	0.748	0.760
L <sub>2</sub> .....	0.882	0.668	1.118	1.270	1.014	0.808	0.988	1.179	1.169	0.994
M <sub>1</sub> .....	1.987	2.176	1.503	1.012	1.535	1.777	1.428	0.870	0.874	1.361
M <sub>2</sub> * , N <sub>2</sub> , 2N, λ <sub>2</sub> , μ <sub>2</sub> , ν <sub>2</sub> .....	0.966	0.973	0.983	0.995	1.008	1.020	1.029	1.035	1.038	1.036
M <sub>3</sub> .....	0.950	0.960	0.975	0.993	1.012	1.029	1.044	1.054	1.057	1.054
M <sub>4</sub> , MN.....	0.934	0.948	0.967	0.991	1.016	1.039	1.059	1.072	1.077	1.073
M <sub>6</sub> .....	0.903	0.922	0.951	0.986	1.024	1.060	1.090	1.110	1.118	1.112
M <sub>8</sub> .....	0.873	0.898	0.935	0.981	1.032	1.081	1.122	1.149	1.160	1.151
O <sub>1</sub> , Q <sub>1</sub> , 2Q, p <sub>1</sub> .....	1.170	1.143	1.101	1.047	0.984	0.920	0.863	0.822	0.806	0.819
OO.....	1.716	1.575	1.380	1.159	0.940	0.750	0.607	0.517	0.485	0.512
MK.....	1.068	1.059	1.045	1.024	0.998	0.970	0.943	0.923	0.915	0.922
2MK.....	1.032	1.031	1.028	1.020	1.006	0.989	0.970	0.956	0.950	0.955
Mf.....	1.417	1.341	1.233	1.102	0.962	0.831	0.723	0.652	0.625	0.647
Mm.....	0.882	0.906	0.940	0.982	1.025	1.067	1.100	1.123	1.131	1.124
Constituent	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989
J <sub>1</sub> .....	0.877	0.930	0.989	1.045	1.093	1.130	1.153	1.164	1.163	1.148
K <sub>1</sub> .....	0.913	0.948	0.987	1.026	1.060	1.086	1.104	1.112	1.111	1.100
K <sub>2</sub> .....	0.799	0.864	0.949	1.045	1.142	1.226	1.285	1.315	1.310	1.270
L <sub>2</sub> .....	0.848	1.001	1.238	1.157	0.745	0.811	1.263	1.244	0.749	0.746
M <sub>1</sub> .....	1.656	1.468	0.974	1.323	2.050	2.032	1.292	1.367	2.142	2.122
M <sub>2</sub> * , N <sub>2</sub> , 2N, λ <sub>2</sub> , μ <sub>2</sub> , ν <sub>2</sub> .....	1.030	1.021	1.009	0.997	0.984	0.974	0.967	0.964	0.964	0.969
M <sub>3</sub> .....	1.045	1.031	1.013	0.994	0.977	0.962	0.951	0.946	0.947	0.954
M <sub>4</sub> , MN.....	1.061	1.042	1.018	0.993	0.969	0.949	0.935	0.928	0.930	0.939
M <sub>6</sub> .....	1.092	1.063	1.027	0.989	0.954	0.924	0.904	0.894	0.896	0.910
M <sub>8</sub> .....	1.125	1.085	1.036	0.986	0.939	0.901	0.874	0.862	0.864	0.881
O <sub>1</sub> , Q <sub>1</sub> , 2Q, p <sub>1</sub> .....	0.858	0.915	0.979	1.041	1.096	1.140	1.168	1.182	1.180	1.161
OO.....	0.596	0.735	0.921	1.137	1.361	1.560	1.706	1.778	1.766	1.668
MK.....	0.941	0.967	0.996	1.022	1.043	1.058	1.068	1.072	1.071	1.065
2MK.....	0.969	0.987	1.005	1.019	1.027	1.031	1.032	1.032	1.032	1.032
Mf.....	0.715	0.820	0.949	1.088	1.221	1.333	1.412	1.450	1.443	1.392
Mm.....	1.103	1.070	1.029	0.986	0.944	0.909	0.884	0.872	0.874	0.891
Constituent	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999
J <sub>1</sub> .....	1.120	1.080	1.030	0.972	0.914	0.864	0.833	0.829	0.852	0.896
K <sub>1</sub> .....	1.079	1.051	1.015	0.976	0.937	0.905	0.886	0.883	0.897	0.926
K <sub>2</sub> .....	1.203	1.115	1.016	0.922	0.842	0.785	0.754	0.750	0.772	0.821
L <sub>2</sub> .....	1.216	1.248	0.898	0.801	1.077	1.208	1.107	0.921	0.893	1.096
M <sub>1</sub> .....	1.334	1.156	1.778	1.829	1.282	0.800	1.083	1.487	1.560	1.214
M <sub>2</sub> * , N <sub>2</sub> , 2N, λ <sub>2</sub> , μ <sub>2</sub> , ν <sub>2</sub> .....	0.977	0.988	1.000	1.013	1.024	1.032	1.037	1.038	1.034	1.027
M <sub>3</sub> .....	0.966	0.982	1.000	1.019	1.036	1.048	1.056	1.057	1.051	1.040
M <sub>4</sub> , MN.....	0.955	0.976	1.000	1.025	1.048	1.065	1.075	1.076	1.069	1.054
M <sub>6</sub> .....	0.932	0.964	1.000	1.038	1.072	1.099	1.115	1.117	1.105	1.082
M <sub>8</sub> .....	0.911	0.952	1.000	1.051	1.098	1.134	1.156	1.159	1.143	1.111
O <sub>1</sub> , Q <sub>1</sub> , 2Q, p <sub>1</sub> .....	1.128	1.081	1.024	0.960	0.897	0.844	0.812	0.808	0.832	0.879
OO.....	1.505	1.296	1.072	0.863	0.688	0.565	0.498	0.489	0.538	0.643
MK.....	1.054	1.038	1.015	0.988	0.959	0.934	0.918	0.916	0.928	0.950
2MK.....	1.030	1.025	1.015	1.000	0.982	0.964	0.952	0.951	0.959	0.976
Mf.....	1.303	1.184	1.048	0.910	0.786	0.691	0.636	0.629	0.669	0.752
Mm.....	0.918	0.956	0.998	1.042	1.081	1.110	1.128	1.130	1.117	1.091

\* Factor *f* of MS, 2SM, and MSf are each equal to factor *f* of M<sub>2</sub>.  
Factor *f* of P<sub>1</sub>, R<sub>1</sub>, S<sub>1</sub>, S<sub>2</sub>, S<sub>4</sub>, S<sub>6</sub>, T<sub>2</sub>, Sa, and Ssa are each unity.

Table 15.—Equilibrium argument ( $V_o + u$ ) for meridian of Greenwich at beginning of each calendar year, 1850 to 2000

Constituent	1850	1851	1852	1853	1854	1855	1856	1857	1858	1859	1860	1861	1862	1863	1864	1865	1866	1867	1868	1869
J <sub>1</sub>	°	°	°	°	°	°	°	°	°	°	°	°	°	°	°	°	°	°	°	°
K <sub>1</sub>	29.7	116.0	204.0	307.5	38.1	129.4	221.2	327.3	59.4	151.2	242.6	347.4	77.0	165.2	251.7	350.6	74.0	157.0	240.8	340.1
K <sub>2</sub>	3.5	1.6	0.9	2.2	3.3	5.0	7.1	10.4	12.8	14.9	16.7	18.9	19.3	18.7	17.0	15.2	11.7	8.0	4.6	3.1
L <sub>2</sub>	187.8	183.8	184.0	186.0	189.5	194.0	200.9	205.8	210.4	214.0	218.2	220.0	217.0	213.2	209.7	203.2	196.1	190.0	187.0	187.0
M <sub>1</sub>	155.7	350.6	161.5	330.9	177.7	26.0	194.6	353.3	204.2	54.4	229.0	26.0	229.0	22.0	22.0	22.0	22.0	249.4	86.0	283.9
M <sub>1</sub>	6.8	273.5	165.2	50.6	346.9	274.8	170.8	55.4	347.2	281.7	178.9	61.0	341.0	280.0	179.7	65.8	311.8	235.5	162.6	47.2
M <sub>2</sub>	299.8	40.1	140.6	217.0	181.0	59.3	160.6	237.7	288.2	80.6	181.8	258.5	359.3	99.8	200.2	276.0	16.0	316.2	292.0	258.0
M <sub>3</sub>	269.7	240.2	211.0	145.6	317.0	89.8	61.0	356.6	328.6	300.9	271.2	207.7	178.8	99.7	120.2	53.9	24.0	354.1	324.5	324.5
M <sub>4</sub>	239.6	80.2	281.3	74.1	276.1	118.5	321.3	115.4	318.4	161.2	13.6	156.9	358.5	169.6	40.3	191.9	32.1	232.4	72.3	224.0
M <sub>6</sub>	179.4	120.3	61.9	291.1	234.1	177.8	121.0	353.5	297.6	241.7	185.4	155.4	239.7	240.5	77.0	48.1	348.4	288.5	156.0	144.7
N <sub>2</sub>	270.0	281.6	293.4	268.0	280.3	292.8	305.5	280.8	306.5	318.7	293.6	305.6	317.5	329.9	314.5	325.8	337.2	311.2	330.4	
N <sub>2</sub>	240.3	163.2	86.2	191.9	242.6	166.4	90.3	328.0	247.9	171.8	95.6	328.7	215.2	98.0	330.3	252.9	175.5	98.1	330.4	
O <sub>1</sub>	299.9	42.6	143.8	218.5	317.9	242.6	166.4	227.3	325.4	63.7	162.3	236.0	335.9	77.0	178.5	258.2	3.6	109.3	214.1	292.7
O <sub>0</sub>	240.0	132.5	23.8	318.0	223.1	129.6	37.5	333.4	242.0	150.0	56.8	348.9	250.7	41.8	317.2	201.3	84.1	329.1	245.8	
P <sub>1</sub>	349.7	349.9	350.2	349.4	349.7	349.9	350.2	319.4	349.6	349.9	350.1	349.4	349.6	349.8	349.3	349.6	349.8	350.1	349.3	
Q <sub>1</sub>	270.0	281.4	286.6	269.5	280.0	289.9	270.4	279.8	289.3	289.2	271.1	282.3	294.7	308.5	285.4	302.0	319.0	335.4	311.9	
Q <sub>2</sub>	240.4	165.6	89.4	320.9	242.0	163.4	84.3	313.8	234.1	154.9	76.1	306.2	228.2	75.2	312.6	240.5	168.8	96.4	331.0	
R <sub>2</sub>	179.9	179.7	179.4	180.2	179.9	179.6	179.4	180.1	179.8	179.6	179.3	180.1	179.8	179.6	179.3	180.0	179.8	179.3	180.0	
S <sub>1</sub>	180.0	180.0	180.0	180.0	180.0	180.0	180.0	180.0	180.0	180.0	180.0	180.0	180.0	180.0	180.0	180.0	180.0	180.0	180.0	
S <sub>0,48</sub>	0.1	0.3	0.6	0.359.8	0.1	0.4	0.6	0.359.9	0.2	0.4	0.7	0.359.9	0.2	0.4	0.7	0.0	0.2	0.5	0.0	0.0
T <sub>2</sub>	0.1	0.3	0.6	0.359.8	0.1	0.4	0.6	0.359.9	0.2	0.4	0.7	0.359.9	0.2	0.4	0.7	0.0	0.2	0.5	0.0	0.0
$\lambda_2$	-148.8	59.6	330.7	228.9	140.4	52.2	324.1	223.0	135.0	46.9	318.7	217.2	128.5	39.6	310.5	208.1	118.7	29.3	299.9	197.6
$p_2$	-241.0	82.1	283.4	76.2	277.0	119.9	322.0	115.5	317.7	159.8	1.8	154.9	356.4	197.7	38.8	191.0	33.4	22.9	312.4	226.4
$p_2$	-270.8	200.6	130.1	26.1	315.6	246.3	177.2	72.4	3.4	294.2	224.9	119.9	50.0	340.1	249.2	163.8	33.4	22.9	312.4	207.2
$p_1$	-270.9	203.9	133.8	26.1	315.3	243.4	171.1	62.0	349.6	277.3	205.4	97.3	26.7	317.2	249.2	146.1	80.9	16.2	310.7	207.2
MK	303.3	41.8	141.6	219.2	321.4	64.3	167.8	248.2	352.0	95.5	198.5	277.4	18.5	118.5	217.2	291.2	27.7	124.0	220.8	295.1
2MK	236.1	78.6	280.4	71.9	272.8	113.5	314.1	105.0	305.6	146.2	346.9	138.1	339.2	181.0	23.3	176.7	20.4	224.4	67.7	220.9
MM	206.8	321.8	74.1	125.1	238.1	352.1	106.1	158.5	272.7	80.6	181.3	258.6	359.3	99.8	200.2	276.0	16.0	116.1	193.3	245.2
MS	249.8	40.1	140.6	217.0	318.0	59.3	160.6	237.7	339.7	115.5	317.7	227.6	140.5	192.0	319.1	330.5	81.9	232.6	73.4	225.6
2SM	60.2	319.9	219.4	143.0	42.0	300.7	199.4	122.3	20.8	279.4	178.2	101.5	0.7	260.2	159.8	84.0	344.0	243.9	143.8	68.0
ML	240.0	134.9	33.0	220.1	222.7	126.7	300.7	191.5	323.1	228.3	133.2	37.3	326.4	227.4	125.8	21.1	299.4	188.8	327.3	246.5
MF	60.2	319.8	143.0	42.0	300.7	199.4	122.3	20.8	279.4	178.2	101.5	0.7	260.2	159.8	84.0	344.0	243.9	143.8	68.0	
Mm	20.8	118.5	207.2	309.0	37.7	126.4	216.2	316.9	45.7	134.4	223.1	324.9	53.6	142.3	322.1	332.8	61.6	150.3	239.0	340.8
Sa	280.3	280.1	279.8	280.6	280.3	280.1	279.8	280.6	280.4	280.1	279.9	280.6	280.4	280.2	280.7	280.4	280.2	279.9	280.7	280.4
Ssa	200.6	200.1	199.6	201.1	200.7	200.2	199.7	201.2	200.7	200.2	199.8	201.3	200.3	200.8	200.3	199.8	201.3	200.8	199.9	201.4

Table 15.—*Equilibrium argument ( $V_o + u$ ) for meridian of Greenwich at beginning of each calendar year, 1850 to 2000*—Con.

Constituent	1870	1871	1872	1873	1874	1875	1876	1877	1878	1879	1880	1881	1882	1883	1884	1885	1886	1887	1888	1889
$J_1$	°	°	°	°	°	°	°	°	°	°	°	°	°	°	°	°	°	°	°	°
$K_1$	67.1	155.7	245.6	350.5	456.3	562.1	674.0	786.9	900.0	1013.9	112.2	123.9	134.6	145.3	156.0	166.7	177.4	188.1	190.8	194.4
$K_2$	1.7	1.4	2.0	4.4	6.3	8.5	10.9	14.1	16.2	17.7	18.6	19.5	20.4	21.3	22.2	23.1	24.0	1.9	2.0	
$L_1$	183.7	182.2	188.2	192.1	196.9	201.8	208.6	212.9	216.0	217.6	219.2	216.7	212.5	206.2	201.5	194.8	188.9	184.4	184.0	
$L_2$	271.7	100.6	138.4	129.4	206.9	121.8	310.8	331.4	151.9	159.2	180.8	184.1	174.7	204.3	201.2	211.7	219.8	235.4	235.3	
$M_1$	297.1	201.0	145.3	48.0	301.8	202.5	145.5	55.4	310.0	207.8	142.2	58.2	312.8	203.4	108.2	31.1	301.4	192.6	83.7	344.5
$M_2$	32.4	133.0	233.9	310.6	51.9	153.3	254.8	321.8	73.2	174.4	275.3	351.6	92.1	192.4	292.5	8.2	108.2	308.6	24.7	
$M_3$	228.6	199.5	170.8	165.9	77.8	30.0	22.2	317.8	289.8	291.5	232.9	167.4	138.1	108.5	78.7	12.2	342.3	312.5	282.9	
$M_4$	64.8	266.0	211.8	261.6	107.8	103.8	149.6	273.5	146.4	219.6	163.0	105.9	334.3	8.8	24.7	225.0	163.6	210.4	257.2	
$M_6$	97.2	39.0	331.6	211.2	155.6	99.9	44.3	275.6	219.9	274.7	292.7	337.4	21.2	326.5	8.3	49.4	89.9	32.6	72.8	113.3
$M_8$	126.6	172.1	215.5	162.4	207.5	253.2	299.1	247.4	292.7	292.7	337.4	217.4	157.4	217.1	24.3	44.3	205.8	72.1	98.8	98.8
$N_2$	322.9	334.8	366.9	321.8	324.4	347.1	359.8	335.1	347.8	1.2	12.4	347.0	358.7	10.2	21.6	355.5	6.9	18.3	4.1	
$2N$	34.8	135.6	256.5	308.8	47.3	145.5	243.6	316.4	104.9	180.0	109.5	326.3	262.3	186.0	109.5	342.3	265.5	188.2	111.0	
$O_1$	140.3	39.3	302.0	234.8	141.9	50.2	318.8	224.6	162.2	68.3	54.7	153.5	252.8	327.8	69.4	172.5	277.2	357.5	103.1	207.6
$O_0$	349.6	349.8	350.0	349.3	349.5	349.8	350.0	349.2	349.5	349.0	349.2	349.5	349.0	349.2	349.7	349.9	349.7	349.9	343.6	
$P_1$	325.3	337.3	348.3	320.0	323.8	339.3	348.7	319.6	329.3	323.9	323.9	323.9	323.0	323.2	323.2	323.2	323.2	323.2	323.2	
$Q_1$	225.8	161.3	331.3	252.3	173.1	93.7	322.9	243.9	105.2	87.1	318.5	212.6	168.3	95.5	332.2	260.4	187.5	113.1	345.7	
$2Q$	179.7	179.5	180.0	179.7	179.4	179.2	179.0	179.7	179.4	179.2	179.7	179.4	179.2	179.6	179.1	179.8	179.6	179.3	179.8	
$R_2$	180.0	180.0	180.0	180.0	180.0	180.0	180.0	180.0	180.0	180.0	180.0	180.0	180.0	180.0	180.0	180.0	180.0	180.0	180.0	
$S_1$	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
$S_2$	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
$T_2$	0.3	0.0	0.5	0.0	0.3	0.0	0.6	0.0	0.1	0.3	0.0	0.6	0.0	0.1	0.4	0.0	0.2	0.4	0.0	
$\lambda_2$	108.5	19.6	291.0	189.6	101.4	13.3	285.3	184.2	96.1	7.8	279.2	177.4	88.4	359.2	269.8	167.4	77.9	348.6	259.4	157.3
$\mu_2$	66.5	208.2	109.8	104.9	104.9	104.9	104.9	104.9	104.9	104.9	104.9	104.9	104.9	104.9	104.9	104.9	104.9	104.9	104.9	
$p_2$	136.3	66.4	256.8	251.6	182.4	113.3	44.2	290.5	230.3	160.9	91.3	345.8	275.8	205.5	135.1	217.1	158.0	224.1	177.8	72.1
$P_K$	34.1	134.4	235.9	315.0	58.2	161.8	265.6	346.0	89.3	192.1	233.9	11.3	110.7	208.9	305.9	18.9	115.3	212.3	310.6	26.7
$2MK$	63.1	264.6	105.7	256.8	97.5	298.1	138.7	289.6	130.2	174.5	287.7	338.6	90.8	202.6	323.6	165.5	8.2	211.5	52.7	255.3
$MN$	355.3	107.8	220.8	272.4	25.3	140.4	254.6	307.0	60.9	174.5	275.3	351.6	92.1	192.4	292.5	5.6	209.3	115.1	226.6	38.8
$MS$	322.4	133.0	233.3	310.6	51.9	153.3	254.8	321.8	73.2	174.4	286.8	185.6	64.7	267.9	167.6	67.5	351.8	251.8	151.7	355.3
$MS'$	327.6	227.0	126.1	49.4	308.1	206.7	105.2	23.2	260.8	184.7	192.4	278.7	323.6	102.1	193.2	227.2	82.1	333.0	150.9	
$ML$	147.2	41.9	303.4	233.0	137.3	42.3	307.6	239.1	143.7	47.4	309.6	236.0	133.3	27.3	278.1	193.2	82.1	333.0	227.2	
$MS'$	327.6	227.0	126.1	49.4	308.1	206.7	105.2	23.2	260.8	184.7	192.4	278.7	323.6	102.1	193.2	227.2	82.1	333.0	150.9	
$Mm$	69.5	158.2	247.0	348.6	77.5	166.2	254.9	356.7	85.4	174.2	262.9	4.7	93.4	182.1	278.8	12.6	101.3	190.1	278.8	20.6
$Sa$	280.4	280.2	280.0	280.7	280.5	280.2	280.0	280.8	280.5	280.3	280.0	280.8	280.5	280.3	280.1	280.8	280.6	280.3	280.8	
$Ss$	200.9	200.4	200.0	201.4	201.0	200.5	200.0	201.5	200.6	201.0	200.5	200.6	201.1	200.6	201.1	200.6	200.7	200.7	201.7	

Table 15.—Equilibrium argument ( $V_o + u$ ) for meridian of Greenwich at beginning of each calendar year, 1850 to 2000—Con.

Constituent	1890	1891	1892	1893	1894	1895	1896	1897	1898	1899	1900	1901	1902	1903	1904	1905	1906	1907	1908	1909
J <sub>1</sub>	°	°	°	°	°	°	°	°	°	°	°	°	°	°	°	°	°	°	°	°
J <sub>2</sub>	107.6	107.6	107.9	289.0	34.8	126.9	219.0	310.9	56.4	147.3	237.2	325.7	52.7	137.8	221.5	304.6	42.1	127.0	213.6	301.8
K <sub>1</sub>	3.0	3.0	4.6	7.6	9.9	12.2	14.7	17.3	18.6	19.2	18.9	17.5	14.9	11.6	7.8	5.3	2.6	0.9	0.4	1.8
K <sub>2</sub>	183.8	185.4	188.6	194.8	199.7	204.7	209.4	215.2	217.8	217.5	214.2	209.0	202.7	195.9	191.4	186.0	182.3	180.7	183.2	182.5
L <sub>2</sub>	65.8	66.3	69.0	94.0	295.3	86.6	268.4	118.4	317.1	127.1	309.0	147.1	345.0	174.1	339.6	162.5	1.2	204.8	17.5	90.6
M <sub>1</sub>	287.7	192.3	86.3	310.4	288.6	199.8	93.4	344.8	289.6	206.1	97.5	551.6	271.7	203.3	101.6	338.1	232.1	158.1	90.6	336.2
M <sub>2</sub>	125.4	226.4	327.6	47.5	146.0	247.4	348.9	65.8	166.8	267.7	8.3	108.7	208.9	49.0	124.7	47.9	325.3	65.8	125.4	
M <sub>3</sub>	188.1	159.6	131.3	66.8	39.0	11.2	333.3	278.6	250.5	221.6	192.5	163.0	133.5	103.5	73.6	49.1	337.4	278.7	213.4	
M <sub>4</sub>	250.8	92.8	80.0	291.9	134.9	337.7	131.5	333.7	175.4	16.6	217.4	57.8	258.0	98.1	249.4	86.8	290.5	131.6	284.5	
M <sub>6</sub>	16.5	319.1	262.7	77.9	230.2	178.1	226.3	197.3	140.5	88.3	24.9	326.0	207.0	147.1	142.2	326.5	197.5	66.7	20.0	
M <sub>8</sub>	141.7	165.6	230.2	178.1	226.3	80.8	315.4	263.0	307.4	350.8	33.2	74.8	115.6	156.0	196.1	138.9	179.6	221.0	263.3	
N <sub>2</sub>	16.1	28.4	40.8	16.0	28.7	41.5	54.2	29.3	41.6	53.8	65.7	77.3	88.8	100.2	111.5	85.4	96.8	108.5	120.3	95.0
N <sub>2N</sub>	266.8	190.3	114.1	347.5	271.5	194.5	119.5	352.8	276.4	199.8	213.0	251.4	174.0	46.1	328.8	174.8	47.7	174.8	144.0	
O <sub>0</sub>	127.2	226.5	325.2	38.2	136.4	234.4	332.7	45.8	144.5	244.5	345.2	345.2	87.4	161.2	296.3	42.0	122.0	328.5	69.3	
O <sub>0</sub>	49.4	313.2	219.4	164.4	62.8	331.5	239.7	174.1	79.4	342.0	240.9	135.2	249.3	24.3	269.2	152.0	68.5	311.8	206.1	133.5
P <sub>1</sub>	319.4	349.6	349.4	349.1	349.4	349.6	349.8	349.3	349.6	349.8	350.0	350.3	350.8	350.5	350.0	350.3	350.5	350.7	350.0	
Q <sub>2</sub>	17.9	28.5	38.5	9.7	19.1	28.5	38.0	9.3	19.6	30.5	42.6	56.0	71.1	87.5	104.5	82.6	98.8	111.7	124.0	
R <sub>3</sub>	179.6	179.3	178.0	179.8	179.5	179.3	179.0	179.7	179.5	179.2	178.7	178.4	178.2	177.9	178.7	178.4	178.2	177.9	178.6	
S <sub>1</sub>	180.0	180.0	180.0	180.0	180.0	180.0	180.0	180.0	180.0	180.0	180.0	180.0	180.0	180.0	180.0	180.0	180.0	180.0	180.0	
S <sub>2</sub>	0.4	0.4	0.7	1.0	0.2	0.5	0.7	0.1	0.3	0.5	0.8	1.0	1.3	1.6	1.8	2.1	1.3	1.6	2.1	1.4
X <sub>2</sub>	68.6	340.1	251.8	150.6	62.5	334.5	246.5	145.2	56.8	328.2	239.4	150.3	61.0	331.6	212.2	139.7	50.4	321.3	232.4	130.6
A <sub>2</sub>	253.0	304.7	296.6	89.9	292.2	134.4	336.5	129.8	331.6	173.3	14.6	215.8	66.7	257.6	98.4	260.4	91.4	292.5	133.8	286.6
P <sub>1</sub>	4.0	2.3	292.8	221.0	111.2	39.8	327.4	256.1	146.3	74.8	4.0	294.2	225.8	159.1	93.7	28.9	220.5	152.2	82.9	335.6
M <sub>K</sub>	127.5	229.4	332.1	52.1	155.9	259.7	3.3	83.1	185.5	287.0	27.2	126.2	223.8	320.6	56.8	130.0	227.5	323.2	144.1	
M <sub>MN</sub>	141.6	226.4	327.6	44.5	146.0	247.4	348.9	65.8	166.8	267.7	8.3	108.7	208.9	309.0	49.0	120.5	210.1	186.2	237.2	
M <sub>MS</sub>	125.4	226.4	327.6	44.5	146.0	247.4	348.9	65.8	166.8	267.7	8.3	108.7	208.9	309.0	49.0	124.7	224.9	325.3	65.8	
P <sub>2M</sub>	234.6	133.6	32.4	315.5	214.0	112.6	11.1	294.2	193.2	92.3	351.7	261.3	151.1	51.0	311.0	235.3	135.1	34.7	294.2	217.8
M <sub>F</sub>	51.1	313.3	217.1	148.1	53.2	223.5	154.2	57.3	318.8	217.8	113.9	6.6	256.4	145.0	60.8	311.0	232.7	134.3	34.3	
M <sub>Si</sub>	234.6	133.6	32.4	315.5	214.0	112.6	11.1	294.2	193.2	92.3	351.7	261.3	151.1	51.0	311.0	235.3	135.1	34.7	294.2	217.8
M <sub>Mm</sub>	109.3	186.0	286.7	28.5	117.2	206.0	294.7	36.5	125.2	213.9	302.6	31.4	120.1	208.8	287.5	39.3	128.0	216.8	305.6	
S <sub>a</sub>	230.6	280.4	280.1	280.9	280.6	280.4	280.9	280.7	280.4	280.2	280.0	279.7	279.5	280.0	279.7	279.5	279.3	280.0	279.3	
S <sub>sa</sub>	201.2	200.7	200.3	201.3	200.8	200.3	201.8	200.3	200.9	200.4	199.9	199.4	199.5	200.0	199.5	199.5	199.0	198.5	200.0	

Table 15.—*Equilibrium argument ( $V_0 + u$ ) for meridian at Greenwich at beginning of each calendar year, 1850 to 2000—Con.*

Constituent	1910	1911	1912	1913	1914	1915	1916	1917	1918	1919	1920	1921	1922	1923	1924	1925	1926	1927	1928	1929
J <sub>1</sub>	◦	◦	◦	◦	◦	◦	◦	◦	◦	◦	◦	◦	◦	◦	◦	◦	◦	◦	◦	◦
K <sub>1</sub>	136.2	227.6	319.5	65.6	167.7	249.5	340.8	85.4	174.8	262.7	349.0	87.6	170.9	253.9	337.9	77.4	164.7	253.6	343.7	88.7
K <sub>2</sub>	3.0	4.8	7.0	10.3	12.6	14.7	16.4	18.5	18.7	16.0	14.2	10.6	6.9	3.6	2.3	1.1	0.9	1.7	4.2	
K <sub>3</sub>	186.5	186.1	183.7	200.6	205.6	210.4	217.4	215.5	211.4	207.1	201.0	194.3	188.1	185.3	182.9	181.4	182.9	182.9	182.9	
L <sub>2</sub>	179.7	22.0	235.0	65.4	206.6	46.0	260.2	81.1	245.3	75.8	277.1	10.6	284.2	101.7	291.2	119.9	139.1	133.4	136.7	141.4
M <sub>1</sub>	229.6	146.2	92.1	343.4	237.0	149.2	350.1	242.0	142.6	78.4	344.0	236.0	124.6	25.1	309.9	226.5	120.0	15.3	282.0	
M <sub>2</sub>	243.3	344.5	85.9	163.0	264.5	9.5	107.7	183.5	284.4	25.0	217.9	187.7	121.6	91.7	41.2	141.3	58.2	159.1	253.9	
M <sub>3</sub>	184.9	156.8	128.9	64.5	36.7	8.8	340.6	275.5	246.7	217.5	187.7	121.6	91.7	61.1	31.9	235.7	238.7	173.8		
M <sub>4</sub>	126.6	323.1	171.9	169.0	11.7	214.2	7.4	208.9	49.9	250.6	42.1	242.2	82.3	282.5	74.2	275.1	116.4	318.2	111.7	
M <sub>5</sub>	9.9	238.2	257.8	129.1	73.3	17.7	210.2	191.1	133.8	74.9	141.1	84.2	183.3	123.5	63.3	291.4	232.6	174.6	117.3	
M <sub>6</sub>	253.2	343.8	292.1	343.8	23.4	68.3	14.8	57.8	99.9	141.1	84.2	124.4	164.6	205.1	148.5	100.2	232.8	276.5	223.4	
N <sub>2</sub>	107.3	119.8	132.5	107.8	120.5	133.2	145.7	120.5	132.5	144.3	155.9	129.9	142.1	152.6	164.0	138.0	149.7	161.7	173.8	
2N	331.3	255.1	170.0	52.6	336.6	260.5	184.3	57.3	340.6	263.7	186.6	58.8	341.4	264.0	186.6	68.9	341.9	265.1	188.6	
O <sub>0</sub>	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
O <sub>1</sub>	297.4	204.2	112.2	48.2	316.7	224.6	131.1	62.8	324.1	221.3	113.6	28.2	271.9	154.9	40.4	317.9	303.6	61.1	204.2	
P <sub>1</sub>	350.2	350.5	350.7	350.0	350.4	350.7	349.9	350.2	350.4	350.6	349.9	350.1	350.4	350.6	349.9	350.1	350.3	350.6	349.8	
Q <sub>1</sub>	107.1	116.9	126.4	97.4	106.8	116.4	123.3	98.3	109.7	109.7	122.3	136.3	130.2	147.2	163.4	139.7	152.8	164.6	175.4	
Q <sub>2</sub>	331.1	252.2	173.0	42.2	322.8	243.7	164.9	35.5	317.8	241.6	167.0	42.4	258.6	186.3	60.6	345.6	190.2	147.1		
R <sub>2</sub>	178.4	178.1	177.9	178.6	178.3	178.1	177.8	178.3	178.0	177.8	178.3	178.0	177.8	178.3	178.0	177.8	178.2	178.7	178.4	
S <sub>1</sub>	180.0	180.0	180.0	180.0	180.0	180.0	180.0	180.0	180.0	180.0	180.0	180.0	180.0	180.0	180.0	180.0	180.0	180.0	180.0	
S <sub>2</sub> , <sub>3</sub> , <sub>6</sub>	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
T <sub>2</sub>	1.6	1.9	2.1	1.4	1.7	2.2	1.8	2.2	1.4	1.7	2.0	2.2	1.0	1.7	2.0	2.2	1.5	2.0	1.6	
A <sub>2</sub>	42.2	314.0	225.9	124.8	36.8	308.7	220.5	118.9	30.2	301.2	212.1	109.7	20.3	290.8	201.5	99.2	10.1	281.3	192.8	
p <sub>1</sub>	128.4	330.4	172.5	328.6	168.2	10.3	212.3	5.3	206.8	48.1	249.1	41.3	242.1	82.9	283.7	76.0	277.1	118.6	320.2	
v <sub>2</sub>	264.4	195.1	126.0	21.2	312.2	243.0	173.7	68.5	358.7	288.7	218.5	112.4	41.9	331.5	261.0	155.1	85.0	151.1	200.4	
MK	246.3	349.4	92.9	173.3	277.1	20.6	123.5	202.2	303.2	43.0	141.4	215.2	311.7	48.0	144.9	219.4	318.6	59.2	160.8	
MK	123.6	324.3	164.9	315.8	166.4	357.0	197.7	348.9	190.2	32.0	234.5	27.9	231.6	75.4	278.9	72.0	274.0	115.6	316.5	
MN	243.3	344.5	85.9	163.0	264.5	5.9	107.1	183.7	284.4	50.0	125.3	335.0	234.7	193.7	305.2	355.2	107.3	333.0	219.9	
MN	116.7	15.5	274.1	197.0	95.5	354.1	252.9	176.3	75.6	335.0	159.0	58.9	318.8	218.7	142.9	42.4	301.8	124.1		
2SM	297.1	201.3	106.2	37.8	303.0	207.8	111.7	40.6	301.2	199.2	94.0	11.8	260.9	149.5	39.9	319.6	115.9	17.7	307.5	
MSI	116.7	116.7	15.5	274.1	197.0	95.5	354.1	252.9	176.3	75.6	335.0	159.0	58.9	318.8	218.7	142.9	42.4	301.8	124.1	
MM	136.0	136.0	136.0	136.0	136.0	136.0	136.0	136.0	136.0	136.0	136.0	136.0	136.0	136.0	136.0	136.0	136.0	136.0	136.0	
S <sub>3</sub>	279.8	279.5	280.0	270.8	270.6	270.3	280.1	270.8	270.6	270.3	270.8	270.6	270.3	270.8	270.4	280.1	270.9	270.4	280.2	
S <sub>3a</sub>	199.6	199.1	200.1	199.6	199.1	200.0	199.6	199.1	199.7	200.2	198.7	199.2	199.7	200.2	198.7	200.3	198.8	199.3	200.3	

Table 15.—Equilibrium argument ( $V_0 + u$ ) for meridian of Greenwich at beginning of each calendar year, 1850 to 2000—Con.

Constituent	1930	1931	1932	1933	1934	1935	1936	1937	1938	1939	1940	1941	1942	1943	1944	1945	1946	1947	1948	1949
$J_1$	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
$K_1$	-180.4	272.3	4.4	110.5	202.1	238.1	23.2	126.2	213.5	299.1	23.1	120.2	203.5	288.0	0	0	0	205.5	296.0	27.3
$K_2$	-6.1	8.4	10.7	14.0	15.5	19.4	18.2	19.1	17.9	15.6	12.4	9.6	6.0	3.1	1.4	1.2	116.2	105.5	133.1	
$L_2$	191.8	196.6	201.6	208.3	212.4	215.3	216.6	217.9	215.1	210.3	204.2	199.6	192.7	187.0	182.9	182.8	184.8	188.2	194.6	
$M_1$	352.2	170.5	330.8	210.6	167.4	18.8	201.7	6.0	193.1	35.9	227.4	215.6	249.6	82.0	147.3	118.7	126.2	114.4	127.7	
$M_2$	228.3	21.0	293.3	234.4	134.1	27.0	255.7	132.6	227.4	132.2	220.0	260.9	173.4	109.5	12.9	252.0	151.1	91.3	14.5	
$M_3$	337.2	78.6	180.1	257.1	358.4	99.6	200.5	276.8	17.2	117.5	217.6	293.2	33.3	133.4	233.8	306.9	50.6	151.6	252.8	
$M_4$	145.8	117.9	90.1	154.3	156.9	169.2	44.0	133.6	34.4	234.9	75.1	226.5	66.6	260.9	250.0	254.8	227.4	109.2	134.7	
$M_5$	314.3	157.2	0.2	154.3	355.4	356.9	169.8	241.5	110.3	51.6	352.4	292.7	159.7	44.3	201.5	230.6	239.6	145.7	299.6	
$M_6$	291.5	25.8	180.2	51.4	355.4	356.8	208.8	241.5	109.8	109.8	150.3	92.9	133.2	215.0	159.5	202.5	246.5	291.3	239.3	
$M_8$	268.7	314.4	0.3	308.6	333.8	338.4	82.0	27.1	68.9	109.8	150.3	133.2	173.8	215.0	159.5	202.5	246.5	291.3	239.3	
$N_2$	161.4	174.1	186.9	162.1	174.7	187.1	199.3	173.8	185.5	197.1	208.4	182.3	193.6	205.1	276.7	190.6	191.7	227.8	203.0	
$N_3$	345.6	166.6	183.6	67.1	271.0	198.2	309.0	353.2	276.6	199.2	208.4	182.3	216.2	210.7	278.0	222.8	227.0	202.8	203.0	
$O_1$	332.6	70.8	168.9	241.7	340.1	274.1	178.4	233.6	355.8	98.8	203.6	284.0	23.5	133.8	236.6	312.6	52.7	151.9	250.6	
$O_2$	216.6	124.9	33.6	329.2	297.7	142.4	45.7	329.9	228.3	118.6	4.2	274.6	158.2	297.5	279.0	222.0	123.2	27.4	238.9	
$P_1$	350.1	350.3	350.6	349.8	350.0	350.5	349.8	350.0	350.2	350.7	349.8	350.0	350.5	349.7	350.2	350.5	350.2	350.4	349.7	
$Q_1$	156.8	166.3	175.6	146.7	156.4	166.5	177.2	130.6	163.7	178.4	194.5	173.1	80.5	219.5	205.1	215.6	225.6	196.7	196.9	
$Q_2$	341.0	261.8	182.4	51.7	332.6	254.0	176.1	47.6	332.0	258.0	185.4	62.1	350.2	277.1	202.5	74.9	357.5	279.3	200.5	
$R_2$	178.2	177.9	177.7	178.4	178.2	177.9	177.6	178.4	178.1	177.9	178.3	178.1	177.8	177.6	178.3	178.3	177.8	177.5	178.3	
$S_1$	180.0	180.0	180.0	180.0	180.0	180.0	180.0	180.0	180.0	180.0	180.0	180.0	180.0	180.0	180.0	180.0	180.0	180.0	180.0	
$S_2, 4, 6$	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
$T_2$	1.8	2.1	2.3	1.6	1.8	2.1	2.4	1.6	1.9	2.1	2.4	1.6	1.9	2.1	2.4	1.7	2.0	2.2	1.7	
$\lambda_2$	3.2	275.1	187.1	86.0	357.9	269.5	181.0	79.1	350.0	260.8	171.4	69.0	339.5	250.2	161.0	59.0	330.3	241.8	153.6	
$\mu_2$	315.6	157.6	359.8	153.2	353.3	248.2	179.0	109.6	40.0	294.5	224.4	154.1	83.7	337.5	267.0	196.7	126.4	207.7	111.0	
$\nu_2$	131.2	62.1	353.0	348.3	300.0	248.2	179.0	109.6	40.0	294.5	224.4	154.1	83.7	337.5	267.0	196.7	126.4	207.7	111.0	
$\rho_1$	126.6	54.3	341.8	232.8	160.7	89.0	18.0	271.3	202.6	135.4	69.8	328.3	263.2	197.1	129.4	23.4	313.0	241.8	169.9	
$MK$	343.3	87.0	190.8	271.1	14.4	171.0	218.7	205.8	35.1	133.0	229.9	302.8	39.3	136.5	234.9	311.3	52.2	154.3	237.3	
$2MK$	308.2	148.8	349.5	140.3	341.0	181.8	22.8	174.5	20.0	174.5	219.4	62.8	216.8	60.6	263.8	288.4	99.6	300.6	141.3	
$MN$	138.6	252.7	7.0	59.3	173.2	286.7	39.8	90.6	202.8	314.5	66.0	115.6	226.9	338.5	90.4	140.9	253.7	6.9	120.6	
$MS$	337.2	78.6	180.1	257.1	358.4	99.6	200.5	276.8	17.2	117.5	217.6	293.2	33.3	133.4	233.8	309.9	50.6	151.6	232.8	
$2SM$	22.8	281.4	179.9	102.9	358.1	159.5	83.2	342.8	242.5	142.4	66.8	326.7	226.6	126.2	50.1	309.4	208.4	107.2	30.2	
$Mf$	212.0	117.1	22.3	313.8	218.3	121.8	23.6	309.7	206.4	99.3	350.3	265.3	154.4	45.8	300.4	224.7	125.2	27.7	201.6	
$Ms$	22.8	281.4	179.9	102.9	1.6	260.4	180.1	159.5	83.2	342.8	242.5	142.4	66.8	326.7	226.6	126.2	50.1	309.4	208.4	
$Mm$	175.8	264.5	353.2	95.0	183.7	272.4	1.2	103.0	191.7	280.4	9.1	110.9	199.6	288.4	17.1	118.9	207.6	226.3	25.0	
$Sa$	279.9	279.7	279.4	280.2	280.0	279.7	279.5	280.2	280.0	279.8	280.3	280.0	279.8	279.5	280.3	280.0	279.8	279.6	280.3	
$Ss$	199.9	199.4	188.9	198.4	199.4	200.4	199.9	199.4	199.0	200.5	200.0	199.5	199.0	200.5	200.0	199.5	199.1	200.6	199.6	

Table 15.—*Equilibrium argument ( $V_0 + u$ ) for meridian of Greenwich at beginning of each calendar year, 1850 to 2000—Con.*

Constituent	1950	1951	1952	1953	1954	1955	1956	1957	1958	1959	1960	1961	1962	1963	1964	1965	1966	1967	1968	1969
$J_1$	◦	◦	◦	◦	◦	◦	◦	◦	◦	◦	◦	◦	◦	◦	◦	◦	◦	◦	◦	◦
$K_2$	225.2	317.2	49.1	154.6	245.3	355.0	63.3	164.0	249.0	332.5	55.5	163.2	2.7	0.9	325.2	53.7	157.6	248.4	340.0	71.9
$K_2'$	9.8	12.1	14.2	17.0	18.3	18.7	18.2	17.6	14.9	11.4	7.7	5.3	2.7	0.9	2.4	3.8	5.6	7.8	11.2	
$K_2''$	169.5	204.4	209.0	214.7	217.0	216.0	214.4	209.0	202.6	195.8	191.1	186.2	182.8	181.6	184.4	186.9	190.8	195.4	202.4	
$L_2$	86.6	298.5	147.4	306.3	118.5	322.8	165.4	338.9	153.8	342.4	179.5	5.4	191.0	2.4	194.4	30.4	230.4	26.6	215.8	57.9
$M_1$	165.3	91.6	22.2	265.5	159.0	85.3	21.9	266.9	155.7	52.4	341.3	252.2	146.5	36.5	302.6	147.6	40.3	302.7	237.8	
$M_1'$	71.3	172.8	274.4	351.0	92.1	192.9	293.0	9.4	109.6	209.7	309.8	25.5	125.7	43.1	144.2	42.5	346.9	64.0		
$M_3$	166.9	70.2	51.3	346.6	318.1	299.4	260.5	194.2	164.6	134.6	104.6	38.2	8.5	339.1	310.0	244.7	216.3	188.2	160.3	
$M_4$	152.6	345.5	188.3	312.1	184.2	255.8	227.0	18.9	219.3	58.4	259.5	50.9	251.3	92.1	235.3	86.2	298.4	150.9	323.8	
$M_6$	213.8	158.3	102.5	333.6	276.3	218.7	160.4	38.4	328.3	76.4	208.3	76.4	101.8	141.8	129.4	72.6	16.4	320.7	101.9	
$M_8$	211.1	16.7	324.2	8.4	51.6	31.9	37.8	78.6	118.9	169.9	101.8	142.6	184.4	226.6	172.5	216.8	21.6	307.6	255.9	
$N_2'$	215.7	228.5	241.2	216.3	228.6	240.7	252.5	226.7	238.2	249.5	260.9	234.8	246.2	257.9	244.5	256.8	269.4	282.1	257.4	
$N_2''$	0.0	0.2	284.2	208.2	81.5	285.5	211.6	84.0	6.7	289.4	212.0	84.1	6.8	289.8	213.0	85.8	9.5	263.2	217.3	
$O_0$	0.0	0.0	159.8	238.0	331.2	70.3	170.1	271.1	313.1	234.4	92.2	197.4	236.3	23.0	329.0	44.1	143.1	241.6	339.8	
$O_0'$	137.6	46.2	314.2	248.5	153.4	165.3	4.1	122.8	2.2	7.1	250.0	162.0	51.1	305.2	204.0	133.8	39.0	306.1	214.3	
$P_1$	349.9	350.9	235.0	346.6	349.9	350.1	350.4	349.6	349.9	350.1	350.3	327.7	254.3	247.4	245.5	255.8	275.5	246.0		
$Q_1$	206.1	215.5	225.0	196.4	206.8	217.9	230.2	220.7	220.4	220.7	220.4	223.3	247.4	260.8	245.5	286.8	289.4	210.2	79.4	
$Q_2$	330.6	271.3	192.0	61.6	343.3	245.5	189.2	62.7	349.6	277.1	205.4	81.6	292.7	216.1	186.1	86.8	8.4	294.7	177.8	
$R_2$	178.0	177.7	178.2	178.0	177.7	177.4	178.2	177.9	177.7	177.4	178.1	177.9	177.6	177.4	178.1	177.8	177.6	177.3	178.1	
$S_1$	180.0	180.0	180.0	180.0	180.0	180.0	180.0	180.0	180.0	180.0	180.0	180.0	180.0	180.0	180.0	180.0	180.0	180.0	180.0	
$S_{1/4}, 6$	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
$T_2$	0.0	2.0	2.3	2.5	1.8	2.0	2.3	2.6	1.8	2.1	2.3	2.6	1.9	2.1	2.4	2.6	1.9	2.2	2.4	
$N_3$	324.4	236.4	148.3	47.0	318.6	229.9	141.0	38.8	308.5	220.1	130.7	28.2	299.0	9.1	212.0	19.3	290.9	202.7	114.7	
$A_2$	142.6	344.9	187.1	340.3	182.1	23.7	225.0	17.4	218.3	59.1	259.9	52.0	253.0	94.1	265.4	88.3	246.9	216.1	334.3	
$p_2$	348.6	289.2	276.2	238.9	95.3	333.1	233.8	243.6	138.8	72.3	302.2	200.2	133.2	65.1	325.4	247.9	246.9	104.1	32.0	282.9
$M_K$	81.0	184.9	288.4	8.1	110.4	211.6	311.7	208.7	1.3	204.4	48.0	251.8	45.7	248.6	90.8	222.4	83.8	284.6	125.3	326.0
$2MK$	132.8	333.5	115.1	45.6	335.9	266.0	160.1	89.8	19.3	308.8	202.7	112.4	62.2	323.3	246.9	177.5	108.2	39.1	220.4	
$MN$	71.3	172.8	274.2	351.0	92.1	182.9	293.5	9.4	109.6	236.7	210.6	210.6	11.9	260.2	112.4	226.0	41.0	154.8	114.5	
$MS$	285.7	187.2	85.8	9.0	267.9	167.1	66.5	350.6	250.4	150.3	30.2	334.5	124.3	134.0	33.3	216.9	116.8	321.4		
$2SM$	184.0	184.9	288.4	8.1	110.4	211.6	311.7	208.7	1.3	204.4	48.0	251.8	45.7	248.6	90.8	222.4	83.8	284.6	125.3	326.0
$Mf$	128.0	33.2	298.1	228.6	131.6	32.7	291.3	213.1	105.3	354.8	243.4	52.1	308.1	207.1	134.8	38.0	302.3	207.2	138.9	
$Mst$	215.6	304.3	33.0	134.8	223.5	312.2	41.0	142.7	231.1	48.0	9.9	150.7	239.1	134.0	33.3	216.9	114.5	13.1	206.0	
$Mm$	280.1	279.8	280.4	280.1	279.9	279.6	279.6	279.6	280.4	280.1	280.0	279.7	280.4	280.4	280.4	280.4	280.4	280.4	280.4	
$Sg$	200.2	199.7	200.2	199.2	199.2	200.2	199.3	199.3	200.3	199.3	200.3	199.3	200.3	199.3	200.3	199.3	200.4	199.4	200.4	
$Ssa$	200.2	199.7	200.2	199.2	199.2	200.2	199.3	199.3	200.3	199.3	200.3	199.3	200.3	199.3	200.3	199.3	200.4	199.4	200.4	

Table 15.—Equilibrium argument ( $V_o + u$ ) for meridian of Greenwich at beginning of each calendar year, 1850 to 2000—Con.

Constituent	1970	1971	1972	1973	1974	1975	1976	1977	1978	1979	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989
$J_1$	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
$K_1$	-270.0	1.8	93.1	197.4	286.6	14.3	100.2	198.6	281.8	4.9	89.1	188.9	276.5	5.6	95.8	201.0	292.7	24.7	116.8	222.8
$K_2$	-13.4	15.5	17.1	19.0	19.5	218.4	218.2	211.5	14.1	10.5	6.8	2.5	1.5	1.5	2.4	4.9	9.2	11.6	14.8	o
$L_1$	-207.2	211.5	214.8	218.4	218.2	215.8	211.5	207.6	200.9	194.2	188.2	185.6	183.1	182.6	184.2	189.3	193.5	198.3	203.3	210.0
$L_2$	-262.8	60.3	255.2	81.8	282.4	99.0	276.0	99.7	297.9	130.9	310.2	116.6	313.4	169.2	350.3	135.8	334.5	188.0	25.5	166.0
$M_1$	-155.2	46.8	307.6	234.9	158.7	51.8	302.5	198.6	135.7	291.5	170.8	41.8	30.0	291.6	171.7	82.0	31.2	299.1	179.8	o
$M_2$	-165.4	266.8	8.0	84.5	185.3	285.8	26.0	101.8	201.8	301.9	42.0	117.9	218.4	319.1	60.0	136.8	238.1	339.6	81.0	158.1
$M_3$	-68.1	40.2	11.9	306.8	277.9	28.6	219.0	152.7	122.7	92.8	63.0	356.8	327.6	298.6	270.0	205.2	177.2	149.3	121.6	57.1
$M_4$	-330.9	173.6	15.9	169.1	51.5	211.5	52.1	203.6	43.5	243.8	84.0	235.8	76.7	278.1	120.0	273.6	219.1	223.1	162.1	316.2
$M_5$	-136.3	80.4	23.9	166.6	195.8	137.2	78.1	305.3	245.5	185.7	126.0	353.7	25.5	237.2	180.0	50.3	364.3	298.7	233.1	114.2
$M_6$	-301.7	347.1	31.8	338.2	21.0	63.0	104.1	47.1	87.3	127.5	168.0	111.6	153.4	196.2	240.0	187.1	232.4	278.2	324.1	272.3
$N_2$	-270.1	298.7	295.2	270.0	282.0	295.3	305.3	297.3	290.6	301.9	313.3	287.4	299.2	311.2	323.4	298.4	311.0	323.7	336.4	311.7
$O_1$	-14.8	298.2	222.4	95.4	18.7	301.8	224.6	296.8	19.4	296.9	41.6	119.4	221.1	321.4	60.8	134.2	322.6	330.7	68.8	141.6
$O_2$	-150.7	249.0	347.8	61.7	162.0	263.6	6.5	85.7	191.3	296.9	41.6	119.4	221.1	321.4	60.8	134.2	322.6	330.7	68.8	141.6
$O_3$	-58.7	326.4	232.6	163.9	64.6	321.1	212.6	126.5	9.8	253.0	139.2	57.5	313.6	213.9	117.6	51.0	138.5	227.0	135.6	71.2
$P_1$	-349.8	350.0	350.2	349.5	349.7	350.0	350.2	349.5	349.0	349.4	349.7	349.9	349.5	350.2	349.4	349.6	349.9	350.1	349.4	
$Q_1$	-255.4	265.0	275.0	247.2	258.7	271.5	280.5	263.3	280.7	297.0	312.9	289.0	301.9	313.5	324.2	295.8	314.8	324.2	295.8	
$Q_2$	-0.1	281.0	262.2	72.6	355.4	279.5	205.1	80.7	8.8	297.0	224.3	98.5	227.7	227.5	97.4	18.3	299.0	219.6	58.9	o
$R_2$	-177.8	177.6	177.3	178.0	177.8	177.5	178.0	177.7	177.5	177.2	178.0	177.7	177.5	177.2	177.4	177.9	177.7	177.4	177.2	177.9
$S_1$	-180.0	180.0	180.0	180.0	180.0	180.0	180.0	180.0	180.0	180.0	180.0	180.0	180.0	180.0	180.0	180.0	180.0	180.0	180.0	
$S_2$	-0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
$S_{p, 4, 6}$	-2.2	2.4	2.7	2.0	2.2	2.5	2.7	2.0	2.3	2.0	2.5	2.0	2.3	2.0	2.6	2.0	2.3	2.0	2.6	2.1
$T_2$	-225.3	172.1	146.0	341.5	86.8	341.5	271.7	201.7	131.4	25.3	314.9	244.4	174.0	68.0	358.0	288.2	218.6	113.5	44.3	335.2
$\lambda_3$	-285.6	197.4	109.1	7.6	278.8	189.8	100.6	358.2	268.8	179.4	90.0	347.8	258.8	170.0	81.4	340.0	251.9	163.9	75.9	334.8
$\mu_3$	-330.0	156.1	14.0	271.7	8.4	299.7	50.7	423.6	43.6	237.6	78.8	280.2	121.9	275.1	131.4	319.4	121.7	227.8	172.8	
$\rho_1$	-210.6	138.4	66.6	318.7	248.4	179.4	111.9	9.2	304.3	239.4	173.6	69.6	0.7	290.5	219.4	110.9	38.8	326.4	294.0	
$M_K$	-178.9	282.3	25.1	103.6	204.4	304.0	42.2	115.9	212.3	308.7	45.7	120.4	210.8	320.5	62.4	141.7	245.0	348.8	92.6	172.8
$2MK$	-317.4	158.1	358.8	150.0	351.4	193.3	35.9	189.4	33.2	237.0	80.4	233.3	75.3	276.6	117.6	268.6	109.3	309.9	130.5	301.9
$MN$	-75.6	189.5	303.8	354.5	107.2	219.5	331.3	21.1	132.4	243.8	355.3	45.3	157.5	242.2	23.4	75.1	185.1	303.8	57.5	109.8
$MS$	-165.4	266.8	8.0	185.3	255.8	26.0	101.8	201.8	301.9	308.7	42.0	117.9	218.4	319.1	60.0	136.8	238.1	339.6	81.0	158.1
$2SM$	-194.6	93.2	332.0	275.5	174.7	74.2	334.0	258.2	158.2	58.1	318.0	242.1	141.6	40.9	300.0	223.2	121.9	20.4	279.0	201.9
$M_L$	-44.0	308.7	212.4	141.1	41.3	286.8	193.0	110.4	359.3	248.0	138.8	59.0	316.2	216.2	118.4	48.4	313.0	218.1	123.4	54.8
$M_{SF}$	-194.6	93.2	332.0	275.5	174.7	74.2	334.0	258.2	158.2	58.1	318.0	242.1	141.6	40.9	300.0	223.2	121.9	20.4	279.0	201.9
$M_m$	-255.3	344.0	272.8	174.6	263.3	352.0	80.7	182.5	271.2	0.0	88.7	190.5	7.9	96.6	198.4	287.1	15.9	104.6	206.4	
$S_a$	-280.2	280.0	279.8	280.5	280.3	280.0	279.8	280.5	280.3	280.1	279.8	280.6	280.1	279.8	280.4	280.1	279.9	280.6	200.2	200.7
$S_{sa}$	-200.5	200.0	199.5	201.0	200.5	200.0	199.5	201.0	200.5	200.0	199.6	201.1	200.6	200.1	199.7	201.2	200.6	200.2	199.8	201.3

Table 15.—*Equilibrium argument ( $V_o + u$ ) for meridian of Greenwich at beginning of each calendar year, 1850 to 2000—Con.*

Constituent	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000
J <sub>1</sub>	°	°	°	°	°	°	°	°	°	°	°
K <sub>1</sub>	314.4	45.2	135.2	237.9	325.0	50.2	134.0	231.1	322.5	39.3	125.8
K <sub>2</sub>	16.7	18.0	18.7	19.4	18.0	15.6	12.2	9.5	6.0	3.2	1.5
L <sub>2</sub>	213.9	216.6	217.6	218.3	215.4	210.3	204.0	199.2	192.7	187.2	183.4
M <sub>1</sub>	2.2	2.2	4.9	4.9	30.4	229.4	66.7	242.7	59.3	244.2	83.7
M <sub>2</sub>	85.9	33.5	305.0	184.1	79.3	4.9	293.5	176.9	64.8	319.9	251.4
M <sub>3</sub>	259.4	0.5	101.3	177.6	278.0	18.2	118.3	194.0	294.0	34.2	134.5
M <sub>4</sub>	158.7	0.9	202.7	266.4	237.0	207.7	177.4	110.9	81.0	51.3	21.8
M <sub>6</sub>	58.1	1.4	304.0	172.8	114.0	54.6	356.9	221.9	162.1	102.6	43.6
M <sub>8</sub>	317.5	1.9	45.3	350.5	72.8	113.2	55.5	96.1	136.8	178.1	
N <sub>2</sub>	324.3	336.7	348.8	323.3	334.9	346.4	357.8	331.7	83.0	354.5	6.1
N <sub>2N</sub>	229.2	312.8	236.3	108.9	31.9	314.6	237.3	109.4	32.0	314.7	237.6
O <sub>1</sub>	240.1	338.0	78.7	154.0	269.7	369.7	104.8	185.2	290.5	34.6	137.2
O <sub>0</sub>	338.4	243.8	146.6	73.2	327.7	217.2	102.3	12.5	256.5	144.4	37.4
P <sub>1</sub>	349.6	349.8	350.1	349.3	349.6	349.8	350.1	349.3	349.6	349.8	350.0
Q <sub>1</sub>	305.0	315.2	326.1	293.7	313.0	327.9	344.3	322.9	339.5	354.9	8.8
2Q	9.9	291.4	213.6	85.4	10.0	296.2	223.8	100.6	28.5	316.0	240.3
R <sub>2</sub>	177.6	177.4	177.1	177.6	177.3	177.3	177.3	177.8	177.5	177.3	177.0
S <sub>1</sub>	180.0	180.0	180.0	180.0	180.0	180.0	180.0	180.0	180.0	180.0	180.0
S <sub>2</sub> , 4, 6	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
T <sub>2</sub>	2.4	2.6	2.9	2.2	2.4	2.4	2.7	2.9	2.2	2.5	3.0
$\lambda_2$	246.6	158.2	69.6	327.7	238.6	149.4	60.0	317.5	228.1	138.8	49.6
$\mu_2$	157.1	358.1	200.5	353.2	194.3	35.5	236.2	28.2	229.0	69.9	271.6
$p_2$	92.2	22.7	313.1	207.5	137.4	67.0	356.6	250.4	180.0	106.6	39.4
$p_1$	72.9	1.3	290.4	183.9	115.4	48.6	343.1	241.6	176.5	110.1	42.1
MK	276.0	18.5	120.0	197.0	296.0	33.8	130.5	203.4	300.0	37.4	136.0
2MK	142.1	342.9	184.0	335.8	177.9	20.8	224.4	18.4	222.1	65.2	207.6
MN	223.6	337.1	90.1	140.8	252.9	4.6	116.1	165.6	277.0	28.7	140.6
MS	259.4	0.5	101.3	177.6	278.0	18.2	118.3	194.0	294.0	34.2	134.6
2SM	100.6	359.5	258.7	182.4	303.8	31.8	120.5	222.3	311.0	35.7	225.5
M <sub>6</sub>	319.2	222.4	124.0	49.6	305.9	198.8	88.8	3.7	253.0	144.9	40.1
M <sub>8</sub>	100.6	359.5	258.7	182.4	303.8	31.8	120.5	222.3	311.0	35.7	225.5
M <sub>m</sub>	295.1	23.8	112.5	214.3	308.3	31.8	120.5	222.3	311.0	35.7	225.5
S <sub>8</sub>	280.4	280.2	279.9	280.4	279.9	280.4	280.4	280.4	280.4	280.4	280.0
S <sub>8a</sub>	200.8	200.3	199.8	201.3	200.7	200.8	200.4	201.4	200.4	200.4	200.0

Table 16.—*Differences to adapt table 15 to beginning of each calendar month*

Constituent	Month of year*											
	Jan.	Feb.	Mar.	Apr.	May	June	July	Aug.	Sept.	Oct.	Nov.	Dec.
J <sub>1</sub>	○	○	○	○	○	○	○	○	○	○	○	○
K <sub>1</sub>	0.00	75.57	108.99	184.56	246.08	321.65	23.17	98.74	174.31	255.82	311.39	12.91
K <sub>2</sub>	0.00	30.66	58.15	88.71	118.28	178.40	208.96	239.51	269.08	299.64	329.21	299.21
L <sub>2</sub>	0.00	61.11	116.31	177.42	236.56	356.66	57.91	178.16	238.27	288.41	288.41	288.41
M <sub>1</sub> †	0.00	9.19	52.33	61.54	82.02	91.21	111.71	120.90	130.09	150.69	159.75	180.29
M <sub>1</sub>	0.00	345.54	7.32	342.86	320.48	336.02	321.64	319.18	304.72	302.94	287.88	285.50
M <sub>1</sub>	0.00	342.09	0.75	342.83	337.11	319.20	313.47	295.56	277.65	271.93	254.01	248.59
M <sub>2</sub>	0.00	324.17	1.49	321.66	314.22	278.39	266.95	231.12	185.85	148.02	136.58	136.58
M <sub>3</sub>	0.00	306.26	2.24	291.33	267.59	240.42	166.68	162.94	93.78	42.04	296.05	273.16
M <sub>4</sub>	0.00	288.35	2.98	291.33	268.44	196.79	173.90	102.24	30.59	7.70	296.05	273.16
M <sub>6</sub>	0.00	252.52	4.48	257.00	222.66	115.18	80.86	333.37	232.89	191.55	84.07	49.74
M <sub>8</sub>	0.00	216.69	5.97	222.66	176.88	33.58	347.80	204.49	61.18	15.40	232.10	186.32
N <sub>2</sub>	0.00	279.16	310.66	223.82	186.42	105.58	62.18	341.34	260.50	217.11	136.27	92.87
N <sub>2</sub>	0.00	234.14	259.82	133.97	58.62	217.42	91.51	235.71	250.36	124.51	49.16	49.16
O <sub>1</sub>	0.00	293.62	303.34	236.96	195.94	129.56	88.55	22.16	315.78	274.77	208.39	167.37
O <sub>0</sub>	0.00	127.49	172.97	300.46	40.61	168.10	268.26	163.24	263.39	30.89	131.04	131.04
P <sub>1</sub>	0.00	329.44	301.85	271.29	241.72	211.17	181.60	151.04	120.49	90.92	60.36	30.79
Q <sub>1</sub>	0.00	248.60	252.50	141.11	68.14	316.75	182.39	20.99	306.63	196.63	123.67	123.67
Q <sub>2</sub>	0.00	203.59	201.67	45.26	300.34	143.93	39.02	232.61	86.20	341.28	184.87	184.87
R <sub>2</sub>	0.00	30.66	58.15	88.71	118.28	148.83	178.40	208.96	239.51	269.08	299.21	299.21
S <sub>4</sub>	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
T <sub>2</sub>	0.00	329.44	301.85	271.29	241.72	211.17	181.60	151.04	120.49	90.92	60.36	30.79
X <sub>2</sub>	0.00	314.98	309.16	264.15	232.20	187.19	155.24	110.22	65.21	33.26	316.24	316.24
μ <sub>2</sub>	0.00	288.35	302.98	291.33	268.44	196.79	173.90	102.24	7.70	296.05	273.16	273.16
ν <sub>2</sub>	0.00	333.36	305.83	271.18	36.24	18.66	352.02	325.38	334.44	307.61	316.87	316.87
ρ <sub>1</sub>	0.00	302.81	355.66	285.47	277.96	200.26	143.07	86.87	65.36	8.17	347.66	347.66
ρ <sub>2</sub>	0.00	354.73	59.64	54.37	72.50	67.23	85.35	80.05	74.81	92.83	87.66	87.66
M <sub>K</sub>	0.00	257.79	304.83	202.62	150.16	47.96	365.50	253.29	161.08	98.62	356.41	303.95
M <sub>N</sub>	0.00	243.33	312.15	195.48	140.64	23.97	329.13	212.47	95.80	40.96	284.29	229.45
M <sub>S</sub>	0.00	324.17	1.49	314.22	265.66	278.39	266.95	231.12	195.30	183.85	148.02	136.58
S <sub>M</sub>	0.00	35.83	358.51	34.34	45.78	81.61	93.05	128.88	164.70	176.15	211.98	223.42
M <sub>f</sub>	0.00	96.94	114.82	211.75	282.34	19.27	89.86	186.79	283.73	364.31	91.25	161.83
M <sub>IS</sub>	0.00	36.83	45.02	50.84	95.85	127.80	204.76	128.88	164.70	176.15	211.98	223.42
M <sub>M</sub>	0.00	30.56	58.15	88.71	118.28	148.83	178.40	208.96	236.74	269.79	11.76	43.71
S <sub>a</sub>	0.00	61.11	116.31	177.42	236.56	267.66	356.80	356.80	239.51	289.08	298.64	329.21
S <sub>ss</sub>	0.00											298.41

\*This table was designed for direct use for common years. For a leap year the values given for the months of March to December, inclusive, apply to the last day of the preceding month, but may be used directly, provided an allowance is made in the day of month as indicated in the following table.

†The first line for constituent M<sub>1</sub> gives the difference as based upon the formula in table 2; the second line gives the differences as derived from the half speed of constituent M<sub>2</sub>.

Table 17.—*Differences to adapt table 15 to beginning of each day of month*

Constituent	Day of month*										
	1	2	3	4	5	6	7	8	9	10	11
J <sub>1</sub>	°	°	°	°	°	°	°	°	°	°	°
K <sub>1</sub>	0.00	14.05	28.10	42.15	56.20	70.25	84.30	98.35	112.41	126.46	140.51
K <sub>2</sub>	0.00	0.99	5.91	2.96	3.94	5.93	6.90	7.88	8.87	9.86	19.71
L <sub>2</sub>	0.00	1.97	3.94	5.91	7.88	9.86	11.83	13.80	15.77	17.74	246.84
M <sub>2</sub>	0.00	337.37	326.05	314.73	303.42	292.10	280.78	269.47	258.15	251.37	239.11
M <sub>3</sub>	0.00	335.84	323.76	311.68	299.60	287.52	275.44	263.37	251.29	240.28	238.09
M <sub>4</sub>	0.00	347.92	335.81	323.43	311.24	299.05	286.86	274.66	262.47	250.28	240.51
M <sub>5</sub>	0.00	347.81	335.62	311.24	286.86	262.47	231.71	189.33	164.95	140.57	354.85
M <sub>6</sub>	0.00	323.43	286.86	250.28	213.71	177.14	140.57	103.99	67.42	30.85	232.37
M <sub>7</sub>	0.00	311.24	262.47	213.71	164.95	116.18	67.42	18.66	326.90	281.13	346.56
M <sub>8</sub>	0.00	286.86	213.71	140.57	67.42	354.28	281.13	207.99	134.84	61.70	104.74
N <sub>2</sub>	0.00	262.47	184.95	67.42	329.90	232.37	134.84	37.32	259.79	202.27	220.98
N <sub>3</sub>	0.00	322.55	285.11	247.66	210.21	165.32	107.44	56.93	6.42	315.91	265.44
N <sub>4</sub>	0.00	309.49	268.98	208.47	167.95	107.44	56.93	10.43	214.89	106.33	214.89
O <sub>1</sub>	0.00	334.63	309.27	283.90	258.63	233.16	202.43	157.06	131.70	242.44	229.35
O <sub>2</sub>	0.00	27.34	54.68	82.02	109.35	136.69	164.03	191.37	218.71	246.05	273.35
P <sub>1</sub>	0.00	-0.99	-1.97	-2.96	-3.94	-5.91	-6.90	-7.88	-8.87	-9.86	335.08
Q <sub>1</sub>	0.00	321.57	283.14	244.70	206.27	167.84	129.41	90.98	52.54	14.11	205.03
2Q	0.00	308.50	257.01	205.51	164.01	102.61	51.02	359.52	308.02	266.53	205.03
R <sub>2</sub>	0.00	0.99	1.97	2.96	3.94	4.93	5.91	6.90	7.88	8.87	9.86
S <sub>1</sub>	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
T <sub>2</sub>	0.00	-0.99	-1.97	-2.96	-3.94	-5.91	-6.90	-7.88	-8.87	-9.86	-10.85
X <sub>3</sub>	0.00	346.93	333.87	307.74	294.68	281.61	265.48	242.44	229.35	213.71	205.03
Y <sub>2</sub>	0.00	311.24	262.47	213.71	164.95	116.18	67.42	18.66	326.90	281.13	232.37
Z <sub>2</sub>	0.00	324.30	288.60	232.91	217.21	176.58	139.90	103.21	66.53	29.85	353.16
M <sub>1</sub>	0.00	323.32	286.63	249.95	213.27	176.58	139.90	103.21	66.53	29.85	126.04
M <sub>2</sub>	0.00	336.60	313.21	288.81	266.42	243.02	219.62	172.83	149.44	122.51	222.56
M <sub>3</sub>	0.00	310.25	260.50	210.75	161.00	111.26	61.51	11.76	322.01	225.38	163.55
M <sub>4</sub>	0.00	298.17	266.34	174.52	112.69	50.86	349.03	287.20	189.33	164.95	101.72
M <sub>5</sub>	0.00	335.62	311.24	286.86	262.47	238.09	213.71	170.67	195.05	219.43	116.18
2S <sub>M</sub>	0.00	24.38	48.76	73.14	97.63	121.91	146.29	184.47	210.82	237.18	243.82
M <sub>6</sub>	0.00	26.35	52.71	79.06	105.41	131.76	158.12	121.91	146.29	170.67	219.43
M <sub>7</sub>	0.00	24.38	48.76	73.14	97.63	121.91	146.29	184.47	210.82	237.18	243.82
M <sub>8</sub>	0.00	13.07	26.13	39.20	52.26	65.32	78.39	91.46	104.52	117.58	130.65
S <sub>8</sub>	0.00	0.99	1.97	2.96	3.94	5.91	6.90	7.88	8.87	9.86	19.71
S <sub>8a</sub>	0.00	1.97	3.94	5.91	7.88	11.83	13.80	15.77	17.74	19.71	19.71

\*The table is adapted directly for use with common years, but if the required date falls between Mar. 1 and Dec. 31, inclusive, in a leap year the day of month should be increased by one before entering the table.

†The first line for constituent M<sub>1</sub> gives the differences as based upon the formula in table 2, the second line gives the differences as derived from the half speed of constituent M<sub>2</sub>.

‡The differences for constituents S<sub>1</sub>, S<sub>2</sub>, S<sub>3</sub>, etc., are each zero for the beginning of every day.

Table 17.—*Differences to adapt table 15 to beginning of each day of month—Continued*

Constituent	Day of month*										
	12	13	14	15	16	17	18	19	20	21	22
J	°	°	°	°	°	°	°	°	°	°	°
J1	154.56	168.61	182.66	196.71	210.76	224.81	238.86	252.91	266.96	281.01	295.06
E1	10.84	11.83	12.81	13.80	14.78	15.77	16.76	17.74	18.73	19.71	20.70
E2	21.68	23.66	25.63	27.61	29.57	31.51	33.48	35.48	38.43	41.40	44.35
I1	235.52	224.20	212.88	201.57	190.25	178.94	167.62	156.30	144.99	133.67	122.35
M1†	227.13	215.05	202.97	190.89	178.81	166.73	154.65	142.57	130.49	118.41	106.33
M2	225.90	213.74	201.52	189.33	177.14	164.95	152.76	140.57	128.38	116.19	103.99
N1	91.80	67.47	43.04	18.66	354.96	329.90	305.92	281.13	266.75	232.37	207.99
N2	317.80	281.13	244.66	207.99	171.42	134.84	98.27	61.70	25.13	111.98	348.56
M3	183.61	134.84	86.08	37.32	348.56	299.79	251.03	202.27	153.50	104.74	55.98
M4	275.41	202.27	129.12	52.83	123.83	102.69	166.54	200.26	337.11	263.97	200.48
M5	7.7	269.69	172.16	74.64	337.11	299.58	142.06	44.53	307.01	111.99	111.99
M6	308.09	270.64	233.20	195.75	158.30	120.86	85.41	45.96	8.52	331.07	293.62
N1	164.37	113.86	63.35	12.84	271.82	221.30	170.79	120.28	69.77	19.71	19.71
O	80.96	55.59	30.23	4.86	339.49	314.13	288.76	263.38	238.02	212.66	187.29
O0	300.72	328.06	355.40	22.74	50.08	77.42	104.75	132.09	159.43	186.77	214.11
P1	-10.84	-11.83	-12.81	-13.80	-14.78	-15.77	-16.74	-17.73	-18.71	-19.71	-20.70
Q1	297.25	258.81	220.38	181.95	143.52	105.09	66.65	28.22	346.79	311.36	272.92
Q2	153.53	102.63	50.54	359.04	307.54	266.05	204.55	153.65	101.55	50.06	258.56
R2	10.84	11.83	12.81	13.80	14.78	15.77	16.76	17.74	18.73	19.71	20.70
S1	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
T1	-10.84	-11.83	-12.81	-13.80	-14.78	-15.77	-16.74	-17.73	-18.71	-19.70	-20.70
M2	216.28	203.22	190.16	177.09	164.02	150.96	137.98	124.83	111.76	104.74	85.64
M3	183.61	134.84	86.08	37.32	348.56	299.79	251.03	202.27	153.50	104.74	55.98
M4	327.32	291.64	255.93	220.23	184.53	148.83	113.13	77.44	41.74	6.04	350.34
M5	316.48	279.33	243.11	206.43	169.75	133.06	96.38	59.69	346.33	309.04	288.69
M6	102.66	79.25	55.86	32.46	9.06	345.67	322.27	298.88	275.48	252.08	235.28
2MK	172.76	123.02	73.27	23.52	333.06	284.02	234.27	184.52	134.78	105.03	141.61
MN	39.80	338.06	276.24	214.41	152.58	90.75	28.92	327.10	265.27	203.44	207.99
MS	91.80	67.47	43.04	18.66	354.96	329.90	305.52	281.13	256.75	232.37	207.99
2SM	268.20	292.58	316.96	341.34	5.72	30.10	54.38	78.87	103.25	127.63	152.01
Mf	289.88	316.23	342.59	8.94	35.29	61.64	88.00	114.35	140.70	167.06	193.41
Msf	268.20	292.58	316.96	311.34	5.72	30.10	54.48	78.87	103.25	127.63	152.01
Mm	143.72	156.78	169.84	182.91	195.98	209.04	222.10	248.24	261.30	19.71	20.70
Sa	10.84	11.83	12.81	13.80	14.78	15.77	16.76	17.74	18.73	19.71	20.70
Ssa	21.68	23.66	25.63	27.61	29.57	31.51	33.51	35.48	37.46	39.43	41.40

\*The table is adapted directly for use with common years, but if the required date falls between Mar. 1 and Dec. 31, inclusive, in a leap year the day of month should be increased by one before entering the table.

†The first line for constituent M gives the differences as based upon the formula in table 2; the second line gives the differences as derived from the half speed of constituent M<sub>2</sub>.

Table 17.—*Differences to adapt table 16 to beginning of each day of month*—Continued

Constituent	Day of month*										
	23	24	25	26	27	28	29	30	31	32	
J <sub>1</sub>	○	○	○	○	○	○	○	○	○	○	■
K <sub>1</sub>	309.11	323.16	337.22	351.27	5.32	19.37	33.42	61.52	75.57		
K <sub>2</sub>	21.68	22.67	23.66	24.64	25.63	26.61	27.60	28.68	30.56		
K <sub>3</sub>	43.37	45.34	47.31	49.28	51.26	53.22	55.20	57.17	59.14	61.11	
L <sub>2</sub>	111.04	99.72	88.40	77.00	65.45	43.14	31.32	20.50	9.19		
M <sub>1</sub> †	94.25	82.18	70.10	58.02	45.94	33.86	21.78	9.70	357.62	345.54	
M <sub>1</sub>	91.80	79.61	67.42	55.23	43.04	30.85	18.66	6.47	354.28	342.09	
M <sub>2</sub>	183.61	159.23	134.84	101.46	68.08	61.70	52.32	42.94	348.56	324.17	
M <sub>3</sub>	275.41	238.84	202.27	165.69	129.12	92.55	65.98	19.40	342.88	306.26	
M <sub>4</sub>	7.21	318.45	269.69	220.92	172.16	123.40	74.64	25.87	337.11	288.35	
M <sub>6</sub>	190.82	117.68	44.53	331.39	268.24	185.10	111.95	38.81	325.66	252.52	
M <sub>8</sub>	14.43	276.90	176.38	81.85	314.32	246.80	149.27	61.75	314.22	216.69	
N <sub>2</sub>	256.18	218.73	181.28	133.84	106.39	68.94	31.50	354.05	216.60	279.15	
2N	328.75	278.24	227.72	177.21	126.70	76.19	25.68	335.17	284.66	234.15	
O <sub>1</sub>	161.92	136.56	111.19	85.82	60.45	35.09	9.72	344.35	293.62		
O <sub>0</sub>	241.45	268.78	296.12	323.46	350.80	381.80	18.14	45.48	372.81	100.15	
P <sub>1</sub>	-21.68	-22.67	-23.66	-24.64	-25.63	-26.61	-27.60	-28.58	-29.57	-30.56	
Q <sub>1</sub>	234.49	196.06	171.63	119.20	80.76	42.33	3.47	287.04	225.87	203.59	
2Q	307.06	255.67	204.07	152.57	101.07	49.58	358.05	306.88	255.09	203.59	
R <sub>2</sub>	21.68	22.67	23.66	24.64	25.63	26.61	27.60	28.58	29.57	30.56	
S <sub>1</sub>	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	
T <sub>2</sub>	-21.68	-22.67	-23.66	-24.64	-25.63	-26.61	-27.60	-28.58	-29.57	-30.56	
X <sub>2</sub>	7.21	59.50	46.44	33.38	20.31	7.24	354.18	341.12	328.05	314.98	
A <sub>2</sub>	294.64	258.95	223.25	187.55	151.86	116.15	80.46	44.76	25.87	337.11	
v <sub>2</sub>	272.96	236.28	199.59	162.91	126.22	89.54	52.86	16.17	339.49	302.81	
P <sub>1</sub>	205.29	181.90	158.50	135.10	103.10	88.31	64.92	41.52	18.12	354.73	
M <sub>K</sub>	345.53	295.78	246.03	196.28	146.54	96.79	47.04	357.29	307.54		
2MK	70.78	17.96	316.13	254.30	192.47	130.64	68.82	6.99	305.16	243.33	
M <sub>N</sub>	183.61	159.23	134.84	110.46	86.08	61.70	37.32	12.94	348.66	324.17	
2S <sub>M</sub>	176.39	200.77	225.16	249.64	273.92	298.30	322.68	347.06	11.44	35.83	
M <sub>f</sub>	219.76	246.11	272.47	298.82	325.17	351.52	17.88	44.23	70.58	96.64	
M <sub>st</sub>	176.39	200.77	225.16	249.54	273.92	298.30	322.68	347.06	11.44	35.83	
M <sub>m</sub>	287.43	300.50	313.56	326.62	339.69	362.76	5.82	18.88	45.02	45.02	
S <sub>a</sub>	21.68	22.67	23.66	24.64	25.63	26.61	27.60	28.58	29.57	30.56	
s <sub>sa</sub>	43.37	45.34	47.31	49.28	51.26	53.22	55.20	57.17	59.14	61.11	

\*The table is adapted directly for use with common years but if the required date falls between Mar. 1 and Dec. 31, inclusive, in a leap year the day of month should be increased by one before entering the table.

†The first line for constituent M<sub>1</sub> gives the differences as based upon the formula in Table 2; the second line gives the differences as derived from the half speed of constituent M<sub>1</sub>.

Table 18.—*Differences to adapt table 15 to beginning of each hour of day*

Constituent	Hour of day										
	0	1	2	3	4	5	6	7	8	9	10
J <sub>1</sub>	o	o	o	o	o	o	o	o	o	o	o
K <sub>1</sub>	0.00	15.59	31.17	46.76	62.34	77.93	93.51	109.10	124.68	140.27	155.85
K <sub>2</sub>	0.00	15.04	30.08	45.12	60.16	75.21	90.25	105.29	120.33	135.37	150.41
I <sub>2</sub>	0.00	30.08	60.16	90.25	120.33	150.41	180.49	210.57	246.66	270.74	300.82
I <sub>1</sub>	0.00	25.53	59.06	88.59	118.11	147.64	177.17	206.70	236.23	265.28	290.90
M <sub>1</sub> A <sub>1</sub>	0.00	14.60	28.99	43.49	57.99	72.48	86.98	101.48	115.97	130.47	144.97
M <sub>1</sub>	0.00	14.49	28.98	43.48	57.97	72.46	86.95	101.44	115.94	130.43	144.92
M <sub>2</sub>	0.00	28.98	57.97	86.95	115.94	144.92	173.90	217.38	260.86	298.84	318.83
M <sub>3</sub>	0.00	43.48	86.95	130.43	173.90	217.38	260.86	304.33	347.81	381.29	418.24
M <sub>4</sub>	0.00	57.97	115.94	173.90	231.87	289.84	347.81	45.78	103.75	161.71	218.68
M <sub>5</sub>	0.00	86.95	173.90	260.86	347.81	49.76	161.71	248.67	335.62	419.57	277.65
M <sub>6</sub>	0.00	115.94	173.90	231.87	289.84	347.81	49.76	161.71	248.67	335.62	419.57
N <sub>2</sub>	0.00	28.44	56.88	85.32	113.76	142.20	170.64	199.08	227.52	255.96	284.40
N <sub>2</sub> N	0.00	27.90	55.79	83.69	111.68	139.48	167.37	195.27	223.16	251.06	278.96
O <sub>1</sub>	0.00	13.94	27.89	41.83	55.77	69.52	83.66	97.60	111.54	125.49	138.43
O <sub>2</sub>	0.00	16.14	32.28	48.42	64.56	80.70	96.83	112.97	129.11	145.25	161.39
P <sub>1</sub>	0.00	14.96	29.92	44.88	59.84	74.79	89.75	104.71	119.67	134.63	149.59
Q <sub>1</sub>	0.00	13.40	26.80	40.20	53.69	66.99	80.39	93.79	107.19	120.59	133.99
Q <sub>2</sub>	0.00	12.85	24.71	38.56	51.42	64.27	77.13	89.98	102.83	115.69	128.54
R <sub>2</sub>	0.00	30.04	60.08	90.12	120.16	150.21	180.25	210.29	240.33	270.37	300.41
S <sub>1</sub>	0.00	15.00	30.00	45.00	60.00	75.00	90.00	105.00	120.00	135.00	150.00
S <sub>2</sub>	0.00	30.00	60.00	90.00	120.00	150.00	180.00	210.00	240.00	270.00	300.00
S <sub>4</sub>	0.00	60.00	120.00	180.00	240.00	300.00	0.00	60.00	120.00	180.00	240.00
S <sub>6</sub>	0.00	90.00	180.00	270.00	360.00	450.00	90.00	180.00	270.00	360.00	450.00
T <sub>2</sub>	0.00	29.96	59.92	89.88	119.84	149.79	179.76	209.71	239.67	269.63	299.59
$\lambda_2$	0.00	29.46	58.91	88.37	117.82	147.28	176.73	206.19	235.65	265.10	294.01
$\mu_2$	0.00	27.97	55.94	83.90	111.87	139.84	167.81	195.78	223.75	251.71	278.68
$p_2$	0.00	28.51	57.63	85.54	114.05	142.56	171.08	199.59	228.10	256.61	285.13
$\rho_1$	0.00	13.47	26.94	40.41	53.89	67.36	80.83	94.30	107.77	121.24	134.72
M <sub>K</sub>	0.00	44.03	88.05	132.08	176.10	220.13	264.15	308.18	352.20	386.23	424.28
2M <sub>K</sub>	0.00	42.93	85.85	128.78	171.71	214.64	257.56	300.49	343.42	386.34	428.27
M <sub>N</sub>	0.00	57.42	114.85	172.27	229.70	287.12	344.54	41.97	99.39	156.81	214.24
M <sub>S</sub>	0.00	58.98	117.97	176.95	235.94	293.90	52.89	111.87	170.96	229.84	271.66
2S <sub>M</sub>	0.00	31.92	62.63	93.06	124.06	155.08	186.10	217.11	248.13	278.14	311.17
M <sub>T</sub>	0.00	1.10	2.20	3.29	4.39	5.49	6.59	7.69	8.78	10.98	12.08
MSL	0.00	0.02	2.03	3.05	4.06	5.08	6.10	7.11	8.13	9.14	10.16
M <sub>M</sub>	0.00	0.54	1.09	1.63	2.18	2.72	3.27	3.81	4.35	5.44	5.99
Sa	0.00	0.04	0.08	0.12	0.16	0.25	0.33	0.41	0.37	0.41	0.45
Ss <sub>a</sub>	0.00	0.08	0.16	0.25	0.33	0.41	0.49	0.57	0.66	0.74	0.90

<sup>†</sup>The first line for constituent M<sub>1</sub> gives the differences as based upon the formula in table 2; the second line gives the differences as derived from the half speed of constituent M<sub>3</sub>.

Table 18.—*Differences to adapt table 15 to beginning of each hour of day—Continued*

Constituent	Hour of day											
	12	13	14	15	16	17	18	19	20	21	22	23
$J_1$	○	□	○	○	○	○	○	○	○	○	○	○
$K_1$	187.03	202.61	218.30	233.78	249.37	264.95	280.54	296.12	311.71	327.29	342.88	358.47
$K_2$	180.49	195.53	210.57	225.62	240.66	255.74	270.74	285.78	300.82	315.86	330.90	345.94
$L_1$	0.99	31.07	61.15	91.23	121.31	151.40	181.48	211.56	241.64	271.72	301.81	331.89
$M_{1f}$	354.34	323.57	33.40	82.93	112.46	141.98	171.04	201.04	230.57	260.10	286.63	319.16
$M_1$	173.96	188.46	202.95	217.45	231.95	246.94	260.94	275.44	289.95	304.55	318.95	333.42
$M_2$	173.90	188.40	202.89	217.38	231.87	246.36	260.86	275.35	289.84	304.33	318.83	333.32
$M_3$	347.81	16.79	45.78	74.76	103.75	132.75	161.79	190.74	219.68	248.67	277.65	306.63
$M_4$	161.71	205.19	248.07	292.14	335.62	19.09	62.57	106.05	149.62	183.00	226.48	279.95
$M_5$	385.62	33.69	91.55	149.52	265.46	201.49	232.43	381.40	79.36	137.33	195.30	253.27
$M_6$	323.43	50.38	137.33	224.38	311.24	38.19	125.14	212.08	299.05	25.00	112.95	199.90
$M_8$	311.24	67.17	183.11	299.05	54.98	170.92	286.96	42.79	188.73	274.66	30.60	146.54
$N_1$	341.28	9.72	88.76	66.60	98.04	163.48	161.92	180.35	208.79	237.23	265.67	294.11
$N_2$	334.74	2.64	30.55	88.43	114.22	142.12	170.11	197.91	225.80	253.70	281.70	310.63
$O_1$	167.32	181.26	206.15	228.09	242.09	258.23	274.36	290.50	328.80	320.75	320.69	320.69
$O_2$	193.67	209.81	225.85	242.09	258.23	274.36	290.50	306.64	322.78	338.92	355.66	371.20
$P_1$	179.51	194.47	209.43	224.38	239.34	254.30	269.26	284.22	299.18	314.14	329.10	344.06
$Q_1$	160.78	174.18	187.68	200.98	214.38	227.78	241.18	254.57	267.97	284.10	304.17	320.17
$Q_2$	154.25	167.11	179.96	192.81	205.67	218.52	231.38	244.23	257.09	269.94	282.79	295.65
$R_3$	270.49	30.53	60.57	90.62	120.66	180.74	210.78	240.82	270.86	300.90	330.94	345.00
$S_1$	180.00	195.00	210.00	225.00	240.00	255.00	270.00	285.00	300.00	315.00	330.00	345.00
$S_2$	0.00	30.00	60.00	90.00	120.00	150.00	180.00	210.00	240.00	270.00	300.00	330.00
$S_4$	0.00	60.00	120.00	180.00	240.00	300.00	0.00	60.00	120.00	180.00	240.00	300.00
$T_3$	369.51	29.47	69.43	89.38	119.34	149.20	179.26	200.00	270.00	0.00	180.00	270.00
$\lambda_2$	333.47	22.92	52.38	81.88	111.29	140.75	170.20	199.66	239.22	269.14	299.22	320.77
$p_2$	335.62	3.59	31.56	59.52	87.49	141.46	171.43	171.40	199.36	227.33	255.30	283.27
$p_2$	342.16	10.66	39.18	67.69	96.20	124.71	153.23	181.74	210.25	238.76	267.28	295.96
$\rho$	161.66	175.13	188.60	202.07	215.54	229.02	242.49	255.96	269.43	282.90	286.37	309.84
$MK$	168.30	212.33	265.35	300.38	344.40	28.43	72.45	116.48	160.50	204.53	248.45	292.58
$2MK$	155.13	198.05	240.98	283.91	326.88	9.76	52.69	138.54	181.47	224.40	267.32	300.00
$MN$	83.93	26.51	105.78	164.76	223.75	256.21	313.63	11.05	68.48	125.90	183.32	240.75
$MS$	347.81	46.79	74.22	105.24	136.25	167.27	198.29	229.30	260.32	291.33	322.35	353.37
$2SM$	12.19	13.18	14.27	15.37	16.47	17.57	18.67	20.86	21.96	24.16	25.25	27.37
$MT$	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....
$MSL$	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....
$Mm$	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....
$Sa$	0.49	0.53	7.08	8.17	9.25	9.80	10.34	10.89	11.43	11.98	12.52	13.94
$Ssa$	0.99	1.07	0.57	0.62	0.66	0.70	0.74	0.82	0.86	0.90	1.04	1.18

The first line for constituent  $M_1$  gives the differences as based upon the formula in table 2; the second line gives the differences as derived from the half speed of constituent  $M_1$ .

Table 19.—*Products for Form 194*

[Multiplier=sin 15°=0.259]

	0	1	2	3	4	5	6	7	8	9
0.00.....	0.000	0.259	0.518	0.777	1.036	1.295	1.554	1.813	2.072	2.331
.01.....	.003	.262	.521	.780	1.039	1.298	1.557	1.816	2.075	2.334
.02.....	.005	.264	.523	.782	1.041	1.300	1.559	1.818	2.077	2.336
.03.....	.008	.267	.526	.785	1.044	1.303	1.562	1.821	2.080	2.339
.04.....	.010	.269	.528	.787	1.046	1.305	1.564	1.823	2.082	2.341
.05.....	.013	.272	.531	.790	1.049	1.308	1.567	1.826	2.085	2.344
.06.....	.016	.275	.534	.793	1.052	1.311	1.570	1.829	2.088	2.347
.07.....	.018	.277	.536	.795	1.054	1.313	1.572	1.831	2.090	2.349
.08.....	.021	.280	.539	.798	1.057	1.316	1.575	1.834	2.093	2.352
.09.....	.023	.282	.541	.800	1.059	1.318	1.577	1.836	2.095	2.354
.10.....	.026	.285	.544	.803	1.062	1.321	1.580	1.839	2.098	2.357
.11.....	.028	.287	.546	.805	1.064	1.323	1.582	1.841	2.100	2.359
.12.....	.031	.290	.549	.808	1.067	1.326	1.585	1.844	2.103	2.362
.13.....	.034	.293	.552	.811	1.070	1.329	1.588	1.847	2.106	2.365
.14.....	.036	.295	.554	.813	1.072	1.331	1.590	1.849	2.108	2.367
.15.....	.039	.298	.557	.816	1.075	1.334	1.593	1.852	2.111	2.370
.16.....	.041	.300	.559	.818	1.077	1.336	1.595	1.854	2.113	2.372
.17.....	.044	.303	.562	.821	1.080	1.339	1.598	1.857	2.116	2.375
.18.....	.047	.306	.565	.824	1.083	1.342	1.601	1.860	2.119	2.378
.19.....	.049	.308	.567	.826	1.085	1.344	1.603	1.862	2.121	2.380
.20.....	.052	.311	.570	.829	1.088	1.347	1.606	1.865	2.124	2.383
.21.....	.054	.313	.572	.831	1.090	1.349	1.608	1.867	2.126	2.385
.22.....	.057	.316	.575	.834	1.093	1.352	1.611	1.870	2.129	2.388
.23.....	.060	.319	.578	.837	1.096	1.355	1.614	1.873	2.132	2.391
.24.....	.062	.321	.580	.839	1.098	1.357	1.616	1.875	2.134	2.393
.25.....	.065	.324	.583	.842	1.101	1.360	1.619	1.878	2.137	2.396
.26.....	.067	.326	.585	.844	1.103	1.362	1.621	1.880	2.139	2.398
.27.....	.070	.329	.588	.847	1.106	1.365	1.624	1.883	2.142	2.401
.28.....	.073	.332	.591	.850	1.109	1.368	1.627	1.886	2.145	2.404
.29.....	.075	.334	.593	.852	1.111	1.370	1.629	1.888	2.147	2.406
.30.....	.078	.337	.596	.855	1.114	1.373	1.632	1.891	2.150	2.409
.31.....	.080	.339	.598	.857	1.116	1.375	1.634	1.893	2.152	2.411
.32.....	.083	.342	.601	.860	1.119	1.378	1.637	1.896	2.155	2.414
.33.....	.085	.344	.603	.862	1.121	1.380	1.639	1.898	2.157	2.416
.34.....	.088	.347	.606	.865	1.124	1.383	1.642	1.901	2.160	2.419
.35.....	.091	.350	.609	.868	1.127	1.386	1.645	1.904	2.163	2.422
.36.....	.093	.352	.611	.870	1.129	1.388	1.647	1.906	2.165	2.424
.37.....	.096	.355	.614	.873	1.132	1.391	1.650	1.909	2.168	2.427
.38.....	.098	.357	.616	.875	1.134	1.393	1.652	1.911	2.170	2.429
.39.....	.101	.360	.619	.878	1.137	1.396	1.655	1.914	2.173	2.432
.40.....	.104	.363	.622	.881	1.140	1.399	1.658	1.917	2.176	2.435
.41.....	.106	.365	.624	.883	1.142	1.401	1.660	1.919	2.178	2.437
.42.....	.109	.368	.627	.886	1.145	1.404	1.663	1.922	2.181	2.440
.43.....	.111	.370	.629	.888	1.147	1.406	1.665	1.924	2.183	2.442
.44.....	.114	.373	.632	.891	1.150	1.409	1.668	1.927	2.186	2.445
.45.....	.117	.376	.635	.894	1.153	1.412	1.671	1.930	2.189	2.448
.46.....	.119	.378	.637	.896	1.155	1.414	1.673	1.932	2.191	2.450
.47.....	.122	.381	.640	.899	1.158	1.417	1.676	1.935	2.194	2.453
.48.....	.124	.383	.642	.901	1.160	1.419	1.678	1.937	2.196	2.455
.49.....	.127	.386	.645	.904	1.163	1.422	1.681	1.940	2.199	2.458
.50.....	.130	.388	.648	.906	1.166	1.424	1.684	1.942	2.202	2.460
	0	1	2	3	4	5	6	7	8	9

Table 19.—*Products for Form 194—Continued*[Multiplier= $\sin 15^\circ = 0.259$ ]

	0	1	2	3	4	5	6	7	8	9
.50	0.130	0.388	0.648	0.906	1.166	1.424	1.684	1.942	2.202	2.460
.51	.132	.391	.650	.909	1.168	1.427	1.686	1.945	2.204	2.463
.52	.135	.394	.653	.912	1.171	1.430	1.689	1.948	2.207	2.466
.53	.137	.396	.655	.914	1.173	1.432	1.691	1.950	2.209	2.468
.54	.140	.399	.658	.917	1.176	1.435	1.694	1.953	2.212	2.471
.55	.142	.401	.660	.919	1.178	1.437	1.696	1.955	2.214	2.473
.56	.145	.404	.663	.922	1.181	1.440	1.699	1.958	2.217	2.476
.57	.148	.407	.666	.925	1.184	1.443	1.702	1.961	2.220	2.479
.58	.150	.409	.668	.927	1.186	1.445	1.704	1.963	2.222	2.481
.59	.153	.412	.671	.930	1.189	1.448	1.707	1.966	2.225	2.484
.60	.155	.414	.673	.932	1.191	1.450	1.709	1.968	2.227	2.486
.61	.158	.417	.676	.935	1.194	1.453	1.712	1.971	2.230	2.489
.62	.161	.420	.679	.938	1.197	1.456	1.715	1.974	2.233	2.492
.63	.163	.422	.681	.940	1.199	1.458	1.717	1.976	2.235	2.494
.64	.166	.425	.684	.943	1.202	1.461	1.720	1.979	2.238	2.497
.65	.168	.427	.686	.945	1.204	1.463	1.722	1.981	2.240	2.499
.66	.171	.430	.689	.948	1.207	1.466	1.725	1.984	2.243	2.502
.67	.174	.433	.692	.951	1.210	1.469	1.728	1.987	2.246	2.505
.68	.176	.435	.694	.953	1.212	1.471	1.730	1.989	2.248	2.507
.69	.179	.438	.697	.956	1.215	1.474	1.733	1.992	2.251	2.510
.70	.181	.440	.699	.958	1.217	1.476	1.735	1.994	2.253	2.512
.71	.184	.443	.702	.961	1.220	1.479	1.738	1.997	2.256	2.515
.72	.186	.445	.704	.963	1.222	1.481	1.740	1.999	2.258	2.517
.73	.189	.448	.707	.966	1.225	1.484	1.743	2.002	2.261	2.520
.74	.192	.451	.710	.969	1.228	1.487	1.746	2.005	2.264	2.523
.75	.194	.453	.712	.971	1.230	1.489	1.748	2.007	2.266	2.525
.76	.197	.456	.715	.974	1.233	1.492	1.751	2.010	2.269	2.528
.77	.199	.458	.717	.976	1.235	1.494	1.753	2.012	2.271	2.530
.78	.202	.461	.720	.979	1.238	1.497	1.756	2.015	2.274	2.533
.79	.205	.464	.723	.982	1.241	1.500	1.759	2.018	2.277	2.536
.80	.207	.466	.725	.984	1.243	1.502	1.761	2.020	2.279	2.538
.81	.210	.469	.728	.987	1.246	1.505	1.764	2.023	2.282	2.541
.82	.212	.471	.730	.989	1.248	1.507	1.766	2.025	2.284	2.543
.83	.215	.474	.733	.992	1.251	1.510	1.769	2.028	2.287	2.546
.84	.218	.477	.736	.995	1.254	1.513	1.772	2.031	2.290	2.549
.85	.220	.479	.738	.997	1.256	1.515	1.774	2.033	2.292	2.551
.86	.223	.482	.741	1.000	1.259	1.518	1.777	2.036	2.295	2.554
.87	.225	.484	.743	1.002	1.261	1.520	1.779	2.038	2.297	2.556
.88	.228	.487	.746	1.005	1.264	1.523	1.782	2.041	2.300	2.559
.89	.231	.490	.749	1.008	1.267	1.526	1.785	2.044	2.303	2.562
.90	.233	.492	.751	1.010	1.269	1.528	1.787	2.046	2.305	2.564
.91	.236	.495	.754	1.013	1.272	1.531	1.790	2.049	2.308	2.567
.92	.238	.497	.756	1.015	1.274	1.533	1.792	2.051	2.310	2.569
.93	.241	.500	.759	1.018	1.277	1.536	1.795	2.054	2.313	2.572
.94	.243	.502	.761	1.020	1.279	1.538	1.797	2.056	2.315	2.574
.95	.246	.505	.764	1.023	1.282	1.541	1.800	2.059	2.318	2.577
.96	.249	.508	.767	1.026	1.285	1.544	1.803	2.062	2.321	2.580
.97	.251	.510	.769	1.028	1.287	1.546	1.805	2.064	2.323	2.582
.98	.254	.513	.772	1.031	1.290	1.549	1.808	2.067	2.326	2.585
.99	.256	.515	.774	1.033	1.292	1.551	1.810	2.069	2.328	2.587
1.00	.259	.518	.777	1.036	1.295	1.554	1.813	2.072	2.331	2.590
	0	1	2	3	4	5	6	7	8	9

Table 19.—*Products for Form 194—Continued*[Multiplier= $\sin 30^\circ = 0.500$ ]

	0	1	2	3	4	5	6	7	8	9
0.00.....	0.000	0.500	1.000	1.500	2.000	2.500	3.000	3.500	4.000	4.500
.01.....	.005	.505	1.005	1.505	2.005	2.505	3.005	3.505	4.005	4.505
.02.....	.010	.510	1.010	1.510	2.010	2.510	3.010	3.510	4.010	4.510
.03.....	.015	.515	1.015	1.515	2.015	2.515	3.015	3.515	4.015	4.515
.04.....	.020	.520	1.020	1.520	2.020	2.520	3.020	3.520	4.020	4.520
.05.....	.025	.525	1.025	1.525	2.025	2.525	3.025	3.525	4.025	4.525
.06.....	.030	.530	1.030	1.530	2.030	2.530	3.030	3.530	4.030	4.530
.07.....	.035	.535	1.035	1.535	2.035	2.535	3.035	3.535	4.035	4.535
.08.....	.040	.540	1.040	1.540	2.040	2.540	3.040	3.540	4.040	4.540
.09.....	.045	.545	1.045	1.545	2.045	2.545	3.045	3.545	4.045	4.545
.10.....	.050	.550	1.050	1.550	2.050	2.550	3.050	3.550	4.050	4.550
.11.....	.055	.555	1.055	1.555	2.055	2.555	3.055	3.555	4.055	4.555
.12.....	.060	.560	1.060	1.560	2.060	2.560	3.060	3.560	4.060	4.560
.13.....	.065	.565	1.065	1.565	2.065	2.565	3.065	3.565	4.065	4.565
.14.....	.070	.570	1.070	1.570	2.070	2.570	3.070	3.570	4.070	4.570
.15.....	.075	.575	1.075	1.575	2.075	2.575	3.075	3.575	4.075	4.575
.16.....	.080	.580	1.080	1.580	2.080	2.580	3.080	3.580	4.080	4.580
.17.....	.085	.585	1.085	1.585	2.085	2.585	3.085	3.585	4.085	4.585
.18.....	.090	.590	1.090	1.590	2.090	2.590	3.090	3.590	4.090	4.590
.19.....	.095	.595	1.095	1.595	2.095	2.595	3.095	3.595	4.095	4.595
.20.....	.100	.600	1.100	1.600	2.100	2.600	3.100	3.600	4.100	4.600
.21.....	.105	.605	1.105	1.605	2.105	2.605	3.105	3.605	4.105	4.605
.22.....	.110	.610	1.110	1.610	2.110	2.610	3.110	3.610	4.110	4.610
.23.....	.115	.615	1.115	1.615	2.115	2.615	3.115	3.615	4.115	4.615
.24.....	.120	.620	1.120	1.620	2.120	2.620	3.120	3.620	4.120	4.620
.25.....	.125	.625	1.125	1.625	2.125	2.625	3.125	3.625	4.125	4.625
.26.....	.130	.630	1.130	1.630	2.130	2.630	3.130	3.630	4.130	4.630
.27.....	.135	.635	1.135	1.635	2.135	2.635	3.135	3.635	4.135	4.635
.28.....	.140	.640	1.140	1.640	2.140	2.640	3.140	3.640	4.140	4.640
.29.....	.145	.645	1.145	1.645	2.145	2.645	3.145	3.645	4.145	4.645
.30.....	.150	.650	1.150	1.650	2.150	2.650	3.150	3.650	4.150	4.650
.31.....	.155	.655	1.155	1.655	2.155	2.655	3.155	3.655	4.155	4.655
.32.....	.160	.660	1.160	1.660	2.160	2.660	3.160	3.660	4.160	4.660
.33.....	.165	.665	1.165	1.665	2.165	2.665	3.165	3.665	4.165	4.665
.34.....	.170	.670	1.170	1.670	2.170	2.670	3.170	3.670	4.170	4.670
.35.....	.175	.675	1.175	1.675	2.175	2.675	3.175	3.675	4.175	4.675
.36.....	.180	.680	1.180	1.680	2.180	2.680	3.180	3.680	4.180	4.680
.37.....	.185	.685	1.185	1.685	2.185	2.685	3.185	3.685	4.185	4.685
.38.....	.190	.690	1.190	1.690	2.190	2.690	3.190	3.690	4.190	4.690
.39.....	.195	.695	1.195	1.695	2.195	2.695	3.195	3.695	4.195	4.695
.40.....	.200	.700	1.200	1.700	2.200	2.700	3.200	3.700	4.200	4.700
.41.....	.205	.705	1.205	1.705	2.205	2.705	3.205	3.705	4.205	4.705
.42.....	.210	.710	1.210	1.710	2.210	2.710	3.210	3.710	4.210	4.710
.43.....	.215	.715	1.215	1.715	2.215	2.715	3.215	3.715	4.215	4.715
.44.....	.220	.720	1.220	1.720	2.220	2.720	3.220	3.720	4.220	4.720
.45.....	.225	.725	1.225	1.725	2.225	2.725	3.225	3.725	4.225	4.725
.46.....	.230	.730	1.230	1.730	2.230	2.730	3.230	3.730	4.230	4.730
.47.....	.235	.735	1.235	1.735	2.235	2.735	3.235	3.735	4.235	4.735
.48.....	.240	.740	1.240	1.740	2.240	2.740	3.240	3.740	4.240	4.740
.49.....	.245	.745	1.245	1.745	2.245	2.745	3.245	3.745	4.245	4.745
.50.....	.250	.750	1.250	1.750	2.250	2.750	3.250	3.750	4.250	4.750
	0	1	2	3	4	5	6	7	8	9

Table 19.—*Products for Form 194—Continued*[Multiplier =  $\sin 30^\circ = 0.500$ ]

	0	1	2	3	4	5	6	7	8	9
0.50-----	0.250	0.750	1.250	1.750	2.250	2.750	3.250	3.750	4.250	4.750
.51-----	.255	.755	1.255	1.755	2.255	2.755	3.255	3.755	4.255	4.755
.52-----	.260	.760	1.260	1.760	2.260	2.760	3.260	3.760	4.260	4.760
.53-----	.265	.765	1.265	1.765	2.265	2.765	3.265	3.765	4.265	4.765
.54-----	.270	.770	1.270	1.770	2.270	2.770	3.270	3.770	4.270	4.770
.55-----	.275	.775	1.275	1.775	2.275	2.775	3.275	3.775	4.275	4.775
.56-----	.280	.780	1.280	1.780	2.280	2.780	3.280	3.780	4.280	4.780
.57-----	.285	.785	1.285	1.785	2.285	2.785	3.285	3.785	4.285	4.785
.58-----	.290	.790	1.290	1.790	2.290	2.790	3.290	3.790	4.290	4.790
.59-----	.295	.795	1.295	1.795	2.295	2.795	3.295	3.795	4.295	4.795
.60-----	.300	.800	1.300	1.800	2.300	2.800	3.300	3.800	4.300	4.800
.61-----	.305	.805	1.305	1.805	2.305	2.805	3.305	3.805	4.305	4.805
.62-----	.310	.810	1.310	1.810	2.310	2.810	3.310	3.810	4.310	4.810
.63-----	.315	.815	1.315	1.815	2.315	2.815	3.315	3.815	4.315	4.815
.64-----	.320	.820	1.320	1.820	2.320	2.820	3.320	3.820	4.320	4.820
.65-----	.325	.825	1.325	1.825	2.325	2.825	3.325	3.825	4.325	4.825
.66-----	.330	.830	1.330	1.830	2.330	2.830	3.330	3.830	4.330	4.830
.67-----	.335	.835	1.335	1.835	2.335	2.835	3.335	3.835	4.335	4.835
.68-----	.340	.840	1.340	1.840	2.340	2.840	3.340	3.840	4.340	4.840
.69-----	.345	.845	1.345	1.845	2.345	2.845	3.345	3.845	4.345	4.845
.70-----	.350	.850	1.350	1.850	2.350	2.850	3.350	3.850	4.350	4.850
.71-----	.355	.855	1.355	1.855	2.355	2.855	3.355	3.855	4.355	4.855
.72-----	.360	.860	1.360	1.860	2.360	2.860	3.360	3.860	4.360	4.860
.73-----	.365	.865	1.365	1.865	2.365	2.865	3.365	3.865	4.365	4.865
.74-----	.370	.870	1.370	1.870	2.370	2.870	3.370	3.870	4.370	4.870
.75-----	.375	.875	1.375	1.875	2.375	2.875	3.375	3.875	4.375	4.875
.76-----	.380	.880	1.380	1.880	2.380	2.880	3.380	3.880	4.380	4.880
.77-----	.385	.885	1.385	1.885	2.385	2.885	3.385	3.885	4.385	4.885
.78-----	.390	.890	1.390	1.890	2.390	2.890	3.390	3.890	4.390	4.890
.79-----	.395	.895	1.395	1.895	2.395	2.895	3.395	3.895	4.395	4.895
.80-----	.400	.900	1.400	1.900	2.400	2.900	3.400	3.900	4.400	4.900
.81-----	.405	.905	1.405	1.905	2.405	2.905	3.405	3.905	4.405	4.905
.82-----	.410	.910	1.410	1.910	2.410	2.910	3.410	3.910	4.410	4.910
.83-----	.415	.915	1.415	1.915	2.415	2.915	3.415	3.915	4.415	4.915
.84-----	.420	.920	1.420	1.920	2.420	2.920	3.420	3.920	4.420	4.920
.85-----	.425	.925	1.425	1.925	2.425	2.925	3.425	3.925	4.425	4.925
.86-----	.430	.930	1.430	1.930	2.430	2.930	3.430	3.930	4.430	4.930
.87-----	.435	.935	1.435	1.935	2.435	2.935	3.435	3.935	4.435	4.935
.88-----	.440	.940	1.440	1.940	2.440	2.940	3.440	3.940	4.440	4.940
.89-----	.445	.945	1.445	1.945	2.445	2.945	3.445	3.945	4.445	4.945
.90-----	.450	.950	1.450	1.950	2.450	2.950	3.450	3.950	4.450	4.950
.91-----	.455	.955	1.455	1.955	2.455	2.955	3.455	3.955	4.455	4.955
.92-----	.460	.960	1.460	1.960	2.460	2.960	3.460	3.960	4.460	4.960
.93-----	.465	.965	1.465	1.965	2.465	2.965	3.465	3.965	4.465	4.965
.94-----	.470	.970	1.470	1.970	2.470	2.970	3.470	3.970	4.470	4.970
.95-----	.475	.975	1.475	1.975	2.475	2.975	3.475	3.975	4.475	4.975
.96-----	.480	.980	1.480	1.980	2.480	2.980	3.480	3.980	4.480	4.980
.97-----	.485	.985	1.485	1.985	2.485	2.985	3.485	3.985	4.485	4.985
.98-----	.490	.990	1.490	1.990	2.490	2.990	3.490	3.990	4.490	4.990
.99-----	.495	.995	1.495	1.995	2.495	2.995	3.495	3.995	4.495	4.995
1.00-----	.500	1.000	1.500	2.000	2.500	3.000	3.500	4.000	4.500	5.000
	0	1	2	3	4	5	6	7	8	9

Table 19.—*Products for Form 194—Continued*[Multiplier= $\sin 45^\circ = 0.707$ ]

	0	1	2	3	4	5	6	7	8	9
0.00.....	0.000	0.707	1.414	2.121	2.828	3.535	4.242	4.949	5.656	6.363
.01.....	.007	.714	1.421	2.128	2.835	3.542	4.249	4.956	5.663	6.370
.02.....	.014	.721	1.428	2.135	2.842	3.549	4.256	4.963	5.670	6.377
.03.....	.021	.728	1.435	2.142	2.849	3.556	4.263	4.970	5.677	6.384
.04.....	.028	.735	1.442	2.149	2.856	3.563	4.270	4.977	5.684	6.391
.05.....	.035	.742	1.449	2.156	2.863	3.570	4.277	4.984	5.691	6.398
.06.....	.042	.749	1.456	2.163	2.870	3.577	4.284	4.991	5.698	6.405
.07.....	.049	.756	1.463	2.170	2.877	3.584	4.291	4.998	5.705	6.412
.08.....	.057	.764	1.471	2.178	2.885	3.592	4.299	5.006	5.713	6.420
.09.....	.064	.771	1.478	2.185	2.892	3.599	4.306	5.013	5.720	6.427
.10.....	.071	.778	1.485	2.192	2.899	3.606	4.313	5.020	5.727	6.434
.11.....	.078	.785	1.492	2.199	2.906	3.613	4.320	5.027	5.734	6.441
.12.....	.085	.792	1.499	2.206	2.913	3.620	4.327	5.034	5.741	6.448
.13.....	.092	.799	1.506	2.213	2.920	3.627	4.334	5.041	5.748	6.455
.14.....	.099	.806	1.513	2.220	2.927	3.634	4.341	5.048	5.755	6.462
.15.....	.106	.813	1.520	2.227	2.934	3.641	4.348	5.055	5.762	6.469
.16.....	.113	.820	1.527	2.234	2.941	3.648	4.355	5.062	5.769	6.476
.17.....	.120	.827	1.534	2.241	2.948	3.655	4.362	5.069	5.776	6.483
.18.....	.127	.834	1.541	2.248	2.955	3.662	4.369	5.076	5.783	6.490
.19.....	.134	.841	1.548	2.255	2.962	3.669	4.376	5.083	5.790	6.497
.20.....	.141	.848	1.555	2.262	2.969	3.676	4.383	5.090	5.797	6.504
.21.....	.148	.855	1.562	2.269	2.976	3.683	4.390	5.097	5.804	6.511
.22.....	.156	.863	1.570	2.277	2.984	3.691	4.398	5.105	5.812	6.519
.23.....	.163	.870	1.577	2.284	2.991	3.698	4.405	5.112	5.819	6.526
.24.....	.170	.877	1.584	2.291	2.998	3.705	4.412	5.119	5.826	6.533
.25.....	.177	.884	1.591	2.298	3.005	3.712	4.419	5.126	5.833	6.540
.26.....	.184	.891	1.598	2.305	3.012	3.719	4.426	5.133	5.840	6.547
.27.....	.191	.898	1.605	2.312	3.019	3.726	4.433	5.140	5.847	6.554
.28.....	.198	.905	1.612	2.319	3.026	3.733	4.440	5.147	5.854	6.561
.29.....	.205	.912	1.619	2.326	3.033	3.740	4.447	5.154	5.861	6.568
.30.....	.212	.919	1.626	2.333	3.040	3.747	4.454	5.161	5.868	6.575
.31.....	.219	.926	1.633	2.340	3.047	3.754	4.461	5.168	5.875	6.582
.32.....	.226	.933	1.640	2.347	3.054	3.761	4.468	5.175	5.882	6.589
.33.....	.233	.940	1.647	2.354	3.061	3.768	4.475	5.182	5.889	6.596
.34.....	.240	.947	1.654	2.361	3.068	3.775	4.482	5.189	5.896	6.603
.35.....	.247	.954	1.661	2.368	3.075	3.782	4.489	5.196	5.903	6.610
.36.....	.255	.962	1.669	2.376	3.083	3.790	4.497	5.204	5.911	6.618
.37.....	.262	.969	1.676	2.383	3.090	3.797	4.504	5.211	5.918	6.625
.38.....	.269	.976	1.683	2.390	3.097	3.804	4.511	5.218	5.925	6.632
.39.....	.276	.983	1.690	2.397	3.104	3.811	4.518	5.225	5.932	6.639
.40.....	.283	.990	1.697	2.404	3.111	3.818	4.525	5.232	5.939	6.646
.41.....	.290	.997	1.704	2.411	3.118	3.825	4.532	5.239	5.946	6.653
.42.....	.297	1.004	1.711	2.418	3.125	3.832	4.539	5.246	5.953	6.660
.43.....	.304	1.011	1.718	2.425	3.132	3.839	4.546	5.253	5.960	6.667
.44.....	.311	1.018	1.725	2.432	3.139	3.846	4.553	5.260	5.967	6.674
.45.....	.318	1.025	1.732	2.439	3.146	3.853	4.560	5.267	5.974	6.681
.46.....	.325	1.032	1.739	2.446	3.153	3.860	4.567	5.274	5.981	6.688
.47.....	.332	1.039	1.746	2.453	3.160	3.867	4.574	5.281	5.988	6.695
.48.....	.339	1.046	1.753	2.460	3.167	3.874	4.581	5.288	5.995	6.702
.49.....	.346	1.053	1.760	2.467	3.174	3.881	4.588	5.295	6.002	6.709
.50.....	.354	1.060	1.768	2.474	3.182	3.888	4.596	5.302	6.010	6.716

0	1	2	3	4	5	6	7	8	9
---	---	---	---	---	---	---	---	---	---

Table 19.—*Products for Form 194—Continued*[Multiplier =  $\sin 45^\circ = 0.707$ ]

	0	1	2	3	4	5	6	7	8	9
0.50	0.354	1.060	1.768	2.474	3.182	3.888	4.596	5.302	6.010	6.716
.51	.361	1.068	1.775	2.482	3.189	3.896	4.603	5.310	6.017	6.724
.52	.368	1.075	1.782	2.489	3.196	3.903	4.610	5.317	6.024	6.731
.53	.375	1.082	1.789	2.496	3.203	3.910	4.617	5.324	6.031	6.738
.54	.382	1.089	1.796	2.503	3.210	3.917	4.624	5.331	6.038	6.745
.55	.389	1.096	1.803	2.510	3.217	3.924	4.631	5.338	6.045	6.752
.56	.396	1.103	1.810	2.517	3.224	3.931	4.638	5.345	6.052	6.759
.57	.403	1.110	1.817	2.524	3.231	3.938	4.645	5.352	6.059	6.766
.58	.410	1.117	1.824	2.531	3.238	3.945	4.652	5.359	6.066	6.773
.59	.417	1.124	1.831	2.538	3.245	3.952	4.659	5.366	6.073	6.780
.60	.424	1.131	1.838	2.545	3.252	3.959	4.666	5.373	6.080	6.787
.61	.431	1.138	1.845	2.552	3.259	3.966	4.673	5.380	6.087	6.794
.62	.438	1.145	1.852	2.559	3.266	3.973	4.680	5.387	6.094	6.801
.63	.445	1.152	1.859	2.566	3.273	3.980	4.687	5.394	6.101	6.808
.64	.452	1.159	1.866	2.573	3.280	3.987	4.694	5.401	6.108	6.815
.65	.460	1.167	1.874	2.581	3.288	3.995	4.702	5.409	6.116	6.823
.66	.467	1.174	1.881	2.588	3.295	4.002	4.709	5.416	6.123	6.830
.67	.474	1.181	1.888	2.595	3.302	4.009	4.716	5.423	6.130	6.837
.68	.481	1.188	1.895	2.602	3.309	4.016	4.723	5.430	6.137	6.844
.69	.488	1.195	1.902	2.609	3.316	4.023	4.730	5.437	6.144	6.851
.70	.495	1.202	1.909	2.616	3.323	4.030	4.737	5.444	6.151	6.858
.71	.502	1.209	1.916	2.623	3.330	4.037	4.744	5.451	6.158	6.865
.72	.509	1.216	1.923	2.630	3.337	4.044	4.751	5.458	6.165	6.872
.73	.516	1.223	1.930	2.637	3.344	4.051	4.758	5.465	6.172	6.879
.74	.523	1.230	1.937	2.644	3.351	4.058	4.765	5.472	6.179	6.886
.75	.530	1.237	1.944	2.651	3.358	4.065	4.772	5.479	6.186	6.893
.76	.537	1.244	1.951	2.658	3.365	4.072	4.779	5.486	6.193	6.900
.77	.544	1.251	1.958	2.665	3.372	4.079	4.786	5.493	6.200	6.907
.78	.551	1.258	1.965	2.672	3.379	4.086	4.793	5.500	6.207	6.914
.79	.559	1.266	1.973	2.680	3.387	4.094	4.801	5.508	6.215	6.922
.80	.566	1.273	1.980	2.687	3.394	4.101	4.808	5.515	6.222	6.929
.81	.573	1.280	1.987	2.694	3.401	4.108	4.815	5.522	6.229	6.936
.82	.580	1.287	1.994	2.701	3.408	4.115	4.822	5.529	6.236	6.943
.83	.587	1.294	2.001	2.708	3.415	4.122	4.829	5.536	6.243	6.950
.84	.594	1.301	2.008	2.715	3.422	4.129	4.836	5.543	6.250	6.957
.85	.601	1.308	2.015	2.722	3.429	4.136	4.843	5.550	6.257	6.964
.86	.608	1.315	2.022	2.729	3.436	4.143	4.850	5.557	6.264	6.971
.87	.615	1.322	2.029	2.736	3.443	4.150	4.857	5.564	6.271	6.978
.88	.622	1.329	2.036	2.743	3.450	4.157	4.864	5.571	6.278	6.985
.89	.629	1.336	2.043	2.750	3.457	4.164	4.871	5.578	6.285	6.992
.90	.636	1.343	2.050	2.757	3.464	4.171	4.878	5.585	6.292	6.999
.91	.643	1.350	2.057	2.764	3.471	4.178	4.885	5.592	6.299	7.006
.92	.650	1.357	2.064	2.771	3.478	4.185	4.892	5.599	6.306	7.013
.93	.658	1.365	2.072	2.779	3.486	4.193	4.900	5.607	6.314	7.021
.94	.665	1.372	2.079	2.786	3.493	4.200	4.907	5.614	6.321	7.028
.95	.672	1.379	2.086	2.793	3.500	4.207	4.914	5.621	6.328	7.035
.96	.679	1.386	2.093	2.800	3.507	4.214	4.921	5.628	6.335	7.042
.97	.686	1.393	2.100	2.807	3.514	4.221	4.928	5.635	6.342	7.049
.98	.693	1.400	2.107	2.814	3.521	4.228	4.935	5.642	6.349	7.056
.99	.700	1.407	2.114	2.821	3.528	4.235	4.942	5.649	6.356	7.063
1.00	.707	1.414	2.121	2.828	3.535	4.242	4.949	5.656	6.363	7.070
	0	1	2	3	4	5	6	7	8	9

Table 19.—*Products for Form 194—Continued*[Multiplier =  $\sin 60^\circ = 0.866$ ]

	0	1	2	3	4	5	6	7	8	9
.00	0.000	0.866	1.732	2.598	3.464	4.330	5.196	6.062	6.928	7.794
.01	.009	.875	1.741	2.607	3.473	4.339	5.205	6.071	6.937	7.803
.02	.017	.883	1.749	2.615	3.481	4.347	5.213	6.079	6.945	7.811
.03	.026	.892	1.758	2.624	3.490	4.356	5.222	6.088	6.954	7.820
.04	.035	.901	1.767	2.633	3.499	4.365	5.231	6.097	6.963	7.829
.05	.043	.909	1.775	2.641	3.507	4.373	5.239	6.105	6.971	7.837
.06	.052	.918	1.784	2.650	3.516	4.382	5.248	6.114	6.980	7.846
.07	.061	.927	1.793	2.659	3.525	4.391	5.257	6.123	6.989	7.855
.08	.069	.935	1.801	2.667	3.533	4.399	5.265	6.131	6.997	7.863
.09	.078	.944	1.810	2.676	3.542	4.408	5.274	6.140	7.006	7.872
.10	.087	.953	1.819	2.685	3.551	4.417	5.283	6.149	7.015	7.881
.11	.095	.961	1.827	2.693	3.559	4.425	5.291	6.157	7.023	7.889
.12	.104	.970	1.836	2.702	3.568	4.434	5.300	6.166	7.032	7.898
.13	.113	.979	1.845	2.711	3.577	4.443	5.309	6.175	7.041	7.907
.14	.121	.987	1.853	2.719	3.585	4.451	5.317	6.183	7.049	7.915
.15	.130	.996	1.862	2.728	3.594	4.460	5.326	6.192	7.058	7.924
.16	.139	1.005	1.871	2.737	3.603	4.469	5.335	6.201	7.067	7.933
.17	.147	1.013	1.879	2.745	3.611	4.477	5.343	6.209	7.075	7.941
.18	.156	1.022	1.888	2.754	3.620	4.486	5.352	6.218	7.084	7.950
.19	.165	1.031	1.897	2.763	3.629	4.495	5.361	6.227	7.093	7.959
.20	.173	1.039	1.905	2.771	3.637	4.503	5.369	6.235	7.101	7.967
.21	.182	1.048	1.914	2.780	3.646	4.512	5.378	6.244	7.110	7.976
.22	.191	1.057	1.923	2.789	3.655	4.521	5.387	6.253	7.119	7.985
.23	.199	1.065	1.931	2.797	3.663	4.529	5.395	6.261	7.127	7.993
.24	.208	1.074	1.940	2.806	3.672	4.538	5.404	6.270	7.136	8.002
.25	.216	1.082	1.948	2.814	3.680	4.546	5.412	6.278	7.144	8.010
.26	.225	1.091	1.957	2.823	3.689	4.555	5.421	6.287	7.153	8.019
.27	.234	1.100	1.966	2.832	3.698	4.564	5.430	6.296	7.162	8.028
.28	.242	1.108	1.974	2.840	3.706	4.572	5.438	6.304	7.170	8.036
.29	.251	1.117	1.983	2.849	3.715	4.581	5.447	6.313	7.179	8.045
.30	.260	1.126	1.992	2.858	3.724	4.590	5.456	6.322	7.188	8.054
.31	.268	1.134	2.000	2.866	3.732	4.598	5.464	6.330	7.196	8.062
.32	.277	1.143	2.009	2.875	3.741	4.607	5.473	6.339	7.205	8.071
.33	.286	1.152	2.018	2.884	3.750	4.616	5.482	6.348	7.214	8.080
.34	.294	1.160	2.026	2.892	3.758	4.624	5.490	6.356	7.222	8.088
.35	.303	1.169	2.035	2.901	3.767	4.633	5.499	6.365	7.231	8.097
.36	.312	1.178	2.044	2.910	3.776	4.642	5.508	6.374	7.240	8.106
.37	.320	1.186	2.052	2.918	3.784	4.650	5.516	6.382	7.248	8.114
.38	.329	1.195	2.061	2.927	3.793	4.659	5.525	6.391	7.257	8.123
.39	.338	1.204	2.070	2.936	3.802	4.668	5.534	6.400	7.266	8.132
.40	.346	1.212	2.078	2.944	3.810	4.676	5.542	6.408	7.274	8.140
.41	.355	1.221	2.087	2.953	3.819	4.685	5.551	6.417	7.283	8.149
.42	.364	1.230	2.096	2.962	3.828	4.694	5.560	6.426	7.292	8.158
.43	.372	1.238	2.104	2.970	3.836	4.702	5.568	6.434	7.300	8.166
.44	.381	1.247	2.113	2.979	3.845	4.711	5.577	6.443	7.309	8.175
.45	.390	1.256	2.122	2.988	3.854	4.720	5.586	6.452	7.318	8.184
.46	.398	1.264	2.130	2.996	3.862	4.728	5.594	6.460	7.326	8.192
.47	.407	1.273	2.139	3.005	3.871	4.737	5.603	6.469	7.335	8.201
.48	.416	1.282	2.148	3.014	3.880	4.746	5.612	6.478	7.344	8.210
.49	.424	1.290	2.156	3.022	3.888	4.754	5.620	6.486	7.352	8.218
.50	.433	1.299	2.165	3.031	3.897	4.763	5.629	6.495	7.361	8.227
	0	1	2	3	4	5	6	7	8	9

Table 19.—*Products for Form 194—Continued*[Multiplier =  $\sin 60^\circ = 0.866$ ]

	0	1	2	3	4	5	6	7	8	9
0.50.	0.433	1.299	2.165	3.031	3.897	4.763	5.629	6.495	7.361	8.227
.51.	.442	1.308	2.174	3.040	3.906	4.772	5.638	6.504	7.370	8.236
.52.	.450	1.316	2.182	3.048	3.914	4.780	5.646	6.512	7.378	8.244
.53.	.459	1.325	2.191	3.057	3.923	4.789	5.655	6.521	7.387	8.253
.54.	.468	1.334	2.200	3.066	3.932	4.798	5.664	6.530	7.396	8.262
.55.	.476	1.342	2.208	3.074	3.940	4.806	5.672	6.538	7.404	8.270
.56.	.485	1.351	2.217	3.083	3.949	4.815	5.681	6.547	7.413	8.279
.57.	.494	1.360	2.226	3.092	3.958	4.824	5.690	6.556	7.422	8.288
.58.	.502	1.368	2.234	3.100	3.966	4.832	5.698	6.564	7.430	8.296
.59.	.511	1.377	2.243	3.109	3.975	4.841	5.707	6.573	7.439	8.305
.60.	.520	1.386	2.252	3.118	3.984	4.850	5.716	6.582	7.448	8.314
.61.	.528	1.394	2.260	3.126	3.992	4.858	5.724	6.590	7.456	8.322
.62.	.537	1.403	2.269	3.135	4.001	4.867	5.733	6.599	7.465	8.331
.63.	.546	1.412	2.278	3.144	4.010	4.876	5.742	6.608	7.474	8.340
.64.	.554	1.420	2.286	3.152	4.018	4.884	5.750	6.616	7.482	8.348
.65.	.563	1.429	2.295	3.161	4.027	4.893	5.759	6.625	7.491	8.357
.66.	.572	1.438	2.304	3.170	4.036	4.902	5.768	6.634	7.500	8.366
.67.	.580	1.446	2.312	3.178	4.044	4.910	5.776	6.642	7.508	8.374
.68.	.589	1.455	2.321	3.187	4.053	4.919	5.785	6.651	7.517	8.383
.69.	.598	1.464	2.330	3.196	4.062	4.928	5.794	6.660	7.526	8.392
.70.	.606	1.472	2.338	3.204	4.070	4.936	5.802	6.668	7.534	8.400
.71.	.615	1.481	2.347	3.213	4.079	4.945	5.811	6.677	7.543	8.409
.72.	.624	1.490	2.356	3.222	4.088	4.954	5.820	6.686	7.552	8.418
.73.	.632	1.498	2.364	3.230	4.096	4.962	5.828	6.694	7.560	8.426
.74.	.641	1.507	2.373	3.239	4.105	4.971	5.837	6.703	7.569	8.435
.75.	.650	1.516	2.382	3.248	4.114	4.980	5.846	6.712	7.578	8.444
.76.	.658	1.524	2.390	3.256	4.122	4.988	5.854	6.720	7.586	8.452
.77.	.667	1.533	2.399	3.265	4.131	4.997	5.863	6.729	7.595	8.461
.78.	.675	1.541	2.407	3.273	4.139	5.005	5.871	7.737	7.603	8.469
.79.	.684	1.550	2.416	3.282	4.148	5.014	5.880	6.746	7.612	8.478
.80.	.693	1.559	2.425	3.291	4.157	5.023	5.889	6.755	7.621	8.487
.81.	.701	1.567	2.433	3.299	4.165	5.031	5.897	6.763	7.629	8.495
.82.	.710	1.576	2.442	3.308	4.174	5.040	5.906	6.772	7.638	8.504
.83.	.719	1.585	2.451	3.317	4.183	5.049	5.915	6.781	7.647	8.513
.84.	.727	1.593	2.459	3.325	4.191	5.057	5.923	6.789	7.655	8.521
.85.	.736	1.602	2.468	3.334	4.200	5.066	5.932	6.798	7.664	8.530
.86.	.745	1.611	2.477	3.343	4.209	5.075	5.941	6.807	7.673	8.539
.87.	.753	1.619	2.485	3.351	4.217	5.083	5.949	6.815	7.681	8.547
.88.	.762	1.628	2.494	3.360	4.226	5.092	5.958	6.824	7.690	8.556
.89.	.771	1.637	2.503	3.369	4.235	5.101	5.967	6.833	7.699	8.565
.90.	.779	1.645	2.511	3.377	4.243	5.109	5.975	6.841	7.707	8.573
.91.	.788	1.654	2.520	3.386	4.252	5.118	5.984	6.850	7.716	8.582
.92.	.797	1.663	2.529	3.395	4.261	5.127	5.993	6.859	7.725	8.591
.93.	.805	1.671	2.537	3.403	4.269	5.135	6.001	6.867	7.733	8.599
.94.	.814	1.680	2.546	3.412	4.278	5.144	6.010	6.876	7.742	8.608
.95.	.823	1.689	2.555	3.421	4.287	5.153	6.019	6.885	7.751	8.617
.96.	.831	1.697	2.563	3.429	4.295	5.161	6.027	6.893	7.759	8.625
.97.	.840	1.706	2.572	3.438	4.304	5.170	6.036	6.902	7.768	8.634
.98.	.849	1.715	2.581	3.447	4.313	5.179	6.045	6.911	7.777	8.643
.99.	.857	1.723	2.589	3.455	4.321	5.187	6.053	6.919	7.785	8.651
1.00.	.866	1.732	2.598	3.464	4.330	5.196	6.062	6.928	7.794	8.660
	0	1	2	3	4	5	6	7	8	9

Table 19.—*Products for Form 194—Continued*[Multiplier= $\sin 75^\circ = 0.966$ ]

	0	1	2	3	4	5	6	7	8	9
.00.	0.000	0.966	1.932	2.898	3.864	4.830	5.796	6.762	7.728	8.694
.01.	.010	.976	1.942	2.908	3.874	4.840	5.806	6.772	7.738	8.704
.02.	.019	.985	1.951	2.917	3.883	4.849	5.815	6.781	7.747	8.713
.03.	.029	.995	1.961	2.927	3.893	4.859	5.825	6.791	7.757	8.723
.04.	.039	1.005	1.971	2.937	3.903	4.869	5.835	6.801	7.767	8.733
.05.	.048	1.014	1.980	2.946	3.912	4.878	5.844	6.810	7.776	8.742
.06.	.058	1.024	1.990	2.956	3.922	4.888	5.854	6.820	7.786	8.752
.07.	.068	1.034	2.000	2.966	3.932	4.898	5.864	6.830	7.796	8.762
.08.	.077	1.043	2.009	2.975	3.941	4.907	5.873	6.839	7.805	8.771
.09.	.087	1.053	2.019	2.985	3.951	4.917	5.883	6.849	7.815	8.781
.10.	.097	1.063	2.029	2.995	3.961	4.927	5.893	6.859	7.825	8.791
.11.	.106	1.072	2.038	3.004	3.970	4.936	5.902	6.868	7.834	8.800
.12.	.116	1.082	2.048	3.014	3.980	4.946	5.912	6.878	7.844	8.810
.13.	.126	1.092	2.058	3.024	3.990	4.956	5.922	6.888	7.854	8.820
.14.	.135	1.101	2.067	3.033	3.999	4.965	5.931	6.897	7.863	8.829
.15.	.145	1.111	2.077	3.043	4.009	4.975	5.941	6.907	7.873	8.839
.16.	.155	1.121	2.087	3.053	4.019	4.985	5.951	6.917	7.883	8.849
.17.	.164	1.130	2.096	3.062	4.028	4.994	5.960	6.926	7.892	8.858
.18.	.174	1.140	2.106	3.072	4.038	5.004	5.970	6.936	7.902	8.868
.19.	.184	1.150	2.116	3.082	4.048	5.014	5.980	6.946	7.912	8.878
.20.	.193	1.159	2.125	3.091	4.057	5.023	5.989	6.955	7.921	8.887
.21.	.203	1.169	2.135	3.101	4.067	5.033	5.999	6.965	7.931	8.897
.22.	.213	1.179	2.145	3.111	4.077	5.043	6.009	6.975	7.941	8.907
.23.	.222	1.188	2.154	3.120	4.086	5.052	6.018	6.984	7.950	8.916
.24.	.232	1.198	2.164	3.130	4.096	5.062	6.028	6.994	7.960	8.926
.25.	.242	1.208	2.174	3.140	4.106	5.072	6.038	7.004	7.970	8.936
.26.	.251	1.217	2.183	3.149	4.115	5.081	6.047	7.013	7.979	8.945
.27.	.261	1.227	2.193	3.159	4.125	5.091	6.057	7.023	7.989	8.955
.28.	.270	1.236	2.202	3.168	4.134	5.100	6.066	7.032	7.998	8.964
.29.	.280	1.246	2.212	3.178	4.144	5.110	6.076	7.042	8.008	8.974
.30.	.290	1.256	2.222	3.188	4.154	5.120	6.086	7.052	8.018	8.984
.31.	.299	1.265	2.231	3.197	4.163	5.129	6.095	7.061	8.027	8.993
.32.	.309	1.275	2.241	3.207	4.173	5.139	6.105	7.071	8.037	9.003
.33.	.319	1.285	2.251	3.217	4.183	5.149	6.115	7.081	8.047	9.013
.34.	.328	1.294	2.260	3.226	4.192	5.158	6.124	7.090	8.056	9.022
.35.	.338	1.304	2.270	3.236	4.202	5.168	6.134	7.100	8.066	9.032
.36.	.348	1.314	2.280	3.246	4.212	5.178	6.144	7.110	8.076	9.042
.37.	.357	1.323	2.289	3.255	4.221	5.187	6.153	7.119	8.085	9.051
.38.	.367	1.333	2.299	3.265	4.231	5.197	6.163	7.129	8.095	9.061
.39.	.377	1.343	2.309	3.275	4.241	5.207	6.173	7.139	8.105	9.071
.40.	.386	1.352	2.318	3.284	4.250	5.216	6.182	7.148	8.114	9.080
.41.	.396	1.362	2.328	3.294	4.260	5.226	6.192	7.158	8.124	9.090
.42.	.406	1.372	2.338	3.304	4.270	5.236	6.202	7.168	8.134	9.100
.43.	.415	1.381	2.347	3.313	4.279	5.245	6.211	7.177	8.143	9.109
.44.	.425	1.391	2.357	3.323	4.289	5.255	6.221	7.187	8.153	9.119
.45.	.435	1.401	2.367	3.333	4.299	5.265	6.231	7.197	8.163	9.129
.46.	.444	1.410	2.376	3.342	4.308	5.274	6.240	7.206	8.172	9.138
.47.	.454	1.420	2.386	3.352	4.318	5.284	6.250	7.216	8.182	9.148
.48.	.464	1.430	2.396	3.362	4.328	5.294	6.260	7.226	8.192	9.158
.49.	.473	1.439	2.405	3.371	4.337	5.303	6.269	7.235	8.201	9.167
.50.	.483	1.449	2.415	3.381	4.347	5.313	6.279	7.245	8.211	9.177
	0	1	2	3	4	5	6	7	8	9

Table 19.—*Products for Form 194—Continued*[Multiplier= $\sin 75^\circ = 0.966$ ]

	0	1	2	3	4	5	6	7	8	9
0.50	0.483	1.449	2.415	3.381	4.347	5.313	6.279	7.245	8.211	9.177
.51	.493	1.459	2.425	3.391	4.357	5.323	6.289	7.255	8.221	9.187
.52	.502	1.468	2.434	3.400	4.366	5.332	6.298	7.264	8.230	9.196
.53	.512	1.478	2.444	3.410	4.376	5.342	6.308	7.274	8.240	9.206
.54	.522	1.488	2.454	3.420	4.386	5.352	6.318	7.284	8.250	9.216
.55	.531	1.497	2.463	3.429	4.395	5.361	6.327	7.293	8.259	9.225
.56	.541	1.507	2.473	3.439	4.405	5.371	6.337	7.303	8.269	9.235
.57	.551	1.517	2.483	3.449	4.415	5.381	6.347	7.313	8.279	9.245
.58	.560	1.526	2.492	3.458	4.424	5.390	6.356	7.322	8.288	9.254
.59	.570	1.536	2.502	3.468	4.434	5.400	6.366	7.332	8.298	9.264
.60	.580	1.546	2.512	3.478	4.444	5.410	6.376	7.342	8.308	9.274
.61	.589	1.555	2.521	3.487	4.453	5.419	6.385	7.351	8.317	9.283
.62	.599	1.565	2.531	3.497	4.463	5.429	6.395	7.361	8.327	9.293
.63	.609	1.575	2.541	3.507	4.473	5.439	6.405	7.371	8.337	9.303
.64	.618	1.584	2.550	3.516	4.482	5.448	6.414	7.380	8.346	9.312
.65	.628	1.594	2.560	3.526	4.492	5.458	6.424	7.390	8.356	9.322
.66	.638	1.604	2.570	3.536	4.502	5.468	6.434	7.400	8.366	9.332
.67	.647	1.613	2.579	3.545	4.511	5.477	6.443	7.409	8.375	9.341
.68	.657	1.623	2.589	3.555	4.521	5.487	6.453	7.419	8.385	9.351
.69	.667	1.633	2.599	3.565	4.531	5.497	6.463	7.429	8.395	9.361
.70	.676	1.642	2.608	3.574	4.540	5.506	6.472	7.438	8.404	9.370
.71	.686	1.652	2.618	3.584	4.550	5.516	6.482	7.448	8.414	9.380
.72	.696	1.662	2.628	3.594	4.560	5.526	6.492	7.458	8.424	9.390
.73	.705	1.671	2.637	3.603	4.569	5.535	6.501	7.467	8.433	9.399
.74	.715	1.681	2.647	3.613	4.579	5.545	6.511	7.477	8.443	9.409
.75	.724	1.690	2.656	3.622	4.588	5.554	6.520	7.486	8.452	9.418
.76	.734	1.700	2.666	3.632	4.598	5.564	6.530	7.496	8.462	9.428
.77	.744	1.710	2.676	3.642	4.608	5.574	6.540	7.506	8.472	9.438
.78	.753	1.719	2.685	3.651	4.617	5.583	6.549	7.515	8.481	9.447
.79	.763	1.729	2.695	3.661	4.627	5.593	6.559	7.525	8.491	9.457
.80	.773	1.739	2.705	3.671	4.637	5.603	6.569	7.535	8.501	9.467
.81	.782	1.748	2.714	3.680	4.646	5.612	6.578	7.544	8.510	9.476
.82	.792	1.758	2.724	3.690	4.656	5.622	5.588	7.554	8.520	9.486
.83	.802	1.768	2.734	3.700	4.666	5.632	6.598	7.564	8.530	9.496
.84	.811	1.777	2.743	3.709	4.675	5.641	6.607	7.573	8.539	9.505
.85	.821	1.787	2.753	3.719	4.685	5.651	6.617	7.583	8.549	9.515
.86	.831	1.797	2.763	3.729	4.695	5.661	6.627	7.593	8.559	9.525
.87	.840	1.806	2.772	3.738	4.704	5.670	6.636	7.602	8.568	9.534
.88	.850	1.816	2.782	3.748	4.714	5.680	6.646	7.612	8.578	9.544
.89	.860	1.826	2.792	3.758	4.724	5.690	6.656	7.622	8.588	9.554
.90	.869	1.835	2.801	3.767	4.733	5.699	6.665	7.631	8.597	9.563
.91	.879	1.845	2.811	3.777	4.743	5.709	6.675	7.641	8.607	9.573
.92	.889	1.855	2.821	3.787	4.753	5.719	6.685	7.651	8.617	9.583
.93	.898	1.864	2.830	3.796	4.762	5.728	6.694	7.660	8.626	9.592
.94	.908	1.874	2.840	3.806	4.772	5.738	6.704	7.670	8.636	9.602
.95	.918	1.884	2.850	3.816	4.782	5.748	6.714	7.680	8.646	9.612
.96	.927	1.893	2.859	3.825	4.791	5.757	6.723	7.689	8.655	9.621
.97	.937	1.903	2.869	3.835	4.801	5.767	6.733	7.699	8.665	9.631
.98	.947	1.913	2.879	3.845	4.811	5.777	6.743	7.709	8.675	9.641
.99	.956	1.922	2.888	3.854	4.820	5.786	6.752	7.718	8.684	9.650
1.00	.966	1.932	2.898	3.864	4.830	5.796	6.762	7.728	8.694	9.660
	0	1	2	3	4	5	6	7	8	9

Table 20.—*Augmenting factors*

SHORT-PERIOD CONSTITUENTS,\* FORMULA (308)

	Augmenting factor	Logarithm	Remarks
Diurnal $J_1$ , $K_1$ , $M_1$ , $O_1$ , $OO$ , $P_1$ , $Q_1$ , $2Q_1$ , $\rho_1$	1.0029	0.001241	
Semidiurnal $K_2$ , $L_2$ , $M_2$ , $N_2$ , $2N$ , $R_2$ , $T_2$ , $\lambda_2$ , $\mu_2$ , $\nu_2$ , $2SM$ .	1.0115	0.004972	
Terdiurnal $M_3$ , $MK$ , $2MK$	1.0262	0.011220	
Quarter-diurnal $M_4$ , $MN$ , $MS$	1.0472	0.020029	
Sixth-diurnal $M_6$	1.1107	0.045605	
Eighth-diurnal $M_8$	1.2092	0.082498	Each tabulated solar hourly height used once and once only in summation; group covers one constituent hour; constituent day represented by 24 means.

SHORT-PERIOD CONSTITUENTS,\* FORMULA (309)

	Augmenting factor	Logarithm		Augmenting factor	Logarithm	Remarks
$J_1$ -----	1.0031	0.00134	$P_1$ -----	1.0028	0.00123	
$K_1$ -----	1.0029	0.00125	$Q_1$ -----	1.0023	0.00099	
$K_2$ -----	1.0116	0.00500	$2Q$ -----	1.0021	0.00091	
$L_2$ -----	1.0112	0.00482	$R_2$ -----	1.0115	0.00499	
$M_1$ -----	1.0027	0.00116	$T_2$ -----	1.0115	0.00496	
$M_1$ -----	1.0107	0.00464	$\lambda_2$ -----	1.0111	0.00479	Each constituent hour of observation period receives one and only one of solar hourly heights in the summation; group covers one solar hour; each constituent day represented by 24 means.
$M_3$ -----	1.0244	0.01047	$\mu_2$ -----	1.0100	0.00432	
$M_4$ -----	1.0440	0.01868	$\nu_2$ -----	1.0104	0.00449	
$M_6$ -----	1.1028	0.04251	$p_1$ -----	1.0023	0.00100	
$M_8$ -----	1.1934	0.07680	$MK$ -----	1.0250	0.01074	
$N_2$ -----	1.0103	0.00447	$2MK$ -----	1.0238	0.01021	
$2N$ -----	1.0099	0.00430	$MN$ -----	1.0431	0.01833	
$O_1$ -----	1.0025	0.00107	$MS$ -----	1.0456	0.01935	
$OO$ -----	1.0033	0.00144	$2SM$ -----	1.0123	0.00532	

LONG-PERIOD CONSTITUENTS, FORMULA (403)

	Augmenting factor	Logarithm	Remarks
$Mm$ -----	1.0050	0.00218	
$Mf$ -----	1.0205	0.00880	
$MSf$ -----	1.0192	0.00825	Daily sums used as units in the summation for the divisional means, and all daily sums used; constituent month for $Mm$ , $Mf$ , $MSf$ , and constituent year for $Sa$ and $Ssa$ represented by 24 means.
$Sa$ -----	1.0029	0.00124	
$Ssa$ -----	1.0115	0.00497	

ANNUAL AND SEMIANNUAL CONSTITUENTS, FORMULA (404)

	Augmenting factor	Logarithm	Remarks
$Sa$ -----	1.0115	0.00497	
$Ssa$ -----	1.0472	0.02003	For analysis of 12 monthly means.

\*For constituents  $S_1$ ,  $S_2$ ,  $S_3$ , etc., augmenting factor is unity.

Table 21.—Acceleration in epoch of  $K_1$  due to  $P_1$ [Argument  $h - \frac{1}{2}\nu'$  refers to beginning of series]

Series		14 days	29 days	58 days	87 days	105 days	134 days	163 days	192 days	221 days	250 days	279 days	297 days	326 days	
$h - \frac{1}{2}\nu'$															
0	180	+6.5	+11.4		+14.6	+12.6	+10.1	+5.1	+0.9	+0.2	+2.4	+3.9	+3.9	+3.3	+1.6
10	190	+13.9	+16.4	+16.0	+12.0	+8.8	+3.4	-0.1	+0.7	+3.0	+4.1	+3.6	+2.7	+0.9	
20	200	+17.9	+18.3	+15.3	+9.9	+6.4	+1.0	-1.0	+0.9	+3.2	+3.7	+2.8	+1.7	0.0	
30	210	+19.0	+17.6	+12.9	+6.7	+3.1	-1.6	-1.7	+1.0	+2.9	+3.0	+1.7	+0.5	-1.0	
40	220	+17.6	+15.2	+9.4	+2.8	-0.5	-3.8	-2.1	+1.0	+2.4	+2.0	+0.3	-0.8	-1.7	
50	230	+14.7	+11.7	+5.2	-1.2	-4.1	-5.4	-2.1	+0.8	+1.7	+0.8	-1.0	-2.0	-2.0	
60	240	+10.8	+7.4	+0.7	-5.2	-7.2	-6.0	-1.9	+0.6	+0.9	-0.4	-2.2	-2.9	-2.1	
70	250	+6.4	+2.7	-3.9	-8.7	-9.3	-5.8	-1.5	+0.4	+0.1	-1.5	-3.2	-3.4	-1.9	
80	260	+1.5	-2.2	-8.2	-11.3	-10.2	-5.0	-1.0	+0.1	-0.8	-2.6	-3.8	-3.3	-1.5	
90	270	-3.5	-6.9	-12.0	-12.5	-9.7	-3.8	-0.5	-0.1	-1.6	-3.5	-3.9	-2.9	-1.1	
100	280	-8.2	-11.3	-14.7	-12.1	-8.1	-2.3	0.0	-0.4	-2.3	-4.0	-3.5	-2.2	-0.5	
110	290	-12.5	-14.9	-16.0	-10.1	-5.6	-0.7	+0.6	-0.6	-2.8	-4.0	-2.7	-1.3	0.6	
120	300	-16.0	-17.5	-15.2	-6.9	-2.7	+1.0	+1.1	-0.8	-3.1	-3.5	-1.6	-0.4	+0.6	
130	310	-18.4	-18.3	-12.2	-2.9	+0.5	+2.6	+1.5	-1.0	-3.1	-2.5	-0.3	+0.7	+1.1	
140	320	-18.9	-16.7	-7.2	+1.3	+3.6	+4.1	+1.9	-1.0	-2.5	-1.1	+0.9	+1.6	+1.6	
150	330	-16.8	-12.1	-1.0	+5.4	+6.4	+5.2	+2.1	-0.9	-1.5	+0.5	+2.1	+2.5	+1.9	
160	340	-11.3	-4.7	+5.4	+8.9	+8.6	+5.9	+2.0	-0.7	-0.1	+1.9	+3.1	+3.1	+2.1	
170	350	-2.8	+3.9	+10.9	+11.5	+10.0	+5.9	+1.7	-0.3	+1.2	+3.1	+3.8	+3.4	+2.0	
180	360	+6.5	+11.4	+14.6	+12.6	+10.1	+5.1	+0.9	+0.2	+2.4	+3.9	+3.9	+3.3	+1.6	

Table 22.—Ratio of increase in amplitude of  $K_1$  due to  $P_1$ [Argument  $h - \frac{1}{2}\nu'$  refers to beginning of series]

Series		14 days	29 days	58 days	87 days	105 days	134 days	163 days	192 days	221 days	250 days	279 days	297 days	326 days
$h - \frac{1}{2}\nu'$														
0	180	-0.31	-0.26	-0.12	+0.01	+0.06	+0.09	+0.06	+0.01	-0.02	0.00	+0.03	+0.05	+0.05
10	190	-0.25	-0.17	-0.02	+0.09	+0.12	+0.12	+0.07	+0.01	0.00	+0.02	+0.06	+0.07	+0.06
20	200	-0.15	-0.06	+0.07	+0.16	+0.17	+0.13	+0.06	+0.02	+0.02	+0.05	+0.08	+0.08	+0.07
30	210	-0.04	+0.04	+0.16	+0.20	+0.20	+0.13	+0.05	+0.02	+0.04	+0.07	+0.09	+0.09	+0.06
40	220	+0.07	+0.14	+0.23	+0.23	+0.21	+0.12	+0.04	+0.03	+0.05	+0.08	+0.09	+0.09	+0.05
50	230	+0.17	+0.23	+0.27	+0.24	+0.19	+0.09	+0.03	+0.03	+0.06	+0.09	+0.09	+0.08	+0.04
60	240	+0.25	+0.28	+0.29	+0.22	+0.15	+0.05	+0.02	+0.04	+0.07	+0.09	+0.08	+0.06	+0.03
70	250	+0.30	+0.31	+0.28	+0.18	+0.10	+0.02	+0.01	+0.04	+0.04	+0.08	+0.09	+0.07	+0.04
80	260	+0.33	+0.32	+0.24	+0.12	+0.05	-0.01	0.00	+0.04	+0.08	+0.08	+0.04	+0.02	+0.01
90	270	+0.32	+0.29	+0.18	+0.04	-0.02	-0.04	0.00	+0.04	+0.07	+0.06	+0.02	0.00	0.00
100	280	+0.28	+0.23	+0.10	-0.03	-0.07	-0.06	-0.01	+0.04	+0.06	+0.03	0.00	-0.01	0.00
110	290	+0.22	+0.15	+0.01	-0.10	-0.11	-0.07	0.00	+0.04	+0.04	+0.01	-0.02	-0.02	-0.01
120	300	+0.13	+0.05	-0.09	-0.15	-0.14	-0.07	0.00	+0.03	+0.02	-0.01	-0.03	-0.03	0.00
130	310	+0.03	-0.06	-0.17	-0.18	-0.14	-0.06	+0.01	+0.03	0.00	-0.03	-0.04	-0.03	0.00
140	320	-0.08	-0.16	-0.23	-0.19	-0.13	-0.04	+0.02	+0.02	-0.01	-0.04	-0.04	-0.02	+0.01
150	330	-0.19	-0.25	-0.26	-0.17	-0.10	-0.01	+0.03	+0.02	-0.02	-0.04	-0.03	-0.01	+0.02
160	340	-0.28	-0.30	-0.25	-0.12	-0.05	+0.03	+0.04	+0.01	-0.03	-0.03	-0.01	+0.01	+0.03
170	350	-0.32	-0.31	-0.19	-0.06	0.00	+0.06	+0.05	+0.01	-0.03	-0.02	+0.01	+0.03	+0.04
180	360	-0.31	-0.26	-0.12	+0.01	+0.06	+0.09	+0.06	+0.01	-0.02	0.00	+0.03	+0.05	+0.05

Table 23.—Acceleration in epoch of  $S_2$  due to  $K_2$ [Argument  $h-v''$  refers to beginning of series]

Series		15 days	29 days	58 days	87 days	105 days	134 days	163 days	192 days	221 days	250 days	279 days	297 days	326 days
$h-v''$		°	°	°	°	°	°	°	°	°	°	°	°	°
0	180	+3.2	+5.9	+10.1	+10.4	+8.0	+3.2	+0.4	+0.1	+1.3	+2.9	+3.2	+2.4	+0.9
10	190	+7.2	+9.6	+12.3	+10.0	+6.7	+2.0	0.0	+0.3	+1.9	+3.3	+2.9	+1.9	+0.5
20	200	+10.9	+12.6	+13.2	+8.4	+4.7	-0.6	-0.5	+0.5	+2.4	+3.3	+2.2	+1.1	0.0
30	210	+13.7	+14.6	+12.5	+5.7	+2.3	-0.9	-0.9	+0.7	+2.6	+2.9	+1.3	+0.3	-0.5
40	220	+15.4	+15.0	+9.9	+2.5	-0.4	-2.2	-1.3	+0.8	+2.5	+2.0	+0.3	-0.6	-1.0
50	230	+15.4	+13.5	+5.8	-1.1	-3.0	-3.4	-1.6	+0.8	+2.0	+0.9	-0.8	-1.4	-1.3
60	240	+13.2	+9.6	+0.8	-4.5	-5.4	-4.4	-1.7	+0.7	+1.2	-0.4	-1.8	-2.1	-1.6
70	250	+8.6	+3.7	-4.4	-7.4	-7.2	-4.9	-1.7	+0.5	+0.1	-1.6	-2.6	-2.6	-1.7
80	260	+1.9	-3.0	-8.8	-9.5	-8.3	-4.9	-1.3	+0.2	-1.0	-2.6	-3.1	-2.8	-1.6
90	270	-5.5	-9.1	-11.9	-10.4	-8.3	-4.2	-0.7	-0.2	-1.9	-3.2	-3.3	-2.7	-1.3
100	280	-11.2	-13.2	-13.2	-9.9	-7.3	-2.7	0.0	-0.5	-2.4	-3.4	-3.0	-2.2	-0.7
110	290	-14.6	-15.0	-12.7	-8.2	-5.2	-0.8	+0.8	-0.7	-2.6	-3.1	-2.3	-1.4	0.0
120	300	-15.6	-14.7	-10.9	-5.6	-2.6	+1.2	+1.4	-0.8	-2.4	-2.5	-1.4	-0.4	+0.8
130	310	-14.7	-12.9	-8.0	-2.4	+0.5	+3.1	+1.7	-0.8	-2.0	-1.7	-0.3	+0.7	+1.3
140	320	-12.4	-10.0	-4.4	+1.0	+3.4	+4.4	+1.7	-0.7	-1.5	-0.7	+0.8	+1.6	+1.7
150	330	-9.1	-6.4	-0.6	+4.4	+5.9	+4.9	+1.6	-0.5	-0.8	+0.3	+1.9	+2.4	+1.7
160	340	-5.3	-2.3	+3.3	+7.3	+7.7	+4.8	+1.3	-0.3	-0.1	+1.3	+2.7	+2.8	+1.6
170	350	-1.1	+1.9	+7.0	+9.4	+8.4	+4.2	+0.9	-0.1	+0.7	+2.2	+3.2	+2.8	+1.3
180	360	+3.2	+5.9	+10.1	+10.4	+8.0	+3.2	+0.4	+0.1	+1.3	+2.9	+3.2	+2.4	+0.9

Table 24.—Ratio of increase in amplitude of  $S_2$  due to  $K_2$ [Argument  $h-v''$  refers to beginning of series]

Series		15 days	29 days	58 days	87 days	105 days	134 days	163 days	192 days	221 days	250 days	279 days	297 days	326 days
$h-v''$		°	°	°	°	°	°	°	°	°	°	°	°	°
0	180	+0.26	+0.24	+0.15	+0.03	-0.02	-0.04	-0.01	+0.03	+0.05	+0.04	+0.01	0.00	0.00
10	190	+0.23	+0.10	+0.08	-0.03	-0.06	-0.05	-0.01	+0.03	+0.04	+0.02	0.00	-0.01	-0.01
20	200	+0.18	+0.12	0.00	-0.09	-0.10	-0.06	-0.01	+0.03	+0.03	0.00	-0.02	-0.02	-0.01
30	210	+0.10	+0.04	-0.08	-0.13	-0.12	-0.06	0.00	+0.02	+0.01	-0.01	-0.03	-0.03	-0.01
40	220	+0.01	-0.05	-0.15	-0.15	-0.13	-0.05	0.00	+0.02	0.00	-0.03	-0.04	-0.03	0.00
50	230	-0.08	-0.14	-0.19	-0.16	-0.11	-0.03	+0.01	+0.02	-0.01	-0.04	-0.03	-0.02	0.00
60	240	-0.17	-0.21	-0.21	-0.14	-0.09	-0.01	+0.02	+0.01	-0.02	-0.04	-0.02	-0.01	+0.01
70	250	-0.23	-0.25	-0.20	-0.10	-0.05	+0.02	+0.03	+0.01	-0.03	-0.03	-0.01	0.00	+0.02
80	260	-0.27	-0.25	-0.16	-0.05	0.00	+0.05	+0.04	0.00	-0.02	-0.02	0.00	+0.02	+0.03
90	270	-0.25	-0.21	-0.10	+0.01	+0.05	+0.08	+0.05	0.00	-0.01	0.00	+0.02	+0.04	+0.04
100	280	-0.20	-0.15	-0.02	+0.07	+0.10	+0.10	+0.05	+0.01	0.00	+0.02	+0.04	+0.05	+0.05
110	290	-0.13	-0.06	+0.05	+0.12	+0.14	+0.11	+0.05	+0.01	+0.01	+0.04	+0.06	+0.06	+0.05
120	300	-0.03	+0.03	+0.13	+0.17	+0.16	+0.11	+0.04	+0.01	+0.03	+0.05	+0.07	+0.07	+0.05
130	310	+0.06	+0.11	+0.18	+0.19	+0.17	+0.09	+0.03	+0.02	+0.04	+0.07	+0.08	+0.07	+0.04
140	320	+0.14	+0.18	+0.22	+0.19	+0.15	+0.07	+0.02	+0.02	+0.05	+0.07	+0.07	+0.06	+0.03
150	330	+0.21	+0.23	+0.24	+0.18	+0.13	+0.04	+0.01	+0.03	+0.06	+0.07	+0.06	+0.05	+0.02
160	340	+0.25	+0.26	+0.23	+0.14	+0.08	+0.01	0.00	+0.03	+0.06	+0.07	+0.05	+0.03	+0.01
170	350	+0.27	+0.26	+0.20	+0.09	+0.03	-0.02	0.00	+0.03	+0.06	+0.06	+0.03	+0.02	0.00
180	360	+0.26	+0.24	+0.15	+0.03	-0.02	-0.04	-0.01	+0.03	+0.05	+0.04	+0.01	0.00	0.00

Table 25.—*Acceleration in epoch of S<sub>2</sub> due to T<sub>2</sub>*[Argument  $h-p_1$  refers to beginning of series]

Series.	15 days	29 days	58 days	87 days	105 days	134 days	163 days	192 days	221 days	250 days	279 days	297 days	326 days
$h-p_1$	°	°	°	°	°	°	°	°	°	°	°	°	°
0	-0.4	-0.8	-1.5	-2.0	-2.2	-2.4	-2.3	-2.0	-1.5	-1.1	-0.6	-0.4	-0.1
10	-1.0	-1.3	-1.9	-2.4	-2.5	-2.6	-2.4	-2.0	-1.4	-0.9	-0.5	-0.3	-0.1
20	-1.5	-1.8	-2.2	-2.7	-2.7	-2.7	-2.3	-1.8	-1.3	-0.7	-0.3	-0.2	0.0
30	-2.0	-2.2	-2.7	-2.9	-2.9	-2.7	-2.2	-1.7	-1.1	-0.6	-0.2	0.0	+0.1
40	-2.4	-2.6	-3.0	-3.0	-2.9	-2.6	-2.0	-1.4	-0.8	-0.4	0.0	+0.1	+0.1
50	-2.7	-2.9	-3.1	-3.1	-2.9	-2.4	-1.8	-1.2	-0.6	-0.2	+0.1	+0.2	+0.2
60	-3.0	-3.2	-3.2	-3.0	-2.8	-2.2	-1.5	-0.9	-0.3	+0.1	+0.3	+0.3	+0.2
70	-3.2	-3.3	-3.2	-2.9	-2.5	-1.9	-1.2	-0.5	0.0	+0.3	+0.4	+0.4	+0.3
80	-3.4	-3.3	-3.1	-2.6	-2.2	-1.5	-0.8	-0.2	+0.3	+0.5	+0.6	+0.5	+0.3
90	-3.4	-3.3	-2.9	-2.3	-1.9	-1.1	-0.4	+0.2	+0.5	+0.7	+0.7	+0.6	+0.4
100	-3.3	-3.1	-2.6	-1.9	-1.4	-0.7	0.0	+0.5	+0.8	+0.9	+0.8	+0.6	+0.4
110	-3.1	-2.8	-2.2	-1.4	-1.0	-0.3	+0.4	+0.8	+1.0	+1.0	+0.9	+0.7	+0.4
120	-2.8	-2.5	-1.8	-0.9	-0.4	+0.3	+0.8	+1.1	+1.2	+1.1	+0.9	+0.7	+0.4
130	-2.4	-2.0	-1.2	-0.4	+0.1	+0.8	+1.2	+1.4	+1.4	+1.2	+0.9	+0.7	+0.4
140	-1.9	-1.5	-0.7	+0.2	+0.6	+1.2	+1.5	+1.6	+1.5	+1.3	+0.9	+0.7	+0.3
150	-1.4	-0.9	-0.1	+0.7	+1.1	+1.6	+1.8	+1.8	+1.6	+1.3	+0.9	+0.7	+0.3
160	-0.8	-0.3	+0.5	+1.2	+1.6	+1.9	+2.1	+2.0	+1.7	+1.3	+0.8	+0.6	+0.3
170	-0.2	+0.3	+1.1	+1.7	+2.0	+2.2	+2.2	+2.0	+1.6	+1.2	+0.8	+0.5	+0.2
180	+0.5	+0.9	+1.6	+2.2	+2.3	+2.5	+2.3	+2.0	+1.6	+1.1	+0.7	+0.4	+0.1
190	+1.1	+1.5	+2.1	+2.5	+2.6	+2.6	+2.4	+2.0	+1.5	+1.0	+0.5	+0.3	+0.1
200	+1.6	+2.0	+2.5	+2.8	+2.8	+2.7	+2.3	+1.8	+1.3	+0.8	+0.4	+0.2	0.0
210	+2.1	+2.4	+2.8	+3.0	+3.0	+2.6	+2.2	+1.7	+1.1	+0.6	+0.2	0.0	-0.1
220	+2.6	+2.8	+3.1	+3.1	+2.9	+2.5	+2.0	+1.5	+0.9	+0.4	0.0	-0.1	-0.1
230	+2.9	+3.1	+3.2	+3.0	+2.9	+2.4	+1.8	+1.2	+0.6	+0.2	-0.1	-0.2	-0.2
240	+3.2	+3.3	+3.2	+3.0	+2.7	+2.1	+1.5	+0.9	+0.3	-0.1	-0.3	-0.3	-0.3
250	+3.3	+3.3	+3.2	+2.8	+2.5	+1.8	+1.2	+0.5	0.0	-0.3	-0.4	-0.4	-0.3
260	+3.4	+3.3	+3.0	+2.5	+2.1	+1.5	+0.8	+0.2	-0.3	-0.5	-0.6	-0.5	-0.3
270	+3.3	+3.2	+2.8	+2.2	+1.8	+1.1	+0.4	-0.2	-0.6	-0.7	-0.7	-0.6	-0.4
280	+3.2	+3.0	+2.5	+1.8	+1.4	+0.6	0.0	-0.5	-0.8	-0.9	-0.8	-0.7	-0.4
290	+2.9	+2.7	+2.1	+1.3	+0.9	+0.2	-0.4	-0.8	-1.1	-1.1	-0.9	-0.7	-0.4
300	+2.6	+2.3	+1.6	+0.9	+0.4	-0.3	-0.8	-1.1	-1.3	-1.2	-0.9	-0.7	-0.4
310	+2.2	+1.9	+1.1	+0.4	-0.1	-0.7	-1.2	-1.4	-1.4	-1.3	-0.9	-0.7	-0.4
320	+1.7	+1.4	+0.6	-0.2	-0.6	-1.1	-1.5	-1.7	-1.5	-1.3	-0.9	-0.7	-0.3
330	+1.2	+0.8	+0.1	-0.7	-1.0	-1.5	-1.8	-1.8	-1.6	-1.3	-0.9	-0.6	-0.3
340	+0.7	+0.3	-0.5	-1.1	-1.5	-1.9	-2.0	-2.0	-1.7	-1.3	-0.8	-0.6	-0.2
350	+0.1	-0.2	-1.0	-1.6	-1.9	-2.2	-2.2	-2.0	-1.6	-1.2	-0.7	-0.5	-0.2
360	-0.4	-0.8	-1.5	-2.0	-2.2	-2.4	-2.3	-2.0	-1.5	-1.1	-0.6	-0.4	-0.1

Table 26.—*Resultant amplitude of S<sub>2</sub> due to T<sub>2</sub>*[Argument h—p<sub>1</sub> refers to beginning of series]

Series h—p <sub>1</sub>	15 days	29 days	58 days	87 days	105 days	134 days	163 days	192 days	221 days	250 days	279 days	297 days	326 days
0	1.06	1.06	1.05	1.04	1.03	1.02	1.01	1.00	0.99	0.99	0.99	0.99	0.99
10	1.06	1.05	1.05	1.03	1.03	1.01	1.00	0.99	0.99	0.99	0.99	0.99	0.99
20	1.05	1.05	1.04	1.03	1.02	1.00	0.99	0.99	0.98	0.98	0.99	0.99	0.99
30	1.05	1.04	1.03	1.02	1.01	1.00	0.99	0.98	0.98	0.98	0.98	0.99	0.99
40	1.04	1.03	1.02	1.01	1.00	0.99	0.98	0.98	0.98	0.98	0.98	0.99	0.99
50	1.03	1.03	1.01	1.00	0.99	0.98	0.97	0.97	0.97	0.98	0.98	0.99	1.00
60	1.02	1.02	1.00	0.99	0.98	0.97	0.97	0.97	0.97	0.98	0.99	0.99	1.00
70	1.01	1.01	0.99	0.98	0.97	0.97	0.96	0.97	0.97	0.98	0.99	0.99	1.00
80	1.00	1.00	0.98	0.97	0.97	0.96	0.96	0.97	0.97	0.98	0.99	0.99	1.00
90	1.00	0.99	0.97	0.96	0.96	0.96	0.96	0.97	0.97	0.98	0.99	0.99	1.00
100	0.99	0.98	0.97	0.96	0.96	0.96	0.96	0.97	0.98	0.98	0.99	1.00	1.00
110	0.98	0.97	0.96	0.95	0.95	0.95	0.96	0.97	0.98	0.98	0.99	1.00	1.00
120	0.97	0.96	0.95	0.95	0.95	0.95	0.96	0.97	0.98	0.99	1.00	1.00	1.00
130	0.96	0.95	0.95	0.95	0.95	0.96	0.97	0.98	0.99	0.99	1.00	1.00	1.00
140	0.95	0.95	0.94	0.95	0.95	0.96	0.97	0.98	0.99	0.99	1.00	1.00	1.00
150	0.95	0.94	0.94	0.95	0.95	0.96	0.97	0.99	1.00	1.00	1.01	1.01	1.01
160	0.94	0.94	0.94	0.95	0.96	0.97	0.98	0.99	1.00	1.01	1.01	1.01	1.01
170	0.94	0.94	0.95	0.96	0.96	0.97	0.99	1.00	1.01	1.01	1.01	1.01	1.01
180	0.94	0.94	0.95	0.96	0.97	0.98	0.99	1.00	1.01	1.01	1.01	1.01	1.01
190	0.95	0.95	0.96	0.97	0.98	0.99	1.00	1.01	1.01	1.02	1.02	1.01	1.01
200	0.95	0.95	0.96	0.98	0.99	1.00	1.01	1.02	1.02	1.02	1.02	1.01	1.01
210	0.96	0.96	0.97	0.99	0.99	1.01	1.02	1.02	1.02	1.02	1.02	1.01	1.01
220	0.96	0.97	0.98	1.00	1.00	1.01	1.02	1.03	1.03	1.02	1.02	1.01	1.01
230	0.97	0.98	0.99	1.00	1.01	1.02	1.03	1.03	1.03	1.02	1.02	1.01	1.01
240	0.98	0.99	1.00	1.01	1.01	1.02	1.03	1.03	1.03	1.02	1.02	1.01	1.01
250	0.99	1.00	1.01	1.02	1.03	1.04	1.04	1.04	1.03	1.02	1.02	1.01	1.01
260	1.00	1.01	1.02	1.03	1.04	1.04	1.04	1.04	1.03	1.02	1.01	1.01	1.00
270	1.01	1.02	1.03	1.04	1.04	1.04	1.04	1.04	1.03	1.02	1.01	1.01	1.00
280	1.02	1.03	1.04	1.04	1.05	1.05	1.04	1.04	1.03	1.02	1.01	1.01	1.00
290	1.03	1.03	1.04	1.05	1.05	1.05	1.04	1.03	1.02	1.01	1.01	1.00	1.00
300	1.04	1.04	1.05	1.05	1.05	1.05	1.04	1.03	1.02	1.01	1.00	1.00	1.00
310	1.04	1.05	1.05	1.05	1.05	1.05	1.04	1.03	1.02	1.01	1.00	1.00	1.00
320	1.05	1.05	1.06	1.05	1.05	1.04	1.03	1.02	1.01	1.00	1.00	1.00	1.00
330	1.06	1.06	1.06	1.05	1.05	1.04	1.03	1.02	1.01	1.00	1.00	1.00	1.00
340	1.06	1.06	1.06	1.05	1.05	1.03	1.02	1.01	1.00	1.00	0.99	0.99	1.00
350	1.06	1.06	1.05	1.05	1.04	1.03	1.02	1.00	1.00	0.99	0.99	0.99	1.00
360	1.06	1.06	1.05	1.04	1.03	1.02	1.01	1.00	0.99	0.99	0.99	0.99	0.99

Table 27.—*Critical logarithms for Form 245*

Natural number	Logarithm								
.000	-----	0.050	8.6947	0.100	8.9979	0.150	9.1747	0.200	9.3000
.001	6.6990	.051	8.7033	.101	9.0022	.151	9.1776	.201	9.3022
.002	7.1761	.052	8.7119	.102	9.0065	.152	9.1805	.202	9.3043
.003	7.3980	.053	8.7202	.103	9.0108	.153	9.1833	.203	9.3065
.004	7.5441	.054	8.7284	.104	9.0150	.154	9.1862	.204	9.3086
.005	7.6533	.055	8.7365	.105	9.0192	.155	9.1890	.205	9.3107
.006	7.7404	.056	8.7443	.106	9.0233	.156	9.1918	.206	9.3129
.007	7.8130	.057	8.7521	.107	9.0274	.157	9.1946	.207	9.3150
.008	7.8751	.058	8.7597	.108	9.0315	.158	9.1973	.208	9.3171
.009	7.9295	.059	8.7672	.109	9.0355	.159	9.2001	.209	9.3192
.010	7.9778	.060	8.7746	.110	9.0395	.160	9.2028	.210	9.3212
.011	8.0212	.061	8.7818	.111	9.0434	.161	9.2055	.211	9.3233
.012	8.0607	.062	8.7889	.112	9.0473	.162	9.2082	.212	9.3254
.013	8.0970	.063	8.7959	.113	9.0512	.163	9.2109	.213	9.3274
.014	8.1304	.064	8.8028	.114	9.0551	.164	9.2136	.214	9.3295
.015	8.1614	.065	8.8096	.115	9.0589	.165	9.2162	.215	9.3315
.016	8.1904	.066	8.8163	.116	9.0626	.166	9.2189	.216	9.3335
.017	8.2175	.067	8.8229	.117	9.0664	.167	9.2215	.217	9.3355
.018	8.2431	.068	8.8294	.118	9.0701	.168	9.2241	.218	9.3375
.019	8.2672	.069	8.8357	.119	9.0738	.169	9.2267	.219	9.3395
.020	8.2901	.070	8.8420	.120	9.0774	.170	9.2292	.220	9.3415
.021	8.3118	.071	8.8482	.121	9.0810	.171	9.2318	.221	9.3435
.022	8.3325	.072	8.8544	.122	9.0846	.172	9.2343	.222	9.3454
.023	8.3522	.073	8.8604	.123	9.0882	.173	9.2368	.223	9.3474
.024	8.3711	.074	8.8663	.124	9.0917	.174	9.2394	.224	9.3493
.025	8.3892	.075	8.8722	.125	9.0952	.175	9.2419	.225	9.3513
.026	8.4066	.076	8.8780	.126	9.0987	.176	9.2443	.226	9.3532
.027	8.4233	.077	8.8837	.127	9.1021	.177	9.2468	.227	9.3551
.028	8.4394	.078	8.8894	.128	9.1056	.178	9.2493	.228	9.3570
.029	8.4549	.079	8.8949	.129	9.1090	.179	9.2517	.229	9.3589
.030	8.4699	.080	8.9004	.130	9.1123	.180	9.2541	.230	9.3608
.031	8.4843	.081	8.9059	.131	9.1157	.181	9.2565	.231	9.3627
.032	8.4984	.082	8.9112	.132	9.1190	.182	9.2589	.232	9.3646
.033	8.5119	.083	8.9165	.133	9.1223	.183	9.2613	.233	9.3665
.034	8.5251	.084	8.9217	.134	9.1255	.184	9.2637	.234	9.3683
.035	8.5379	.085	8.9269	.135	9.1288	.185	9.2661	.235	9.3702
.036	8.5503	.086	8.9320	.136	9.1320	.186	9.2684	.236	9.3720
.037	8.5623	.087	8.9371	.137	9.1352	.187	9.2707	.237	9.3739
.038	8.5741	.088	8.9421	.138	9.1384	.188	9.2731	.238	9.3757
.039	8.5855	.089	8.9470	.139	9.1415	.189	9.2754	.239	9.3775
.040	8.5967	.090	8.9519	.140	9.1446	.190	9.2777	.240	9.3794
.041	8.6075	.091	8.9567	.141	9.1477	.191	9.2799	.241	9.3812
.042	8.6181	.092	8.9615	.142	9.1508	.192	9.2822	.242	9.3830
.043	8.6284	.093	8.9662	.143	9.1539	.193	9.2845	.243	9.3848
.044	8.6385	.094	8.9709	.144	9.1569	.194	9.2867	.244	9.3866
.045	8.6484	.095	8.9755	.145	9.1599	.195	9.2890	.245	9.3883
.046	8.6581	.096	8.9801	.146	9.1629	.196	9.2912	.246	9.3901
.047	8.6675	.097	8.9846	.147	9.1659	.197	9.2934	.247	9.3919
.048	8.6767	.098	8.9891	.148	9.1688	.198	9.2956	.248	9.3936
.049	8.6858	.099	8.9935	.149	9.1718	.199	9.2978	.249	9.3954
.050	8.6947	.100	8.9979	.150	9.1747	.200	9.3000	.250	9.3971

Table 28.—Constituent speed differences ( $b-a$ ) and  $\log(b-a)$ 

DIURNAL CONSTITUENTS

<i>A</i>	<i>B</i>	<i>J<sub>1</sub></i>	<i>K<sub>1</sub></i>	<i>M<sub>1</sub></i>	<i>O<sub>1</sub></i>	<i>OO</i>	<i>P<sub>1</sub></i>	<i>Q<sub>1</sub></i>	<i>2Q</i>	<i>S<sub>1</sub></i>	<i>P<sub>1</sub></i>
<i>J<sub>1</sub></i>	0	-0.544375 9.735898	-1.088749 0.036928	-1.642408 0.215481	+0.553658 9.743242	-0.628552 9.796929	-2.186782 0.339806	-2.731147 0.436347	-0.585443 9.767485	-2.113929 0.325090	
<i>K<sub>1</sub></i>	+0.544375 9.735898	0	-0.544375 9.735898	-1.098033 0.040615	+1.098033 0.040615	-0.082137 8.914539	-1.642408 0.215481	-2.186782 0.339806	-0.041069 8.613514	-1.569554 0.195776	
<i>M<sub>1</sub></i>	+1.088749 0.036928	+0.544375 9.735898	0	-0.553658 9.743242	+1.642408 0.215481	+0.462308 9.664865	-1.098033 0.040615	-1.642408 0.215481	+0.503396 9.701832	-1.025179 0.010800	
<i>O<sub>1</sub></i>	+1.642408 0.040615	+1.098033 0.040615	+0.553658 9.743242	0	+2.160606 0.341045	+1.015896 0.006849	-0.544376 9.735898	-1.088749 0.036928	+1.056964 0.024060	-0.471521 9.675501	
<i>OO</i>	-0.553658 -1.098033	-1.642408 0.040615	-2.160606 0.341045	0	-1.180170 0	-1.180170 0.071945	-2.740441 0.437820	-3.284816 0.616511	-1.139102 0.056563	-2.667587 0.426119	
<i>P<sub>1</sub></i>	+0.625512 9.796929	+0.621337 8.914539	-0.462237 9.664865	-1.015896 0.006849	+1.180170 0.071945	0	-1.560270 0.193200	-2.104645 0.323179	+0.041069 8.613514	-1.487447 0.172433	
<i>Q<sub>1</sub></i>	+2.186782 0.339806	+1.612408 0.215481	+1.098033 0.040615	+0.544375 0.437820	+2.740441 0.163200	+1.560270 0	-0.544375 9.755898	+1.601339 0.204483	+0.072854 0.204483	+0.072854 8.862463	
<i>2Q</i>	+2.731147 0.436347	+2.186782 0.339806	+1.642408 0.215481	+1.088749 0.036928	+3.284816 0.516511	+2.104645 0.323179	+0.644375 9.735898	0	+2.145714 0.331572	+0.617228 0.331572	
<i>S<sub>1</sub></i>	+0.586443 9.767485	+0.041069 8.613514	-0.503306 9.701832	-1.056964 0.024060	+1.139102 0.066863	-0.041069 8.613514	-1.601339 0.204483	-2.145714 0.331572	0	-1.528466 0.184261	
<i>P<sub>1</sub></i>	+2.113929 0.325090	+1.025179 0.010800	+1.599554 0.195776	+0.471521 0.436347	+2.667587 0.426119	+1.478417 0.172433	-0.072854 9.750446	-0.617228 8.862463	+1.528466 0.184261	0	

Table 28.—Constituent speed differences ( $b-a$ ) and log ( $b-a$ )—Continued

SEMI DIURNAL CONSTITUENTS												2S M	
A	R	K <sub>2</sub>	L <sub>2</sub>	M <sub>2</sub>	N <sub>2</sub>	2N	R <sub>2</sub>	S <sub>2</sub>	T <sub>2</sub>	λ <sub>2</sub>	μ <sub>2</sub>	ν <sub>2</sub>	ν <sub>2</sub>
K <sub>2</sub>	0	-0.553658 9.743242	-1.069033 0.646615	-1.642408 0.2151581	-2.186782 0.359806	0.041071 8.914559	-0.082137 0.090625	-0.123204 0.325690	-0.626512 9.796229	-2.113929 0.195776	-5.565554 0.1560270	+0.933759 -1.015896	+0.970235 0.006849
L <sub>2</sub>	+0.553658 9.743242	0	-0.544875 9.735898	-1.088749 0.036928	-1.63124 0.213019	+0.512988 0.9709758	+0.471621 9.673501	+0.430454 9.632927	-1.560270 8.862453	-1.015896 0.193200	+1.487417 0.006849	+1.487417 0.172443	+0.970235 0.006849
M <sub>2</sub>	+1.069033 0.40615	+0.544375 9.735898	0	-0.544375 9.735898	-1.088749 0.036928	+1.015696 0.024059	+0.974829 0.006849	+0.471521 9.673501	-0.471521 0.006849	-2.113929 9.673501	-0.105896 0.006849	+2.031792 0.307879	+0.970235 0.006849
N <sub>2</sub>	+1.642408 0.2151581	+1.088749 0.036928	+0.544375 9.735898	0	-0.544375 9.735898	+1.601337 0.204483	+1.560270 0.183200	+1.519204 0.181616	+1.015896 0.006849	+0.072854 9.673501	+0.072854 8.862453	+2.576166 0.411094	+0.970235 0.006849
2N	+2.186782 0.359806	+1.633124 0.213019	+1.088749 0.036928	+0.544375 9.735898	0	+2.104645 0.331571	+2.145712 0.323179	+2.104645 0.314621	+1.560270 0.193200	+0.072854 8.862453	+0.072854 9.790446	+3.120541 0.494230	+0.970235 0.006849
R <sub>2</sub>	+0.041071 8.613635	-0.512588 9.709768	-1.056962 0.024059	-1.601337 0.204483	-2.147112 0.331571	0	-0.041067 8.613493	-0.082133 8.613493	-0.585451 9.767483	-2.072858 0.316570	-1.523484 0.182401	+0.974823 0.988928	+0.970235 0.006849
S <sub>2</sub>	+0.689237 8.914639	-0.471621 9.673501	-1.015896 0.066849	-1.560270 0.193200	-2.109645 0.323217	+0.041067 8.613493	+0.041067 0	-0.041067 8.613493	-0.54375 9.735898	-2.091792 0.307879	-1.487417 0.172433	+1.015896 0.172433	+0.970235 0.006849
T <sub>2</sub>	+0.123204 9.6906325	-0.430454 9.633827	-0.974829 9.988628	-1.519204 0.181616	-2.063579 0.314621	+0.082133 8.914518	+0.041067 8.613493	0	-0.503308 9.701834	-1.980725 0.269011	-1.446350 0.160273	+1.056962 0.024059	+0.970235 0.006849
M	+0.626512 9.796929	+0.072854 8.862453	-0.471621 9.673501	-1.015896 0.066849	-1.560270 0.193200	+0.585451 0.006849	+0.544375 9.701834	+0.503308 0	0	-1.487417 9.701834	-0.93042	+1.560270 0.172433	+0.970235 0.006849
R <sub>2</sub>	+2.113926 0.325690	+1.560270 0.193200	+1.015896 0.066849	+0.471621 9.673501	-0.072854 8.862453	+0.072854 0.316570	+0.072858 0.307879	+1.487417 0.298011	+1.487417 0.172433	0	+0.544375 0	+3.047687 0.735898	+0.970235 0.006849
v <sub>2</sub>	+1.569554 0.185776	+1.015896 0.066849	+0.471621 9.673501	-0.072854 8.613493	-0.617228 9.790446	+1.523484 0.184261	+1.487417 0.172433	+1.446350 0.160273	+0.943042 9.794531	-0.544375 9.755898	0	+2.503313 0.398615	+0.970235 0.006849
2SM	-0.933759 0.970235	-1.487417 0.172433	-2.031792 0.307879	-2.576166 0.410974	-3.120541 0.493230	-0.974829 0.988628	-1.015896 0.066849	-1.056962 0.024059	-1.560270 0.193200	-3.047687 0.483970	-2.503313 0.398615	0	+0.970235 0.006849

Table 29.—*Elimination factors*

[Upper line for each constituent gives the logarithms of the factors; middle line, corresponding natural numbers; lower line, angles in degrees]

## SERIES 14 DAYS. DIURNAL CONSTITUENTS

Constituent sought (A)	Disturbing constituents (B, C, etc.)									
	J <sub>1</sub>	K <sub>1</sub>	M <sub>1</sub>	O <sub>1</sub>	O <sub>O</sub>	P <sub>1</sub>	Q <sub>1</sub>	2Q	S <sub>1</sub>	$\rho_1$
J <sub>1</sub> -----	9.7968 .626 --- ---	8.2015 .016 --- ---	9.3150 .207 --- ---	9.7890 .615 --- ---	9.7203 .525 --- ---	8.3017 .020 --- ---	9.0913 .123 --- ---	9.7607 .576 --- ---	8.1357 .014 --- ---	--- 185 --- ---
K <sub>1</sub> -----	9.7968 .626 91	----- .626 ---	9.7968 .626 269	8.3839 .024 356	8.3839 .024 4	9.9958 .990 ---	9.3150 .207 ---	8.3017 .020 ---	9.9990 .998 ---	9.3344 .216 276
M <sub>1</sub> -----	8.2015 .016 3	9.7968 .626 91	----- .615 ---	9.7800 .207 267	9.3150 .721 96	9.8578 .024 78	8.3839 .024 356	9.3150 .207 264	9.8290 .675 85	8.6530 .045 188
O <sub>1</sub> -----	9.3150 .207 96	8.3839 .024 4	9.7890 .615 93	----- .024 ---	8.3826 .054 9	8.7358 .626 171	9.7968 .016 269	8.2015 .016 357	8.1361 .014 178	9.8516 .711 281
O <sub>O</sub> -----	9.7890 .615 267	8.3839 .024 356	9.3150 .207 264	8.3826 .024 351	----- .091 ---	8.9571 .091 342	9.0878 .122 260	8.3320 .021 348	8.7710 .059 349	9.1065 .128 272
P <sub>1</sub> -----	9.7203 .525 105	9.9958 .990 14	8.9578 .721 282	8.7358 .054 189	8.9571 .091 18	----- .217 278	9.3355 .217 186	8.2581 .018 7	9.9990 .998 290	9.3331 .215 290
Q <sub>1</sub> -----	8.3017 .020 7	9.3150 .207 96	8.3839 .024 4	9.7968 .626 91	9.0878 .122 100	9.3355 .217 82	----- .217 269	9.7968 .626 89	9.3283 .213 12	9.9967 .992 12
2Q-----	9.0913 .123 99	8.3017 .020 7	9.3150 .207 96	8.2015 .016 3	8.3320 .021 12	8.2581 .018 174	9.7968 .626 91	----- .001 0	7.1244 .001 0	9.7298 .537 104
S <sub>1</sub> -----	9.7607 .576 98	9.9990 .998 7	9.8290 .675 275	8.1361 .014 182	8.7710 .059 11	9.9990 .998 353	9.3283 .213 271	7.1244 .001 0	----- .217 283	9.3369 .217 283
P <sub>1</sub> -----	8.1357 .014 175	9.3344 .216 84	8.6530 .045 172	9.8516 .711 79	9.1065 .128 88	9.3331 .215 70	9.9967 .992 348	9.7298 .537 256	9.3369 .217 77	----- --- ---

Table 29.—*Elimination factors—Continued*  
SERIES 15 DAYS. SEMIDIURNAL CONSTITUENTS

Constituent sought ( <i>A</i> )	Disturbing constituents ( <i>B</i> , <i>C</i> , etc.)											
	K <sub>2</sub>	L <sub>2</sub>	M <sub>2</sub>	N <sub>2</sub>	2N	R <sub>2</sub>	S <sub>2</sub>	T <sub>2</sub>	λ <sub>2</sub>	μ <sub>2</sub>	ν <sub>2</sub>	2SM
K <sub>2</sub> -----	9.7534 .567 --- 260	8.9437 .088 --- 342	9.2424 .175 --- 244	8.9063 .081 --- 326	9.9986 .997 --- 353	9.9950 .989 --- 345	9.9892 .975 --- 338	9.6707 .468 --- 247	8.7223 .053 --- 339	9.2966 .198 --- 257	8.8476 .070 --- 168	
L <sub>2</sub> -----	9.7534 .567 100	9.7627 .579 --- 262	8.9055 .080 --- 344	9.2507 .178 --- 246	9.7927 .080 --- 92	9.8276 .054 --- 85	9.8585 .620 --- 77	9.9961 .722 --- 347	9.3018 .200 --- 259	8.1941 .016 --- 357	9.3301 .016 --- 357	
M <sub>2</sub> -----	8.9437 .088 18	9.7627 .579 --- 98	8.9055 .080 --- 262	9.7627 .579 --- 344	8.7291 .080 --- 10	8.1941 .016 --- 3	8.4114 .026 --- 175	9.8276 .672 --- 85	8.1941 .016 --- 357	9.8276 .672 --- 275	8.1935 .016 --- 6	
N <sub>2</sub> -----	9.2424 .175 116	8.9055 .080 16	9.7627 .579 98	----- --- 262	9.7627 .579 --- 108	9.2760 .189 --- 101	9.3018 .200 --- 93	9.3204 .209 --- 3	8.1941 .016 --- 275	9.8276 .672 --- 13	9.9961 .120 --- 104	
2N-----	8.9063 .081 34	9.2507 .178 114	8.9055 .080 16	9.7627 .579 98	----- --- 26	8.8187 .066 --- 19	8.6888 .049 --- 11	8.4856 .031 --- 101	9.3018 .200 --- 101	9.9961 .991 --- 13	9.6823 .481 --- 111	
R <sub>2</sub> -----	9.9986 .997 7	9.7927 .620 268	8.7291 .054 350	9.2760 .189 252	8.8167 .066 334	----- --- ---	9.9987 .997 353	9.9950 .989 345	9.7195 .524 255	8.5420 .035 347	9.3168 .207 265	
S <sub>2</sub> -----	9.9950 .989 15	9.8276 .672 275	8.1941 .016 357	9.3018 .200 259	8.6888 .049 341	9.9987 .997 7	----- --- ---	9.9987 .997 353	9.7627 .579 262	8.1935 .016 354	9.3301 .214 272	
T <sub>2</sub> -----	9.9892 .975 22	9.8585 .722 283	8.4114 .026 185	9.3204 .209 267	8.4856 .031 349	9.9950 .989 15	9.9987 .997 7	----- --- ---	9.8010 .632 269	7.6684 .005 182	9.3364 .217 280	
λ <sub>2</sub> -----	9.6707 .468 113	9.9961 .991 13	9.8276 .672 275	8.1941 .016 357	9.3018 .200 259	9.7195 .524 105	9.7627 .579 98	9.8010 .632 91	----- --- ---	9.3301 .214 272	8.7786 .060 190	
μ <sub>2</sub> -----	8.7223 .053 21	9.3018 .200 101	8.9141 .016 3	9.8276 .672 85	9.9961 .991 347	8.5420 .035 13	8.1935 .016 6	7.6684 .005 178	9.3301 .214 88	----- --- ---	9.7627 .579 98	
ν <sub>2</sub> -----	9.2966 .198 103	8.1941 .016 3	9.8276 .672 85	9.9961 .991 347	9.6823 .481 249	9.3168 .207 95	9.3301 .214 88	9.3364 .217 80	8.7786 .060 170	9.7627 .579 262	9.1043 .127 91	
2SM-----	8.8476 .070 192	9.3301 .214 272	8.1935 .016 354	9.0793 .120 256	8.5765 .038 338	8.4114 .026 185	8.1941 .016 357	8.7291 .054 350	9.3018 .200 259	8.1926 .016 351	9.1043 .127 269	

Table 29.—*Elimination factors—Continued*  
SERIES 29 DAYS. DIURNAL CONSTITUENTS

Constituent sought (A)	Disturbing constituents (B, C, etc.)									
	J <sub>1</sub>	K <sub>1</sub>	M <sub>1</sub>	O <sub>1</sub>	OO	P <sub>1</sub>	Q <sub>1</sub>	2Q	S <sub>1</sub>	p <sub>1</sub>
J <sub>1</sub> -----	8.6955	8.6896	8.7199	8.8144	9.2092	8.6937	8.6672	9.0538	8.3224	
	.050	.049	.052	.065	.162	.049	.046	.113	.021	
	351	341	328	13	322	319	310	336	.344	
K <sub>1</sub> -----	8.6955	8.6955	8.7517	8.7517	9.9818	8.7199	8.6937	9.9954	8.0542	
	.050	.050	.056	.056	.959	.052	.049	.990	.011	
	9	351	338	22	331	328	319	346	.354	
M <sub>1</sub> -----	8.6896	8.6955	-----	8.8144	8.7199	9.0674	8.7517	8.7199	8.4418	7.9579
	.049	.050	-----	.065	.052	.117	.056	.052	.028	.009
	19	9	-----	347	32	161	338	328	175	.183
O <sub>1</sub> -----	8.7199	8.7517	8.8144	-----	8.7185	8.2616	8.6955	8.6896	8.3262	8.9810
	.052	.056	.065	-----	.052	.018	.050	.049	.021	.096
	32	22	13	-----	44	174	351	341	8	.196
OO-----	8.8144	8.7517	8.7199	8.7185	-----	9.0332	8.6848	8.6504	8.9334	8.4666
	.065	.056	.052	.052	-----	.108	.048	.045	.086	.029
	347	338	328	316	-----	309	306	297	324	.332
P <sub>1</sub> -----	9.2092	9.9818	9.0674	8.2616	9.0332	-----	7.7378	8.2260	9.9954	8.6248
	.162	.959	.117	.018	.108	-----	.005	.017	.990	.042
	38	29	199	186	51	-----	357	348	14	.202
Q <sub>1</sub> -----	8.6937	8.7199	8.7517	8.6955	8.6848	7.7378	-----	8.6955	8.4846	9.9857
	.049	.052	.056	.050	.048	.005	-----	.050	.031	.968
	41	32	22	9	54	3	-----	351	17	.25
2Q-----	8.6672	8.6937	8.7199	8.6896	8.6504	8.2260	8.6955	-----	8.5377	9.1825
	.046	.049	.052	.049	.045	.017	.050	-----	.034	.152
	50	41	32	19	63	12	9	-----	27	.35
S <sub>1</sub> -----	9.0538	9.9954	8.4418	8.3262	8.9334	9.9954	8.4846	8.5377	-----	8.1807
	.113	.990	.028	.021	.086	.990	.031	.034	-----	.015
	24	14	185	352	36	346	343	333	-----	.188
p <sub>1</sub> -----	8.3224	8.0542	7.9579	8.9810	8.4666	8.6248	9.9857	9.1825	8.1807	-----
	.021	.011	.009	.096	.029	.042	.968	.152	.015	-----
	16	6	177	164	28	158	335	325	172	-----

Table 29.—*Elimination factors—Continued*  
SERIES 29 DAYS. SEMIDIURNAL CONSTITUENTS

Constituent sought (A)	Disturbing constituents (B, C, etc.)											
	K <sub>2</sub>	L <sub>2</sub>	M <sub>2</sub>	N <sub>2</sub>	2N	R <sub>2</sub>	S <sub>2</sub>	T <sub>2</sub>	λ <sub>2</sub>	μ <sub>2</sub>	ν <sub>2</sub>	2SM
K <sub>2</sub> -----	8.8144	8.7517	8.7199	8.6937	9.9954	9.9818	9.9587	9.2092	8.3224	8.0542	9.0054	
	.065	.056	.052	.049	.990	.959	.909	.162	.021	.011	.101	
	347	338	328	319	346	331	317	322	344	354	145	
L <sub>2</sub> -----	8.8144	-----	8.6955	8.6896	8.6798	7.9581	8.9810	9.2842	9.9857	7.7378	8.2616	8.6248
	.065	-----	.050	.049	.048	.009	.096	.192	.968	.005	.018	.042
	13	-----	351	341	332	178	164	150	335	357	186	158
M <sub>2</sub> -----	8.7517	8.6955	-----	8.6955	8.6896	8.3262	8.2616	8.7772	8.9810	8.2616	8.9810	8.2588
	.056	.050	-----	.050	.049	.021	.018	.060	.096	.018	.096	.018
	22	9	-----	351	341	8	174	159	164	186	196	167
N <sub>2</sub> -----	8.7199	8.6896	8.6955	-----	8.6955	8.4846	7.7378	8.3278	8.2616	8.9810	9.9857	7.5900
	.052	.049	.050	-----	.050	.031	.005	.021	.018	.096	.968	.004
	32	19	9	-----	351	17	3	169	174	196	25	177
2N-----	8.6937	8.6798	8.6896	8.6955	-----	8.5377	8.2260	7.4179	7.7378	9.9857	9.1825	7.7379
	.049	.048	.049	.050	-----	.034	.017	.003	.005	.968	.152	.005
	41	28	19	9	-----	27	12	178	3	25	35	6
R <sub>2</sub> -----	9.9954	7.9581	8.3262	8.4846	8.5377	-----	9.9954	9.9818	9.0538	7.2754	8.1807	8.7772
	.990	.009	.021	.031	.034	-----	.990	.959	.113	.002	.015	.060
	14	182	352	343	333	-----	346	331	336	359	188	159
S <sub>2</sub> -----	9.9818	8.9810	8.2616	7.7378	8.2260	9.9954	-----	9.9954	8.6955	8.2588	8.6248	8.2616
	.959	.096	.018	.005	.017	.990	-----	.990	.050	.018	.042	.018
	29	196	186	357	348	14	-----	346	351	193	202	174
T <sub>2</sub> -----	9.9587	9.2842	8.7772	8.3278	7.4179	9.9818	9.9954	-----	8.4418	8.5780	8.8324	8.3262
	.909	.192	.060	.021	.003	.959	.990	-----	.028	.038	.068	.021
	43	210	201	191	182	29	14	-----	185	207	217	8
λ <sub>2</sub> -----	9.2092	9.9857	8.9810	8.2616	7.7378	9.0538	8.6955	8.4418	-----	8.6248	8.9640	7.7378
	.162	.968	.096	.018	.005	.113	.050	.028	-----	.042	.092	.005
	38	25	196	186	357	24	9	175	-----	202	212	3
μ <sub>2</sub> -----	8.3224	7.7378	8.2616	8.9810	9.9857	7.2754	8.2588	8.5780	8.6248	-----	8.6955	8.2539
	.021	.005	.018	.096	.968	.002	.018	.038	.042	-----	.050	.018
	16	3	174	164	335	335	1	167	153	158	9	161
ν <sub>2</sub> -----	8.0542	8.2616	8.9810	9.9857	9.1825	8.1807	8.6248	8.8324	8.9640	8.6955	-----	8.5015
	.011	.018	.096	.968	.152	.015	.042	.068	.092	.050	-----	.032
	6	174	164	335	325	172	158	143	148	351	-----	151
2SM-----	9.0054	8.6248	8.2588	7.5900	7.7379	8.7772	8.2616	8.3262	7.7378	8.2539	8.5015	-----
	.101	.042	.018	.004	.005	.060	.018	.021	.005	.018	.032	-----
	215	202	193	183	354	201	186	352	357	199	209	-----

Table 29.—*Elimination factors—Continued*  
SERIES 58 DAYS. DIURNAL CONSTITUENTS

Constituent sought (A)	Disturbing constituents (B, C, etc.)										$p_1$
	J <sub>1</sub>	K <sub>1</sub>	M <sub>1</sub>	O <sub>1</sub>	OO	P <sub>1</sub>	Q <sub>1</sub>	2Q	S <sub>1</sub>		
J <sub>1</sub> -----	8.6896	8.6657	8.6504	8.8039	9.1056	8.5715	8.4713	9.0154	8.3059		
	.049	.046	.045	.064	.128	.037	.030	.104	.020		
	341	322	297	25	284	278	259	313	329		
K <sub>1</sub> -----	8.6896	8.6896	8.7185	8.7185	9.9254	8.6504	8.5715	9.9818	8.0520		
	.049	.049	.052	.052	.842	.045	.037	.959	.011		
	19	341	316	44	303	297	.278	331	348		
M <sub>1</sub> -----	8.6657	8.6896	-----	8.8039	8.6504	9.0427	8.7185	8.6504	8.4403	7.9572	
	.046	.049		.064	.045	.110	.052	.045	.028	.009	
	38	19		335	63	142	316	.297	170	186	
O <sub>1</sub> -----	8.6504	8.7185	8.8039	-----	8.5737	8.2588	8.6896	8.6657	8.3224	8.9640	
	.045	.052	.064		.037	.018	.049	.046	.021	.092	
	63	44	25		88	167	341	322	16	212	
OO-----	8.8039	8.7185	8.6504	8.5737	-----	8.8349	8.4575	8.3057	8.8391	8.4112	
	.064	.052	.045	.037		.068	.029	.020	.069	.026	
	335	316	297	272		259	253	.234	287	303	
P <sub>1</sub> -----	9.1056	9.9254	9.0427	8.2588	8.8349	-----	7.7379	8.2155	9.9818	8.5907	
	.128	.842	.110	.018	.068		.005	.016	.959	.039	
	76	57	218	193	101		354	.335	29	225	
Q <sub>1</sub> -----	8.5715	8.6504	8.7185	8.6896	8.4575	7.7379	-----	8.6896	8.4645	9.9418	
	.037	.045	.052	.049	.029	.005		.049	.029	.875	
	82	63	44	19	107	6		341	35	51	
2Q-----	8.4713	8.5715	8.6504	8.6657	8.3057	8.2155	8.6896	-----	8.4887	9.0969	
	.030	.037	.045	.046	.020	.016	.049		.031	.125	
	101	82	63	38	126	25	19		.53	70	
S <sub>1</sub> -----	9.0154	9.9818	8.4403	8.3224	8.8391	9.9818	8.4645	8.4887	-----	8.1761	
	.104	.959	.028	.021	.069	.959	.029	.031	-----	.015	
	47	29	190	344	73	331	.325	307	---	196	
P <sub>1</sub> -----	8.3059	8.0520	7.9572	8.9640	8.4112	8.5907	9.9418	9.0969	8.1761	-----	
	.020	.011	.009	.092	.026	.039	.875	.125	.015	---	
	34	12	174	148	57	135	309	290	.164	---	

Table 29.—*Elimination factors—Continued*  
SERIES 58 DAYS. SEMIDIURNAL CONSTITUENTS

Constituent sought (A)	Disturbing constituents (B, C, etc.)											
	K <sub>2</sub>	L <sub>2</sub>	M <sub>2</sub>	N <sub>2</sub>	2N	R <sub>2</sub>	S <sub>2</sub>	T <sub>2</sub>	λ <sub>2</sub>	μ <sub>2</sub>	ν <sub>2</sub>	2SM
K <sub>2</sub> -----	8.8039	8.7185	8.6504	8.5715	9.9818	9.9254	9.8237	9.1056	8.3059	8.0520	8.9185	
	.064	.052	.045	.037	.959	.842	.666	.128	.020	.011	.083	
	335	316	297	278	331	303	274	284	329	348	110	
L <sub>2</sub> -----	8.8039	8.6896	8.6657	8.6244	7.9579	8.9640	9.2209	9.9418	7.7379	8.2588	8.5907	
	.064	.049	.046	.042	.009	.092	.166	.875	.005	.018	.039	
	25	---	341	322	303	177	148	120	306	354	193	135
M <sub>2</sub> -----	8.7185	8.6896	-----	8.6896	8.6657	8.3224	8.2588	8.7480	8.9640	8.2588	8.9640	8.2475
	.052	.049	----	.049	.046	.021	.018	.056	.092	.018	.092	.018
	44	19	---	341	322	16	167	138	148	193	212	154
N <sub>2</sub> -----	8.6504	8.6657	8.6896	-----	8.6896	8.4645	7.7379	8.3193	8.2588	8.9640	9.9418	7.5898
	.045	.046	.049	----	.049	.029	.005	.021	.018	.092	.875	.004
	63	38	19	---	341	35	6	157	167	212	51	173
2N-----	8.5715	8.6244	8.6657	8.6896	-----	8.4887	8.2155	7.4165	7.7379	9.9418	9.0969	7.7356
	.037	.042	.046	.049	----	.031	.016	.003	.005	.875	.125	.005
	82	57	38	19	---	53	25	176	6	51	70	12
R <sub>2</sub> -----	9.9818	7.9579	8.3224	8.4645	8.4887	-----	9.9818	9.9254	9.0154	7.2736	8.1761	8.7480
	.959	.009	.021	.029	.031	----	.959	.842	.104	.002	.015	.056
	29	183	344	325	307	---	331	303	313	357	196	138
S <sub>2</sub> -----	9.9254	8.9640	8.2588	7.7379	8.2155	9.9818	-----	9.9818	8.6896	8.2475	8.5907	8.2588
	.842	.092	.018	.005	.016	.959	----	.959	.049	.018	.039	.018
	57	212	193	354	3-5	29	---	331	341	206	225	167
T <sub>2</sub> -----	9.8237	9.2209	8.7480	8.3193	7.4165	9.9254	9.9818	-----	8.4402	8.5270	8.7366	8.3224
	.666	.166	.056	.021	.003	.842	.959	----	.028	.034	.055	.021
	86	240	222	203	184	57	29	---	190	234	253	16
λ <sub>2</sub> -----	9.1056	9.9418	8.9640	8.2588	7.7379	9.0154	8.6896	8.4402	-----	8.5907	8.8933	7.7379
	.128	.875	.092	.018	.005	.104	.049	.028	----	.039	.078	.005
	76	51	212	193	354	47	19	170	----	225	244	6
μ <sub>2</sub> -----	8.3050	7.7379	8.2588	8.9640	9.9418	7.2736	8.2475	8.5270	8.5907	-----	8.6896	8.2286
	.020	.005	.018	.092	.875	.002	.018	.034	.039	----	.049	.017
	31	6	167	148	309	3	154	126	135	----	19	141
ν <sub>2</sub> -----	8.0520	8.2588	8.9640	9.9418	9.0969	8.1761	8.5907	8.7366	8.8933	8.6896	-----	8.4439
	.011	.018	.092	.875	.125	.015	.039	.055	.078	.049	----	.028
	12	167	148	309	290	164	135	107	116	341	----	122
2SM-----	8.9185	8.5907	8.2475	7.5898	7.7356	8.7480	8.2588	8.3224	7.7379	8.2286	8.4439	-----
	.083	.039	.018	.004	.005	.056	.018	.021	.005	.017	.028	-----
	250	225	206	187	348	222	193	344	354	219	238	---

Table 29.—*Elimination factors—Continued*  
SERIES 87 DAYS. DIURNAL CONSTITUENTS

Constituent sought (A)	Disturbing constituents (B, C, etc.)									
	J <sub>1</sub>	K <sub>1</sub>	M <sub>1</sub>	O <sub>1</sub>	O <sub>O</sub>	P <sub>1</sub>	Q <sub>1</sub>	2Q	S <sub>1</sub>	$\rho_1$
J <sub>1</sub> -----	8.6798	8.6244	8.5225	8.7857	8.9030	8.3232	7.9841	8.9481	8.2780	
	.048	.042	.033	.061	.080	.021	.010	.089	.019	
	332	303	265	38	246	237	209	289	313	
K <sub>1</sub> -----	8.6798	-----	8.6798	8.6607	8.6607	9.8237	8.5225	8.3232	9.9587	8.0476
	.048	-----	.048	.046	.046	.666	.033	.021	.909	.011
	28	-----	332	294	66	274	265	237	317	341
M <sub>1</sub> -----	8.6244	8.6798	-----	8.7857	8.5225	9.0002	8.6607	8.5225	8.4376	7.9556
	.042	.048	-----	.061	.033	.100	.046	.033	.027	.009
	57	28	-----	322	95	123	294	265	165	190
O <sub>1</sub> -----	8.5225	8.6607	8.7857	-----	8.2641	8.2539	8.6798	8.6244	8.3155	8.9351
	.033	.046	.061	-----	.018	.018	.048	.042	.021	.086
	95	66	38	-----	133	161	332	303	23	228
O <sub>O</sub> -----	8.7857	8.6607	8.5225	8.2641	-----	8.3377	7.8138	7.4337	8.6579	8.3116
	.061	.046	.033	.018	-----	.022	.007	.003	.045	.020
	322	294	265	227	-----	208	199	351	251	275
P <sub>1</sub> -----	8.9030	9.8237	9.0002	8.2539	8.3377	-----	7.7367	8.1982	9.9587	8.5315
	.080	.666	.100	.018	.022	-----	.005	.016	.909	.034
	114	86	237	199	152	-----	351	323	43	247
Q <sub>1</sub> -----	8.3232	8.5225	8.6607	8.6798	7.8138	7.7367	-----	8.6798	8.4303	9.8640
	.021	.033	.046	.048	.007	.005	-----	.048	.027	.731
	123	95	66	28	161	9	-----	332	52	76
2Q-----	7.9841	8.3232	8.5225	8.6244	7.4337	8.1982	8.6798	-----	8.4014	8.9351
	.010	.021	.033	.042	.003	.016	.048	-----	.025	.086
	151	123	95	57	9	37	28	-----	80	104
S <sub>1</sub> -----	8.9481	9.9587	8.4376	8.3155	8.6579	9.9587	8.4303	8.4014	-----	8.1689
	.089	.909	.003	.021	.045	.909	.027	.025	-----	.015
	71	43	195	337	109	317	308	280	-----	204
P <sub>1</sub> -----	8.2780	8.0476	7.9556	8.9351	8.3116	8.5315	9.8640	8.9351	8.1689	-----
	.019	.011	.009	.086	.020	.034	.731	.086	.015	-----
	47	19	170	132	85	113	284	256	156	-----

Table 29.—*Elimination factors—Continued*  
 SERIES 87 DAYS. SEMIDIURNAL CONSTITUENTS

Constituent sought (A)	Disturbing constituents (B, C, etc.)											
	K <sub>2</sub>	L <sub>2</sub>	M <sub>2</sub>	N <sub>2</sub>	2N	R <sub>2</sub>	S <sub>2</sub>	T <sub>2</sub>	λ <sub>2</sub>	μ <sub>2</sub>	ν <sub>2</sub>	2SM
K <sub>2</sub>	8.7857	8.6607	8.5225	8.3232	9.9587	9.8237	9.5416	8.9030	8.2780	8.0476	8.7538	
	.061	.046	.033	.021	.909	.666	.348	.080	.019	.011	.057	
	322	294	265	237	317	274	231	246	313	341	75	
L <sub>2</sub>	8.7857	8.6798	8.6244	8.5247	7.9576	8.9351	9.1055	9.8640	7.7367	8.2539	8.5315	
	.061	.048	.042	.033	.009	.086	.127	.731	.005	.018	.034	
	38	332	303	275	175	132	89	284	351	199	113	
M <sub>2</sub>	8.6607	8.6798	8.6798	8.6244	8.3155	8.2539	8.6976	8.9351	8.2539	8.9351	8.2286	
	.046	.048	.048	.042	.021	.018	.050	.086	.018	.086	.017	
	66	28	332	303	23	161	118	132	199	228	141	
N <sub>2</sub>	8.5225	8.6244	8.6798	8.6798	8.4303	7.7367	8.3068	8.2539	8.9351	9.8640	7.5883	
	.033	.042	.048	.048	.027	.005	.020	.018	.086	.731	.004	
	95	57	28	332	52	9	146	161	228	76	170	
2N	8.3232	8.5247	8.6244	8.6798	8.4014	8.1982	7.4165	7.7367	9.8640	8.9351	7.7314	
	.021	.033	.042	.048	.025	.016	.003	.005	.731	.086	.005	
	123	85	57	28	80	37	174	9	76	104	18	
R <sub>2</sub>	9.9587	7.9576	8.3155	8.4303	8.4014	9.9587	9.8237	8.9481	7.2740	8.1689	8.6976	
	.909	.009	.021	.027	.025	.909	.666	.089	.002	.015	.050	
	43	185	337	308	280	317	274	289	356	204	118	
S <sub>2</sub>	9.8237	8.9351	8.2539	7.7367	8.1982	9.9587	9.9587	8.6798	8.2286	8.5315	8.2539	
	.666	.086	.018	.005	.016	.909	.009	.048	.017	.034	.018	
	86	228	199	351	323	43	317	332	219	247	161	
T <sub>2</sub>	9.5416	9.1055	8.6976	8.3068	7.4165	9.8237	9.9587	8.4376	8.4358	8.5521	8.3155	
	.348	.127	.050	.020	.003	.666	.909	.027	.027	.036	.021	
	129	271	242	214	186	86	43	195	262	290	23	
λ <sub>2</sub>	8.9030	9.8640	8.9351	8.2539	7.7367	8.9481	8.6798	8.4376	8.5315	8.7629	7.7367	
	.080	.731	.086	.018	.005	.080	.048	.027	—	.034	.058	.005
	114	76	228	199	351	71	28	165	—	247	275	9
μ <sub>2</sub>	8.2780	7.7367	8.2539	8.9351	9.8640	7.2740	8.2286	8.4358	8.5315	8.6798	8.1849	
	.019	.005	.018	.086	.731	.002	.017	.027	.034	—	.048	.015
	47	9	161	132	284	4	141	98	113	—	28	122
ν <sub>2</sub>	8.0476	8.2539	8.9351	9.8640	8.9351	8.1689	8.5315	8.5521	8.7629	8.6798	8.3401	
	.011	.018	.086	.731	.086	.015	.034	.036	.058	.048	—	.022
	19	161	132	284	256	156	113	70	85	332	—	93
2SM	8.7538	8.5315	8.2286	7.5883	7.7314	8.6976	8.2539	8.3155	7.7367	8.1849	8.3401	
	.057	.034	.017	.004	.005	.050	.018	.021	.005	.015	.022	
	285	247	219	190	342	242	199	337	351	238	267	—

Table 29.—*Elimination factors—Continued*  
SERIES 105 DAYS. DIURNAL CONSTITUENTS

Constituent sought (A)	Disturbing constituents (B, C, etc.)									
	J <sub>1</sub>	K <sub>1</sub>	M <sub>1</sub>	O <sub>1</sub>	OO	P <sub>1</sub>	Q <sub>1</sub>	2Q	S <sub>1</sub>	$\rho_1$
J <sub>1</sub> -----	8.6704 .047 ---	8.5885 .039 214	8.4422 .028 248	8.4953 .031 271	8.8322 .068 158	8.2332 .017 291	7.7808 .006 308	8.3722 .024 339	8.1065 .013 342	----- .216
K <sub>1</sub> -----	8.6704 .047 146	8.6704 .047 214	8.5381 .035 236	8.5381 .035 124	9.7311 .538 257	8.4422 .028 271	8.2332 .017 305	9.9393 .870 308	7.0766 .001 182	-----
M <sub>1</sub> -----	8.5885 .039 112	8.6704 .047 146	----- .031 202	8.4422 .028 89	8.8219 .066 42	8.5381 .035 236	8.4422 .028 271	8.9548 .090 94	8.3679 .023 328	-----
O <sub>1</sub> -----	8.4422 .028 89	8.5381 .035 124	8.4953 .031 158	----- .019 ---	8.2803 .015 67	8.1856 .015 20	8.6704 .047 214	8.5885 .039 248	8.6113 .041 72	8.8929 .078 306
OO-----	8.4953 .031 202	8.5381 .035 236	8.4422 .028 271	8.2803 .019 293	----- .015 ---	8.4500 .028 313	7.9556 .009 327	6.4362 .000 181	7.5174 .003 185	8.1640 .015 239
P <sub>1</sub> -----	8.8322 .068 69	9.7311 .538 103	8.8219 .066 318	8.1856 .015 340	8.4500 .028 47	----- .007 ---	7.8500 .007 194	8.2067 .016 228	9.9393 .870 52	8.4685 .029 286
Q <sub>1</sub> -----	8.2332 .017 55	8.4422 .028 89	8.5381 .035 124	8.6704 .047 146	7.9556 .009 33	7.8500 .007 166	----- .007 ---	8.6704 .047 214	8.2396 .017 38	9.7951 .624 92
2Q-----	7.7808 .006 21	8.2332 .017 55	8.4422 .028 89	8.5885 .039 112	6.4362 .000 179	8.2067 .016 132	8.6704 .047 146	----- .007 ---	7.1241 .001 4	8.7943 .062 58
S <sub>1</sub> -----	8.3722 .024 18	9.9393 .870 52	8.9548 .090 266	8.6113 .041 288	7.5174 .003 175	9.9393 .870 308	8.2396 .017 322	7.1241 .001 356	----- .024 ---	8.3820 .024 234
$\rho_1$ -----	8.1065 .013 144	7.0766 .001 178	8.3679 .023 32	8.8929 .078 54	8.1640 .015 121	8.4685 .029 74	9.7951 .624 268	8.7943 .062 302	8.3820 .024 126	----- ---

Table 29.—*Elimination factors—Continued*  
 SERIES 105 DAYS. SEMIDIURNAL CONSTITUENTS

Constituent sought (A)	Disturbing constituents (B, C, etc.)											
	K <sub>2</sub>	L <sub>2</sub>	M <sub>2</sub>	N <sub>2</sub>	2N	R <sub>2</sub>	S <sub>2</sub>	T <sub>2</sub>	λ <sub>2</sub>	μ <sub>2</sub>	ν <sub>2</sub>	2SM
K <sub>2</sub> .....	8.4953 .031 202	8.5381 .035 236	8.4422 .028 271	8.2332 .017 305	9.9392 .869 308	9.7311 .538 257	9.1892 .155 205	8.8322 .068 291	8.1065 .013 216	7.0766 .001 182	8.6847 .048 97	
L <sub>2</sub> .....	8.4953 .031 158	8.6704 .047 214	8.5885 .039 248	8.4347 .027 282	8.9311 .085 106	8.8929 .078 54	7.6403 .004 2	9.7951 .624 268	7.8500 .007 194	8.1856 .015 340	8.4685 .015 74	
M <sub>2</sub> .....	8.5381 .035 124	8.6704 .047 146	8.6704 .047 214	8.5885 .039 248	8.6113 .041 72	8.1856 .015 20	8.3896 .025 148	8.8929 .078 54	8.1856 .015 340	8.8929 .078 306	8.1585 .014 40	
N <sub>2</sub> .....	8.4422 .028 89	8.5885 .039 112	8.6704 .047 146	8.6704 .047 214	8.2395 .017 38	7.8500 .007 166	8.4362 .027 114	8.1856 .015 20	8.8929 .078 306	9.7951 .624 92	7.2638 .002 6	
2N.....	8.2332 .017 55	8.4347 .027 78	8.5885 .039 112	8.6704 .047 146	7.1241 .001 4	8.2067 .016 132	8.3366 .022 80	7.8500 .007 166	9.7951 .624 92	8.7043 .062 58	7.8368 .007 152	
R <sub>2</sub> .....	9.0392 .869 52	8.9311 .085 254	8.6113 .041 288	8.2395 .017 322	7.1241 .001 356	9.9392 .016 308	9.7311 .025 257	8.3722 .014 342	8.3410 .022 268	8.3896 .024 234	8.6609 .024 148	
S <sub>2</sub> .....	9.7311 .538 103	8.8929 .078 306	8.1856 .015 340	7.8500 .007 194	8.2067 .016 228	9.9392 .069 52	8.3722 .047 308	9.9392 .047 214	8.6704 .014 320	8.1585 .029 286	8.1856 .015 20	
T <sub>2</sub> .....	9.1892 .155 155	7.6403 .004 358	8.3896 .025 212	8.4362 .027 246	8.3366 .022 280	9.7311 .538 103	9.9392 .869 52	8.9548 .069 ---	7.6654 .005 266	8.0785 .012 192	8.6113 .041 338	
λ <sub>2</sub> .....	8.8322 .068 69	9.7951 .624 92	8.8929 .078 306	8.1856 .015 340	7.8500 .007 194	8.3722 .024 18	8.6704 .047 146	8.9548 .090 94	8.4685 .029 ---	8.6609 .029 286	7.8500 .007 252	
μ <sub>2</sub> .....	8.1065 .013 144	7.8500 .007 166	8.1856 .015 20	8.8929 .078 54	9.7951 .624 268	8.3410 .022 92	8.1585 .014 40	7.6654 .005 168	8.4685 .029 74	8.6704 .047 ---	8.1117 .013 60	
ν <sub>2</sub> .....	7.0766 .001 178	8.1856 .015 20	8.8929 .078 54	8.1856 .624 268	8.7943 .062 302	8.3820 .024 126	8.4685 .029 74	8.0785 .012 22	8.6609 .046 108	8.6704 .047 214	8.2581 .018 94	
2SM.....	8.6847 .048 263	8.4685 .029 286	8.1585 .014 320	7.2638 .002 354	7.8368 .007 208	8.3896 .025 212	8.1856 .015 340	8.6113 .041 288	7.8500 .007 194	8.1117 .013 300	8.2581 .018 266	

Table 29.—*Elimination factors—Continued*  
 SERIES 134 DAYS. DIURNAL CONSTITUENTS

Constituent sought (A)	Disturbing constituents (B, C, etc.)									
	J <sub>1</sub>	K <sub>1</sub>	M <sub>1</sub>	O <sub>1</sub>	OO	P <sub>1</sub>	Q <sub>1</sub>	2Q	S <sub>1</sub>	p <sub>1</sub>
J <sub>1</sub> -----	8.4360	8.3946	8.2695	8.0361	8.7345	8.2094	8.0930	8.6047	7.7771	
	.027	.025	.019	.011	.054	.016	.012	.040	.006	
	205	229	239	170	253	264	288	319	201	
K <sub>1</sub> -----	8.4360	8.4360	8.2628	8.2628	9.5078	8.2695	8.2094	9.8992	7.1819	
	.027	.027	.018	.018	.322	.019	.016	.793	.002	
	155	205	214	146	228	239	264	294	356	
M <sub>1</sub> -----	8.3946	8.4360	--	8.0361	8.2695	8.4838	8.2628	8.2695	8.8500	8.2196
	.025	.027	-	.011	.019	.030	.018	.019	.071	.017
	131	155	-	190	121	23	214	239	.89	332
O <sub>1</sub> -----	8.2695	8.2628	8.0361	-----	8.1796	7.9151	8.4360	8.3946	8.5206	8.6697
	.019	.018	.011	-----	.015	.008	.027	.025	.033	.047
	121	146	170	---	111	14	205	229	80	322
OO-----	8.0361	8.2628	8.2695	8.1796	-----	8.4760	8.1133	7.9812	8.2156	7.8315
	.011	.018	.019	.015	-----	.030	.013	.010	.016	.007
	190	214	239	249	---	262	273	298	328	211
P <sub>1</sub> -----	8.7345	9.5078	8.4838	7.9151	8.4760	-----	7.6424	7.9951	9.8992	8.2746
	.054	.322	.030	.008	.030	-----	.004	.010	.793	.019
	107	132	337	346	98	---	191	216	.66	308
Q <sub>1</sub> -----	8.2094	8.2695	8.2628	8.4360	8.1133	7.6424	-----	8.4360	8.2605	9.6387
	.016	.019	.018	.027	.013	.004	-----	.027	.018	.435
	96	121	146	155	87	169	---	205	55	117
2Q-----	8.0930	8.2094	8.2695	8.3946	7.9812	7.9951	8.4360	-----	7.9233	8.7610
	.012	.016	.019	.025	.010	.010	.027	-----	.008	.058
	72	96	121	131	62	144	155	---	30	92
S <sub>1</sub> -----	8.6047	9.8992	8.8500	8.5206	8.2156	9.8992	8.2605	7.9233	-----	8.3143
	.040	.793	.071	.033	.016	.793	.018	.008	-----	.021
	41	66	271	280	32	294	305	330	-----	242
p <sub>1</sub> -----	7.7771	7.1819	8.2196	8.6697	7.8315	8.2746	9.6387	8.7610	8.3143	-----
	.006	.002	.017	.047	.007	.019	.435	.058	.021	-----
	159	4	28	38	149	52	243	268	118	---

Table 29.—*Elimination factors—Continued*  
SERIES 134 DAYS. SEMIDIURNAL CONSTITUENTS

Constituent sought (A)	Disturbing constituents (B, C, etc.)											
	K <sub>2</sub>	L <sub>2</sub>	M <sub>2</sub>	N <sub>2</sub>	2N	R <sub>2</sub>	S <sub>2</sub>	T <sub>2</sub>	λ <sub>2</sub>	μ <sub>2</sub>	ν <sub>2</sub>	2SM
K <sub>2</sub> -----	8.0361	8.2628	8.2695	8.2094	9.8992	9.5078	8.9538	8.7345	7.7771	7.1819	8.5254	
----- .011	.018	.019	.016	.793	.322	.090	.054	.006	.002	.034		
---- 190	214	239	264	294	228	342	253	201	356	61		
L <sub>2</sub> -----	8.0361	-----	8.4360	8.3946	8.3215	8.8285	8.6697	8.5871	9.6387	7.6424	7.9151	8.2746
----- .011	-----	.027	.025	.021	.067	.047	.039	.435	.004	.008	.019	
---- 170	205	229	254	104	38	152	243	191	346	346	346	52
M <sub>1</sub> -----	8.2628	8.4360	-----	8.4360	8.3946	8.5206	7.9151	8.4622	8.6697	7.9151	8.6697	7.9208
----- .018	.027	-----	.027	.025	.033	.008	.029	.047	.008	.047	.008	
---- 146	155	-----	205	229	80	14	128	38	346	322	322	27
N <sub>2</sub> -----	8.2698	8.3946	8.4360	-----	8.4360	8.2605	7.6424	8.3592	7.9151	8.6697	9.6387	6.7753
----- .019	.025	.027	-----	.027	.018	.004	.023	.008	.047	.435	.001	
---- 121	131	155	-----	205	55	169	103	14	322	117	117	2
2N-----	8.2094	8.3215	8.3946	8.4360	-----	7.9233	7.9951	8.2280	7.6424	9.6387	8.7610	7.6344
----- .016	.021	.025	.027	-----	.008	.010	.017	.004	.435	.058	.004	
---- 96	106	131	155	-----	30	144	78	169	117	92	92	158
R <sub>2</sub> -----	9.8992	8.8285	8.5206	8.2605	7.9233	-----	9.8992	9.5079	8.6047	8.2346	8.3143	8.4622
----- .793	.067	.033	.018	.008	-----	.793	.322	.040	.017	.021	.029	
---- 66	256	280	305	330	-----	294	228	319	267	242	242	128
S <sub>1</sub> -----	9.5078	8.6697	7.9151	7.6424	7.9951	9.8992	-----	9.8992	8.4360	7.9028	8.2746	7.9151
----- .322	.047	.008	.004	.010	.793	-----	.793	.027	.008	.019	.008	
---- 132	322	346	191	216	66	-----	294	205	333	308	308	14
T <sub>2</sub> -----	8.9538	8.5871	8.4622	8.3592	8.2280	9.5079	9.8992	-----	8.8500	8.0509	7.7834	8.5206
----- .090	.039	.029	.023	.017	.322	.793	-----	.071	.011	.006	.033	
---- 18	208	232	257	282	132	66	-----	271	219	194	194	80
λ <sub>2</sub> -----	8.7345	9.6387	8.6697	7.9151	7.6424	8.6047	8.4360	8.8500	-----	8.2746	8.5650	7.6424
----- .054	.435	.047	.008	.004	.040	.027	.071	-----	.019	.037	.004	
---- 107	117	322	346	191	41	155	89	-----	308	284	284	169
μ <sub>2</sub> -----	7.7771	7.6424	7.9151	8.6697	9.6387	8.2346	7.9028	8.0509	8.2746	-----	8.4360	7.8820
----- .006	.004	.008	.047	.435	.017	.008	.011	.019	-----	.027	.008	
---- 159	169	14	38	243	93	27	141	52	-----	155	155	41
ν <sub>2</sub> -----	7.1819	7.9151	8.6697	9.6387	8.7610	8.3143	8.2746	7.7834	8.5650	8.4360	-----	8.1118
----- .002	.008	.047	.435	.058	.021	.019	.006	.037	.027	-----	.013	
---- 4	14	38	243	268	118	52	166	76	205	205	205	65
2SM-----	8.5254	8.2746	7.9028	6.7753	7.6344	8.4622	7.9151	8.5206	7.6424	7.8820	8.1118	-----
----- .034	.019	.008	.001	.004	.029	.008	.033	.004	.008	.013	-----	
---- 299	308	333	358	202	232	346	280	191	319	295	295	---

Table 29.—*Elimination factors—Continued*  
SERIES 163 DAYS. DIURNAL CONSTITUENTS

Constituent sought (A)	Disturbing constituents (B, C, etc.)									
	J <sub>1</sub>	K <sub>1</sub>	M <sub>1</sub>	O <sub>1</sub>	O O	P <sub>1</sub>	Q <sub>1</sub>	2Q	S <sub>1</sub>	$\rho_1$
J <sub>1</sub> -----	8.1495	8.1341	7.9150	7.4365	8.4234	7.9579	7.9582	8.6570	7.0948	
	.014	.014	.008	.003	.027	.009	.009	.045	.001	
	195	210	207	3	215	223	238	295	185	
K <sub>1</sub> -----	8.1495	8.1495	7.7528	7.7528	9.0723	7.9150	7.9579	9.8470	7.5128	
	.014	.014	.006	.006	.118	.008	.009	.703	.003	
	165	195	192	168	199	207	223	280	350	
M <sub>1</sub> -----	8.1341	8.1495	-----	7.4365	7.9150	7.6604	7.7528	7.9150	8.7629	8.0859
	.014	.014		.003	.008	.005	.006	.008	.058	.012
	150	165		357	153	4	192	207	84	335
O <sub>1</sub> -----	7.9150	7.7528	7.4365	-----	7.7427	7.5513	8.1495	8.1341	8.4422	8.3724
	.008	.006	.003		.006	.004	.014	.014	.028	.024
	153	168	3		156	7	195	210	87	338
O O-----	7.4365	7.7528	7.9150	7.7427	-----	8.1140	7.8343	7.8631	8.3776	6.6248
	.003	.006	.008	.006		.013	.007	.007	.024	.000
	357	192	207	204		212	220	235	292	182
P <sub>1</sub> -----	8.4234	9.0723	7.6604	7.5513	8.1140	-----	7.4230	7.7409	9.8470	7.9852
	.027	.118	.005	.004	.013		.003	.006	.703	.010
	145	161	356	353	148		188	203	80	331
Q <sub>1</sub> -----	7.9579	7.9150	7.7528	8.1495	7.8343	7.4230	-----	8.1495	8.2410	9.3888
	.009	.008	.006	.014	.007	.003		.014	.017	.245
	137	153	168	165	140	172		195	72	142
2Q-----	7.9582	7.9579	7.9150	8.1341	7.8631	7.7409	8.1495	-----	8.0589	8.5769
	.009	.009	.008	.014	.007	.006	.014		.011	.038
	122	137	153	150	125	157	165		57	127
S <sub>1</sub> -----	8.6570	9.8470	8.7629	8.4422	8.3776	9.8470	8.2410	8.0589	-----	8.2562
	.045	.703	.058	.028	.024	.703	.017	.011		.018
	65	80	276	273	68	280	288	57		250
$\rho_1$ -----	7.0948	7.5128	8.0859	8.3724	6.6248	7.9852	9.3888	8.5769	8.2562	-----
	.001	.003	.012	.024	.000	.010	.245	.038	.018	
	175	10	25	22	178	29	218	233	110	---

Table 29.—*Elimination factors—Continued*  
 SERIES 163 DAYS. SEMIDIURNAL CONSTITUENTS

Constituent sought ( <i>A</i> )	Disturbing constituents ( <i>B</i> , <i>C</i> , etc.)											
	K <sub>2</sub>	L <sub>2</sub>	M <sub>2</sub>	N <sub>2</sub>	2N	R <sub>2</sub>	S <sub>2</sub>	T <sub>2</sub>	λ <sub>2</sub>	μ <sub>2</sub>	ν <sub>2</sub>	2SM
K <sub>2</sub>	7.4365 .003 357	7.7528 .006 192	7.9150 .008 207	7.9579 .009 223	9.8470 .703 280	9.0723 .118 199	9.3179 .208 299	8.4234 .027 215	7.0948 .001 185	7.5128 .003 350	8.1450 .014 26	
L <sub>2</sub>	7.4365 .003 3	8.1495 .014 195	8.1341 .014 210	8.1078 .013 226	8.7464 .056 103	8.3724 .024 22	8.7614 .058 122	9.3888 .245 218	7.4230 .003 188	7.5513 .004 353	7.9852 .010 29	
M <sub>2</sub>	7.7528 .006 168	8.1495 .014 165	8.1341 .014 195	8.1495 .014 210	8.4422 .028 87	7.5513 .004 7	8.4590 .029 107	8.3724 .024 22	7.5513 .004 353	8.3724 .024 338	7.5483 .004 14	
N <sub>2</sub>	7.9150 .008 153	8.1341 .014 150	8.1495 .014 165	8.1495 .014 195	8.2410 .017 72	7.4230 .003 172	8.2850 .019 92	7.5513 .004 7	8.3724 .024 338	9.3888 .245 142	6.3062 .000 179	
2N	7.9579 .009 137	8.1078 .013 134	8.1341 .014 150	8.1495 .014 165	----- 8.0588 .011	8.7409 .006 57	8.1397 .014 157	7.4230 .003 76	9.3888 .245 172	8.5769 .038 142	7.4186 .003 164	
R <sub>2</sub>	9.8470 .703 80	8.7464 .056 257	8.4422 .028 273	8.2410 .017 288	8.0588 .011 303	----- 9.8470 .011	9.0725 .118 280	8.6570 .045 199	8.1488 .045 295	8.2563 .018 265	8.4590 .029 250	
S <sub>2</sub>	9.0723 .118 161	8.3724 .024 338	7.5513 .004 353	7.4230 .003 188	7.7409 .006 203	9.8470 .703 80	----- 9.8470 .014	8.1495 .014 280	7.5483 .004 195	7.9852 .010 346	7.5513 .010 331	
T <sub>2</sub>	9.3179 .208 61	8.7614 .058 238	8.4590 .029 253	8.2850 .019 268	8.1397 .014 284	9.0725 .118 161	9.8470 .708 80	----- 8.7629 .058	8.1668 .015 276	8.1966 .016 246	8.4422 .028 231	
λ <sub>2</sub>	8.4234 .027 145	9.3888 .245 142	8.3724 .024 338	7.5513 .004 353	7.4230 .003 188	8.6570 .045 65	8.1495 .014 165	8.7629 .058 84	----- 7.9852 .010	8.3386 .022 331	7.4230 .003 315	
μ <sub>2</sub>	7.0948 .001 175	7.4230 .003 172	7.5513 .004 7	8.3724 .024 22	9.3888 .245 218	8.1488 .014 95	7.5483 .004 14	8.1668 .015 114	7.9852 .010 29	----- 8.1495 .014	7.5426 .003 165	
ν <sub>2</sub>	7.5128 .003 10	7.5513 .004 7	8.3724 .024 22	9.3888 .245 218	8.5769 .038 233	8.2563 .018 110	7.9852 .010 29	8.1966 .016 129	8.3386 .022 45	8.1495 .014 195	7.8424 .007 36	
2SM	8.1450 .014 334	7.9852 .010 331	7.5483 .004 346	6.3062 .000 181	7.4186 .003 196	8.4590 .029 253	7.5513 .004 353	8.4422 .028 273	7.4230 .003 188	7.5426 .003 339	7.8424 .007 324	

Table 29.—*Elimination factors—Continued*  
SERIES 182 DAYS. DIURNAL CONSTITUENTS

Constituent sought (A)	Disturbing constituents (B, C, etc.)									
	J <sub>1</sub>	K <sub>1</sub>	M <sub>1</sub>	O <sub>1</sub>	OO	P <sub>1</sub>	Q <sub>1</sub>	2Q	S <sub>1</sub>	$\rho_1$
J <sub>1</sub> -----	7.6613	7.6591	7.0355	8.0828	7.3819	6.5151	7.0698	8.6281	7.3308	
	.005	.005	.001	.012	.002	.000	.001	.42	.002	
	186	192	356	16	357	182	187	271	350	
K <sub>1</sub> -----	7.6613	7.6613	7.5891	7.5891	8.6868	7.0355	6.5151	9.7807	7.6468	
	.005	.005	.004	.004	.049	.001	.000	.604	.004	
	174	186	350	10	351	356	182	265	344	
M <sub>1</sub> -----	7.6591	7.6613	-----	8.0828	7.0355	8.1441	7.5891	7.0355	8.6868	7.9586
	.005	.005	-----	.012	.001	.014	.004	.001	.049	.009
	168	174	-----	344	4	165	350	356	80	338
O <sub>1</sub> -----	7.0355	7.5891	8.0828	-----	7.5826	6.4230	7.6613	7.6591	8.3698	7.7679
	.001	.004	.012	-----	.004	.000	.005	.005	.023	.006
	4	10	16	-----	20	1	186	192	95	354
OO-----	8.0828	7.5891	7.0355	7.5826	-----	7.8388	7.3409	7.0344	8.3250	7.6132
	.012	.004	.001	.004	-----	.007	.002	.001	.021	.004
	344	350	356	340	-----	341	346	352	256	334
P <sub>1</sub> -----	7.3819	8.6868	8.1441	6.4230	7.8388	-----	7.1547	7.3491	9.7807	7.3097
	.002	.049	.014	.000	.007	-----	.001	.002	.604	.002
	3	9	195	359	19	-----	185	191	95	353
Q <sub>1</sub> -----	6.5151	7.0355	7.5891	7.6613	7.3409	7.1547	-----	7.6613	8.1911	8.8560
	.000	.001	.004	.005	.002	.001	-----	.005	.016	.072
	178	4	10	174	14	175	-----	186	89	168
2Q-----	7.0698	6.5151	7.0355	7.6591	7.0344	7.3491	7.6613	-----	8.0615	8.0931
	.001	.000	.001	.005	.001	.002	.005	-----	.012	.012
	173	178	4	168	8	169	174	-----	84	162
S <sub>1</sub> -----	8.6281	9.7807	8.6868	8.3698	8.3250	9.7807	8.1911	8.0615	-----	8.2024
	.042	.604	.049	.023	.021	.004	.016	.012	-----	.016
	89	95	280	265	104	265	271	276	-----	258
$\rho_1$ -----	7.3308	7.6468	7.9586	7.7679	7.6132	7.3097	8.8560	8.0931	8.2024	-----
	.002	.004	.009	.000	.004	.002	.072	.012	.016	-----
	10	16	22	6	26	7	192	198	102	-----

Table 29.—*Elimination factors—Continued*  
SERIES 192 DAYS. SEMIDIURNAL CONSTITUENTS

Constituent sought ( <i>A</i> )	Disturbing constituents ( <i>B</i> , <i>C</i> , etc.)											
	K <sub>2</sub>	L <sub>2</sub>	M <sub>2</sub>	N <sub>2</sub>	2N	R <sub>2</sub>	S <sub>2</sub>	T <sub>2</sub>	λ <sub>2</sub>	μ <sub>2</sub>	ν <sub>2</sub>	2SM
<b>K<sub>2</sub></b>	8.0828	7.5891	7.0355	6.5151	9.7807	8.6868	9.2922	7.3819	7.3308	7.6468	7.6012	
	.012	.004	.001	.000	.604	.049	.196	.002	.002	.004	.004	
	344	350	356	182	265	351	256	357	350	344	171	
<b>L<sub>2</sub></b>	8.0828	7.6613	7.6591	7.6554	8.6778	7.7679	8.7615	8.8560	7.1547	6.4230	7.3097	
	.012	.005	.005	.005	.048	.006	.058	.072	.001	.000	.002	
	16	186	192	197	101	6	92	192	185	359	7	
<b>M<sub>2</sub></b>	7.5891	7.6613	7.6613	7.6591	8.3698	6.4230	8.4057	7.7679	6.4230	7.7679	6.4265	
	.004	.005	-----	.005	.023	.000	.025	.006	.000	.006	.000	
	10	174	-----	186	192	95	1	86	6	359	354	1
<b>N<sub>2</sub></b>	7.0355	7.6591	7.6613	7.6613	8.1911	7.1547	8.2077	6.4230	7.7679	8.8560	6.8803	
	.001	.005	.005	-----	.005	.016	.001	.016	.000	.006	.072	.001
	4	168	174	-----	186	89	175	80	1	354	168	175
<b>2N</b>	6.5151	7.6554	7.6591	7.6613	-----	8.0615	7.3491	8.0649	7.1547	8.8560	8.0931	7.1525
	.000	.005	.005	.005	-----	.012	.002	.012	.001	.072	.012	.001
	178	163	168	174	-----	84	169	74	175	168	162	170
<b>R<sub>2</sub></b>	9.7807	8.6778	8.3698	8.1911	8.0615	-----	9.7807	8.6863	8.6281	8.0768	8.2024	8.4057
	.604	.048	.023	.016	.012	-----	.604	.049	.042	.012	.016	.025
	95	259	265	271	276	-----	265	351	271	264	258	86
<b>S<sub>2</sub></b>	8.6868	7.7679	6.4230	7.1547	7.3491	9.7807	-----	9.7807	7.6613	6.4265	7.3097	6.4230
	.049	.006	.000	.001	.002	.604	-----	.604	.005	.000	.002	.000
	9	354	359	185	191	95	-----	265	186	359	353	1
<b>T<sub>2</sub></b>	9.2922	8.7615	8.4057	8.2077	8.0649	8.6863	9.7807	-----	8.6866	8.0959	8.2350	8.3698
	.196	.058	.025	.016	.012	.049	.604	-----	.049	.012	.017	.023
	104	268	274	280	286	9	95	-----	280	273	268	95
<b>λ<sub>2</sub></b>	7.3819	8.8560	7.7679	6.4230	7.1547	8.6281	7.6613	8.6866	-----	7.3097	7.7656	7.1547
	.002	.072	.006	.000	.001	.042	.005	.049	-----	.002	.006	.001
	3	168	354	359	185	89	174	80	-----	353	347	175
<b>μ<sub>2</sub></b>	7.3308	7.1547	6.4230	7.7679	8.8560	8.0768	8.4265	8.0859	7.3097	-----	7.6613	6.4253
	.002	.001	.000	.006	.006	.072	.012	.000	.012	.002	.005	.000
	10	175	1	6	192	96	1	87	7	174	2	
<b>ν<sub>2</sub></b>	7.6468	6.4230	7.7679	8.8560	8.0931	8.2024	7.3097	8.2350	7.7656	7.6613	-----	7.1202
	.004	.000	.006	.072	.012	.016	.002	.017	.006	.005	-----	.001
	16	1	6	192	198	102	7	92	13	186	8	
<b>2SM</b>	7.6012	7.3097	6.4265	6.8803	7.1525	8.4057	6.4230	8.3698	7.1547	6.4253	7.1202	
	.004	.002	.000	.001	.001	.025	.000	.023	.001	.000	.001	
	189	353	359	185	190	274	359	265	185	358	352	

Table 29.—*Elimination factors—Continued*  
SERIES 221 DAYS. DIURNAL CONSTITUENTS

Constituent sought (A)	Disturbing constituents (B, C, etc.)									
	J <sub>1</sub>	K <sub>1</sub>	M <sub>1</sub>	O <sub>1</sub>	OO	P <sub>1</sub>	Q <sub>1</sub>	2Q	S <sub>1</sub>	$\rho_1$
J <sub>1</sub> -----	7.4061	7.4052	7.8848	8.2672	8.3590	7.7969	7.7322	8.5324	7.6535	
	.003	.003	.008	.019	.023	.006	.005	.034	.005	
	356	353	324	28	318	321	317	247	334	
K <sub>1</sub> -----	7.4061	7.4061	8.0179	8.0179	9.2077	7.8848	7.7969	9.6969	7.7209	
	.003	.003	.010	.010	.161	.008	.006	.498	.005	
	4	356	328	32	322	324	321	251	338	
M <sub>1</sub> -----	7.4052	7.4061	-----	8.2672	7.8848	8.4189	8.0179	7.8848	8.6172	7.8313
	.003	.003	-----	.019	.008	.026	.010	.008	.041	.007
	7	4	-----	332	36	146	328	324	75	341
O <sub>1</sub> -----	7.8848	8.0179	8.2672	-----	7.9465	7.3352	7.4061	7.4052	8.2991	7.8800
	.008	.010	.019	-----	.009	.002	.003	.003	.020	.008
	36	32	28	-----	64	174	356	353	103	190
OO-----	8.2672	8.0179	7.8848	7.9465	-----	8.2351	7.8628	7.7946	8.0778	7.8188
	.019	.010	.008	.009	-----	.017	.007	.006	.012	.007
	332	328	324	296	-----	290	292	289	219	306
P <sub>1</sub> -----	8.3590	9.2077	8.4189	7.3352	8.2351	-----	6.7176	6.4350	9.6969	7.5854
	.023	.161	.026	.002	.017	-----	.001	.000	.498	.004
	42	38	214	186	70	-----	182	358	109	195
Q <sub>1</sub> -----	7.7969	7.8848	8.0179	7.4061	7.8628	6.7176	-----	7.4061	8.1112	8.8310
	.006	.008	.010	.003	.007	.001	-----	.003	.013	.068
	39	36	32	4	68	178	-----	356	107	13
2Q-----	7.7322	7.7969	7.8848	7.4052	7.7946	6.4350	7.4061	-----	7.9748	8.0073
	.005	.006	.008	.003	.006	.000	.003	-----	.009	.010
	43	39	36	7	71	2	4	-----	110	17
S <sub>1</sub> -----	8.5324	9.6969	8.6172	8.2991	8.0778	9.6969	8.1112	7.9748	-----	8.1495
	.034	.498	.041	.020	.012	.498	.013	.009	-----	.014
	113	109	285	257	141	251	253	250	-----	266
$\rho_1$ -----	7.6535	7.7209	7.8313	7.8800	7.8188	7.5854	8.8310	8.0073	8.1495	-----
	.005	.005	.007	.008	.007	.004	.068	.010	.014	-----
	26	22	19	170	54	165	347	343	94	-----

Table 29.—*Elimination factors—Continued*  
 SERIES 221 DAYS. SEMIDIURNAL CONSTITUENTS

Constituent sought (A)	Disturbing constituents (B, C, etc.)											
	K <sub>2</sub>	L <sub>2</sub>	M <sub>2</sub>	N <sub>2</sub>	2N	R <sub>2</sub>	S <sub>2</sub>	T <sub>2</sub>	λ <sub>2</sub>	μ <sub>2</sub>	ν <sub>2</sub>	2SM
K <sub>2</sub>	8.2872	8.0179	7.8848	7.7969	9.6969	9.2077	8.9831	8.3590	7.6535	7.7209	8.2035	
	.019	.010	.008	.006	.498	.161	.096	.023	.005	.005	.016	
	332	328	324	321	251	322	213	318	334	338	136	
L <sub>2</sub>	8.2672	-----	7.4061	7.4052	7.4037	8.6189	7.8800	8.6448	8.8310	6.7176	7.3352	7.5854
	.019	-----	.003	.003	.003	.042	.008	.044	.068	.001	.002	.004
	28	-----	356	353	349	99	170	62	347	182	186	165
M <sub>2</sub>	8.0179	7.4061	-----	7.4061	7.4052	8.2991	7.3352	8.3038	7.8800	7.3352	7.8800	7.3333
	.010	.003	-----	.003	.003	.020	.002	.020	.008	.002	.008	.002
	32	4	-----	356	353	103	174	65	170	186	190	168
N <sub>2</sub>	7.8848	7.4052	7.4061	-----	7.4061	8.1112	6.7176	8.1229	7.3352	7.8800	8.8310	7.0677
	.008	.003	.003	-----	.003	.013	.001	.013	.002	.008	.068	.001
	36	7	4	-----	356	107	178	69	174	190	13	172
2N	7.7969	7.4037	7.4052	7.4061	-----	7.9748	6.4350	7.9996	6.7176	8.8310	8.0073	6.7183
	.006	.003	.003	.003	-----	.009	.000	.010	.001	.068	.010	.001
	39	11	7	4	-----	110	2	73	178	13	17	176
R <sub>2</sub>	9.6969	8.6189	8.2991	8.1112	7.9748	-----	9.6970	9.2076	8.5324	8.0145	8.1495	8.3038
	.498	.042	.020	.013	.009	-----	.498	.161	.034	.010	.014	.020
	109	261	257	253	250	-----	251	322	247	263	.266	65
S <sub>2</sub>	9.2077	7.8800	7.3352	6.7176	6.4350	9.6970	-----	9.6970	7.4061	7.3333	7.5854	7.3352
	.161	.008	.002	.001	.000	.498	-----	.498	.003	.002	.004	.002
	38	190	186	182	358	109	-----	251	356	192	195	174
T <sub>2</sub>	8.9831	8.6448	8.3038	8.1229	7.9996	9.2076	9.6970	-----	8.6172	7.9704	8.0914	8.2991
	.096	.044	.020	.013	.010	.161	.498	-----	.041	.009	.012	.020
	147	298	295	291	287	38	109	-----	285	301	304	103
λ <sub>2</sub>	8.3590	8.8310	7.8800	7.3352	6.7176	8.5324	7.4061	8.6172	-----	7.5854	7.8738	6.7176
	.023	.068	.008	.002	.001	.034	.003	.041	-----	.004	.007	.001
	42	13	190	186	182	113	4	75	-----	195	199	178
μ <sub>2</sub>	7.6535	6.7176	7.3352	7.8800	8.8310	8.0145	7.3333	7.9704	7.5854	-----	7.4061	7.3294
	.005	.001	.002	.008	.068	.010	.002	.009	.004	-----	.003	.002
	26	178	174	170	347	97	168	59	165	-----	4	162
ν <sub>2</sub>	7.7209	7.3352	7.8800	8.8310	8.0073	8.1495	7.5854	8.0914	7.8738	7.4061	-----	7.4945
	.005	.002	.008	.068	.010	.014	.004	.012	.007	.003	-----	.003
	22	174	170	347	343	94	165	56	161	356	-----	159
2SM	8.2035	7.5854	7.3333	7.0677	6.7183	8.3038	7.3352	8.2991	6.7176	7.3294	7.4945	-----
	.016	.004	.002	.001	.001	.020	.002	.020	.001	.002	.003	-----
	224	195	192	188	184	295	186	257	182	198	201	-----

Table 29.—*Elimination factors—Continued*  
SERIES 250 DAYS. DIURNAL CONSTITUENTS

Constituent sought (A)	Disturbing constituents (B, C, etc.)									
	J <sub>1</sub>	K <sub>1</sub>	M <sub>1</sub>	O <sub>1</sub>	OO	P <sub>1</sub>	Q <sub>1</sub>	2Q	S <sub>1</sub>	$\rho_1$
J <sub>1</sub> -----	7.9011	7.8898	8.0302	8.3544	8.4768	7.9350	7.8438	8.3527	7.7796	
	.008	.008	.011	.023	.030	.009	.007	.023	.006	
	347	334	293	41	280	280	267	224	318	
K <sub>1</sub> -----	7.9011	7.9011	8.1489	8.1489	9.3286	8.0302	7.9350	9.5900	7.7661	
	.008	.008	.014	.014	.213	.011	.009	.389	.006	
	13	347	306	54	294	293	280	237	331	
M <sub>1</sub> -----	7.8898	7.9011	-----	8.3544	8.0302	8.5201	8.1489	8.0302	8.5519	7.6982
	.008	.008	-----	.023	.011	.033	.014	.011	.036	.005
	26	13	-----	319	67	127	306	293	70	344
O <sub>1</sub> -----	8.0302	8.1489	8.3544	-----	7.9171	7.6029	7.9011	7.8898	8.2274	8.2405
	.011	.014	.023	-----	.008	.004	.008	.008	.017	.017
	67	54	41	-----	108	168	347	384	111	205
OO-----	8.3544	8.1489	8.0302	7.9171	-----	8.1443	7.7748	7.6181	6.8959	7.8514
	.023	.014	.011	.008	-----	.014	.006	.004	.001	.007
	319	306	293	252	-----	239	239	226	183	277
P <sub>1</sub> -----	8.4768	9.3286	8.5201	7.6029	8.1443	-----	6.2382	7.3397	9.5900	7.8955
	.030	.213	.033	.004	.014	-----	.000	.002	.389	.008
	80	66	233	192	121	-----	359	346	123	218
Q <sub>1</sub> -----	7.9350	8.0302	8.1489	7.9011	7.7748	6.2382	-----	7.9011	7.9950	9.2133
	.009	.011	.014	.008	.006	.000	-----	.008	.010	.163
	80	67	54	13	121	1	-----	347	124	39
2Q-----	7.8438	7.9350	8.0302	7.8898	7.6181	7.3397	7.9011	-----	7.7821	8.3852
	.007	.009	.011	.008	.004	.002	.008	-----	.006	.024
	93	80	67	26	134	14	13	-----	137	52
S <sub>1</sub> -----	8.3527	9.5900	8.5519	8.2274	6.8959	9.5900	7.9950	7.7821	-----	8.0954
	.023	.389	.036	.017	.001	.389	.010	.006	-----	.012
	136	123	290	249	177	237	236	223	-----	275
$\rho_1$ -----	7.7796	7.7661	7.6982	8.2405	7.8514	7.8955	9.2133	8.3852	8.0954	-----
	.006	.006	.005	.017	.007	.008	.163	.024	.012	-----
	42	29	16	155	83	142	321	308	85	-----

Table 29.—*Elimination factors—Continued*  
SERIES 250 DAYS. SEMIDIURNAL CONSTITUENTS

Constituent sought (A)	Disturbing constituents (B, C, etc.)											
	K <sub>2</sub>	L <sub>2</sub>	M <sub>2</sub>	N <sub>2</sub>	2N	R <sub>2</sub>	S <sub>2</sub>	T <sub>2</sub>	λ <sub>2</sub>	μ <sub>2</sub>	ν <sub>2</sub>	2SM
K <sub>2</sub> -----	8.3544	8.1489	8.0302	7.9350	9.5900	9.3286	8.4129	8.4768	7.7796	7.7661	8.3023	
	.023	.014	.011	.009	.389	.213	.026	.030	.006	.006	.020	
	319	306	293	280	237	294	350	280	318	331	101	
L <sub>2</sub> -----	8.3544	-----	7.9011	7.8898	7.8703	8.5672	8.2405	8.3634	9.2133	6.2382	7.6029	7.8955
	.023	-----	.008	.008	.007	.037	.017	.023	.163	.000	.004	.008
	41	-----	347	334	321	98	155	31	321	359	192	143
M <sub>2</sub> -----	8.1489	7.9011	-----	7.9011	7.8898	8.2274	7.6029	8.1376	8.2405	7.6029	8.2405	7.5928
	.014	.008	-----	.008	.008	.017	.004	.014	.017	.004	.017	.004
	54	13	-----	347	334	111	168	44	155	192	205	155
N <sub>2</sub> -----	8.0302	7.8898	7.9011	-----	7.9011	7.9950	6.2382	8.0259	7.6029	8.2405	9.2133	7.1697
	.011	.008	.008	-----	.008	.010	.000	.011	.004	.017	.163	.001
	67	26	13	-----	347	124	1	58	168	205	39	168
2N-----	7.9350	7.8703	7.8898	7.9011	-----	7.7821	7.3397	7.9414	6.2382	9.2133	8.3852	6.2381
	.009	.007	.008	.008	-----	.006	.002	.009	.000	.163	.024	.000
	80	39	26	13	-----	137	14	71	1	39	52	2
R <sub>2</sub> -----	9.5900	8.5672	8.2274	7.9950	7.7821	-----	9.5901	9.3286	8.3528	7.9596	8.0954	8.1376
	.389	.037	.017	.010	.010	-----	.389	.213	.023	.009	.012	.014
	123	262	249	236	223	-----	237	294	224	261	275	44
S <sub>2</sub> -----	9.3286	8.2405	7.6029	6.2382	7.3397	9.5901	-----	9.5901	7.9011	7.5928	7.8955	7.6029
	.213	.017	.004	.000	.002	.389	-----	.389	.008	.004	.008	.004
	66	205	192	359	346	123	-----	237	347	205	218	168
T <sub>2</sub> -----	8.4129	8.3634	8.1376	8.0259	7.9414	9.3286	9.5901	-----	8.5519	7.7084	7.6345	8.2274
	.026	.023	.014	.011	.009	.213	.389	-----	.036	.005	.004	.017
	10	329	316	302	289	66	123	-----	290	328	341	111
λ <sub>2</sub> -----	8.4768	9.2133	8.2405	7.6029	6.2382	8.3528	7.9011	8.5519	-----	7.8955	8.1962	6.2382
	.030	.163	.017	.004	.000	.023	.008	.036	-----	.008	.016	.000
	80	39	205	192	359	136	13	70	-----	218	231	1
μ <sub>2</sub> -----	7.7796	6.2382	7.6029	8.2405	9.2133	7.9596	7.5928	7.7084	7.8955	-----	7.9011	7.5759
	.006	.000	.004	.017	.163	.009	.004	.005	.008	-----	.008	.004
	42	1	168	155	321	99	155	32	142	-----	13	143
ν <sub>2</sub> -----	7.7661	7.6029	8.2405	9.2133	8.3852	8.0954	7.8955	7.6345	8.1962	7.9011	-----	7.7671
	.006	.004	.017	.163	.024	.012	.008	.004	.016	.008	-----	.006
	29	168	155	321	308	85	142	19	129	347	-----	130
2SM-----	8.3023	7.8955	7.5928	7.1697	6.2381	8.1376	7.6029	8.2274	6.2382	7.5759	7.7671	-----
	.020	.008	.004	.001	.000	.014	.004	.017	.000	.004	.006	-----
	259	218	205	192	358	316	192	249	359	217	230	---

Table 29.—*Elimination factors—Continued*

## SERIES 279 DAYS. DIURNAL CONSTITUENTS

Constituent sought (A)	Disturbing constituents (B, C, etc.)									
	J <sub>1</sub>	K <sub>1</sub>	M <sub>1</sub>	O <sub>1</sub>	O <sub>O</sub>	P <sub>1</sub>	Q <sub>1</sub>	2Q	S <sub>1</sub>	$\rho_1$
J <sub>1</sub> -----	8.0816 .012 --- 337	8.0469 .011 --- 315	8.0127 .025 --- 261	8.3961 .024 --- 54	8.3841 .024 --- 242	7.8250 .007 --- 239	7.5672 .004 --- 216	7.9987 .010 --- 200	7.8339 .007 --- 303	
K <sub>1</sub> -----	8.0816 .012 23	8.0816 .012 --- 337	8.1800 .015 --- 284	8.1800 .015 --- 76	9.3172 .208 --- 265	8.0127 .010 --- 261	7.8250 .007 --- 239	9.4495 .282 --- 222	7.7947 .006 --- 325	
M <sub>1</sub> -----	8.0469 .011 45	8.0816 .012 23	----- --- 306	8.3961 .025 --- 99	8.5477 .010 --- 108	8.1800 .035 --- 284	8.0127 .015 --- 261	8.4890 .010 --- 65	7.5510 .031 --- 348	
O <sub>1</sub> -----	8.0127 .010 99	8.1800 .015 76	8.3961 .025 54	----- --- ---	7.5571 .004 --- 152	7.7343 .005 --- 161	8.0816 .012 --- 337	8.0469 .012 --- 315	8.1523 .014 --- 119	8.3798 .024 --- 221
O <sub>O</sub> -----	8.3961 .025 306	8.1800 .015 284	8.0127 .010 261	7.5571 .004 --- 208	----- --- ---	7.3456 .002 --- 189	6.7358 .001 --- 185	7.1964 .002 --- 342	7.9211 .008 --- 326	7.7771 .006 --- 249
P <sub>1</sub> -----	8.3841 .024 118	9.3172 .208 95	8.5477 .035 252	7.7343 .005 199	7.3456 .002 171	----- --- ---	6.8592 .001 --- 356	7.5574 .004 --- 334	9.4495 .282 --- 138	7.9990 .010 --- 240
Q <sub>1</sub> -----	7.8250 .007 121	8.0127 .010 99	8.1800 .015 76	8.0816 .012 23	6.7358 .001 175	6.8592 .001 4	----- --- ---	8.0816 .012 --- 337	7.8251 .007 --- 141	9.3242 .211 --- 64
2Q-----	7.5672 .004 144	7.8250 .007 121	8.0127 .010 99	8.0469 .011 45	7.1964 .002 18	7.5574 .004 26	8.0816 .012 23	----- --- ---	7.3460 .002 --- 164	8.4421 .028 --- 86
S <sub>1</sub> -----	7.9987 .010 160	9.4495 .282 138	8.4890 .031 295	8.1523 .014 241	7.9211 .008 34	9.4495 .282 222	7.8251 .007 219	7.3460 .002 196	----- --- ---	8.0384 .011 --- 283
P <sub>1</sub> -----	7.8339 .007 57	7.7947 .006 35	7.5510 .004 12	8.3798 .024 139	7.7771 .006 111	7.9990 .010 120	9.3242 .211 296	8.4421 .028 274	8.0384 .011 77	----- --- ---

Table 29.—*Elimination factors—Continued*  
SERIES 279 DAYS. SEMIDIURNAL CONSTITUENTS

Constituent sought ( <i>A</i> )	Disturbing constituents ( <i>B</i> , <i>C</i> , etc.)										
	K <sub>2</sub>	L <sub>2</sub>	M <sub>2</sub>	N <sub>2</sub>	2N	R <sub>2</sub>	S <sub>2</sub>	T <sub>2</sub>	λ <sub>2</sub>	μ <sub>2</sub>	ν <sub>2</sub>
K <sub>2</sub> .....	8.3961 .025 306	8.1800 .015 284	8.0127 .010 261	7.8250 .007 239	9.4494 .281 222	9.3172 .208 265	9.0421 .110 308	8.3841 .024 242	7.8330 .007 303	7.7947 .006 325	8.2246 .017 66
L <sub>2</sub> .....	8.3961 .025 54	8.0816 .012 337	8.0469 .011 315	7.9866 .010 292	8.5211 .038 96	8.3798 .024 139	6.9057 .001 1	9.3242 .211 296	6.8592 .001 356	7.7343 .005 199	7.9990 .010 120
M <sub>2</sub> .....	8.1800 .015 76	8.0816 .012 23	8.0469 .011 337	8.1523 .014 315	7.7433 .119 119	.005 161	.007 24	7.8491 .139	8.3798 .199	7.7343 .221	8.3798 .142
N <sub>2</sub> .....	8.0127 .010 99	8.0469 .011 45	8.0816 .012 23	----- .012	8.0816 .012 337	7.8251 .007 141	6.8592 .001 4	7.9108 .46	7.7343 .161	8.3798 .221	9.3242 .64
2N.....	7.8250 .007 121	7.9866 .010 68	8.0469 .011 45	8.0816 .012 23	----- .002	7.3463 .164	7.8721 .66	7.8885 .008 69	6.8592 .001 4	9.3242 .211 64	6.8588 .028 8
R <sub>2</sub> .....	9.4494 .281 138	8.5211 .033 264	8.1523 .014 241	7.8251 .007 219	7.3463 .002 196	----- .002	9.4496 .282 223	9.3172 .208 265	7.9987 .010 200	7.9102 .008 260	8.0384 .011 283
S <sub>2</sub> .....	9.3172 .208 95	8.3798 .024 221	7.7343 .005 199	6.8592 .001 356	7.8721 .007 294	9.4496 .282 137	----- .282 223	9.4496 .012 237	8.0816 .005 218	7.7105 .010 240	7.9990 .005 161
T <sub>2</sub> .....	9.0421 .110 52	6.9057 .001 359	7.8491 .007 336	7.9108 .008 314	7.8885 .008 291	9.3172 .208 95	9.4496 .282 137	----- .031 295	8.4891 .012 355	6.8703 .001 198	8.1523 .004 119
λ <sub>2</sub> .....	8.3841 .024 118	9.3242 .211 64	8.3798 .024 221	7.7343 .005 199	6.8592 .001 356	7.9987 .010 160	8.0384 .012 123	8.4891 .031 65	----- .010 240	7.9990 .018 263	8.2553 .001 4
μ <sub>2</sub> .....	7.8339 .007 57	6.8592 .001 4	7.7343 .005 161	8.3798 .024 139	9.3242 .211 296	7.9102 .008 100	7.7105 .005 142	6.8703 .001 5	7.9990 .010 120	----- .012 -----	8.0816 .005 23
ν <sub>2</sub> .....	7.7947 .006 35	7.7343 .005 161	8.3798 .024 139	9.3242 .211 296	8.4421 .028 274	8.0384 .011 77	7.9990 .010 120	7.5541 .004 162	8.2553 .018 97	8.0816 .012 337	7.8266 .007 101
2SM.....	8.2246 .017 294	7.9990 .010 240	7.7105 .005 218	7.2354 .002 195	6.8588 .001 352	7.8491 .007 336	7.7343 .005 199	8.1523 .014 241	6.8592 .001 356	7.6697 .005 236	7.8266 .007 259

Table 29.—*Elimination factors—Continued*

## SERIES 297 DAYS. DIURNAL CONSTITUENTS

Constituent sought (A)	Disturbing constituents (B, C, etc.)									
	J <sub>1</sub>	K <sub>1</sub>	M <sub>1</sub>	O <sub>1</sub>	OO	P <sub>1</sub>	Q <sub>1</sub>	2Q	S <sub>1</sub>	$\rho_1$
J <sub>1</sub> -----	8.2770	8.1622	7.9899	7.5338	8.3896	7.7726	7.1486	8.4204	7.5223	
	.019	.015	.010	.003	.025	.006	.001	.026	.003	
	220	260	266	173	287	306	346	253	206	
K <sub>1</sub> -----	8.2770	8.2770	8.0269	8.0269	9.2565	7.9899	7.7726	9.3360	7.3907	
	.019	.019	.011	.011	.181	.010	.006	.217	.002	
	140	220	227	133	247	266	306	214	346	
M <sub>1</sub> -----	8.1622	8.2770	-----	7.5338	7.9899	8.2044	8.0269	7.9899	7.5392	8.1019
	.015	.019	-----	.003	.010	.016	.011	.010	.003	.013
	100	140	-----	187	94	27	227	260	174	306
O <sub>1</sub> -----	7.9899	8.0269	7.5338	-----	7.8638	7.7467	8.2770	8.1622	7.5336	8.4724
	.010	.011	.003	-----	.007	.006	.019	.015	.003	.030
	94	133	173	-----	87	21	220	260	167	300
OO-----	7.5338	8.0269	7.9899	7.8638	-----	8.0954	7.6320	6.7805	8.1433	7.5129
	.003	.011	.010	.007	-----	.012	.004	.001	.014	.003
	187	227	266	273	294	313	353	260	213	
P <sub>1</sub> -----	8.3896	9.2565	8.2044	7.7467	8.0954	-----	7.5300	7.8163	9.3360	8.0286
	.025	.181	.016	.006	.012	-----	.003	.007	.217	.011
	73	113	333	339	66	-----	199	239	146	279
Q <sub>1</sub> -----	7.7726	7.9899	8.0269	8.2770	7.6320	7.5300	-----	8.2770	7.9031	9.3366
	.006	.010	.011	.019	.004	.003	-----	.019	.008	.217
	54	94	133	140	47	161	-----	220	127	80
2Q-----	7.1486	7.7726	7.9899	8.1622	6.7805	7.8163	8.2770	-----	7.8741	8.2220
	.001	.006	.010	.015	.001	.007	.019	-----	.007	.017
	14	54	94	100	7	121	140	-----	87	40
S <sub>1</sub> -----	8.4204	9.3360	7.5392	7.5336	8.1433	9.3360	7.9031	7.8741	-----	7.8897
	.026	.217	.003	.003	.014	.217	.008	.007	-----	.008
	107	146	186	193	100	214	233	273	-----	312
$\rho_1$ -----	7.5223	7.3907	8.1019	8.4724	7.5129	8.0286	9.3366	8.2220	7.8897	-----
	.003	.002	.013	.030	.003	.011	.217	.017	.008	-----
	154	14	54	60	147	81	280	320	48	-----

Table 29.—*Elimination factors—Continued*  
SERIES 297 DAYS. SEMIDIURNAL CONSTITUENTS

Constituent sought (A)	Disturbing constituents (B, C, etc.)										
	K <sub>2</sub>	L <sub>2</sub>	M <sub>2</sub>	N <sub>2</sub>	2N	R <sub>2</sub>	S <sub>2</sub>	T <sub>2</sub>	λ <sub>2</sub>	μ <sub>2</sub>	ν <sub>2</sub>
K <sub>2</sub> -----	7.5338	8.0269	7.9899	7.7726	9.3359	9.2565	9.1076	8.3896	7.5223	7.3907	8.2357
.003	.011	.010	.006	.217	.181	.128	.025	.003	.002	.017	
187	227	266	306	214	247	281	.287	206	346	88	
L <sub>2</sub> -----	7.5338	8.2770	8.1622	7.9326	8.1514	8.4724	8.5711	9.3366	7.5300	7.7467	8.0286
.003	.019	.015	.009	.014	.030	.037	.217	.003	.006	.011	
173	220	260	300	27	60	94	280	199	339	81	
M <sub>2</sub> -----	8.0269	8.2770	-----	8.1622	7.5339	7.7467	8.1268	8.4724	7.7467	8.4724	7.7179
.011	.019	-----	.019	.015	.003	.006	.013	.030	.006	.030	.005
133	140	-----	220	260	167	21	54	60	339	300	41
N <sub>2</sub> -----	7.9899	8.1622	8.2770	-----	8.2770	7.9031	7.5300	7.4214	7.7467	8.4724	9.3366
.010	.015	.019	-----	.019	.008	.003	.003	.006	.030	.217	.000
94	100	140	-----	220	127	161	14	21	300	80	1
2N-----	7.7726	7.9326	8.1622	8.2770	-----	7.8741	7.8163	7.5240	7.5300	9.3366	8.2220
.006	.009	.015	.019	-----	.007	.007	.003	.003	.217	.017	.003
54	60	100	140	-----	87	121	155	161	80	40	142
R <sub>2</sub> -----	9.3359	8.1514	7.5339	7.9031	7.8741	-----	9.3362	9.2566	8.4204	7.0150	7.8897
.217	.014	.003	.008	.007	-----	.217	.181	.026	.001	.008	.013
146	333	193	233	273	-----	214	247	253	352	312	54
S <sub>2</sub> -----	9.2565	8.4724	7.7467	7.5300	7.8163	9.3362	-----	9.3362	8.2770	7.7179	8.0286
.181	.030	.006	.003	.007	.217	-----	.217	.019	.005	.011	.006
113	300	339	199	239	146	-----	214	220	319	279	21
T <sub>2</sub> -----	9.1076	8.5711	8.1268	7.4214	7.5240	9.2566	9.3362	-----	7.5385	7.8920	8.0039
.128	.037	.013	.003	.003	.181	.217	-----	.003	.008	.010	.003
79	266	306	346	205	113	146	-----	186	285	245	167
λ <sub>2</sub> -----	8.3896	9.3366	8.4724	7.7467	7.5300	8.4204	8.2770	7.5385	-----	8.0286	8.1647
.025	.217	.030	.006	.006	.003	.026	.019	.003	-----	.011	.015
73	80	300	339	199	107	140	174	-----	279	239	161
μ <sub>2</sub> -----	7.5223	7.5300	7.7467	8.4724	9.3366	7.0150	7.7179	7.8920	8.0286	-----	8.2770
.003	.003	.006	.030	.217	.001	.005	.008	.011	-----	.019	.005
154	161	21	60	280	8	41	75	81	-----	140	62
ν <sub>2</sub> -----	7.3907	7.7467	8.4724	9.3366	8.2220	7.8897	8.0286	8.0039	8.1647	8.2770	7.7984
.002	.006	.030	.217	.017	.008	.011	.010	.015	.019	-----	.006
14	21	60	280	320	48	81	115	121	220	-----	102
2SM-----	8.2357	8.0286	7.7179	6.2014	7.5050	8.1268	7.7467	7.5339	7.5300	7.6680	7.7984
.017	.011	.005	.000	.003	.013	.006	.003	.003	.005	.006	-----
272	279	319	359	218	306	339	193	199	298	258	-----

Table 29.—*Elimination factors—Continued*  
SERIES 326 DAYS. DIURNAL CONSTITUENTS

Constituent sought: (A)	Disturbing constituents (B, C, etc.)									
	J <sub>1</sub>	K <sub>1</sub>	M <sub>1</sub>	O <sub>1</sub>	OO	P <sub>1</sub>	Q <sub>1</sub>	2Q	S <sub>1</sub>	$\rho_1$
J <sub>1</sub> -----	8.1340 .014 --- 210	8.0698 .012 --- 241	7.8631 .007 --- 235	7.4352 .003 --- 6	8.3392 .022 --- 249	7.8244 .007 --- 265	7.6841 .005 --- 296	8.2809 .019 --- 230	7.0934 .001 --- 190	
K <sub>1</sub> -----	8.1340 .014 150	8.1340 .014 --- 210	7.7427 .006 --- 204	7.7427 .006 --- 156	9.0470 .111 --- 219	7.8631 .007 --- 235	7.8244 .007 --- 265	9.0723 .118 --- 199	7.5061 .003 --- 340	
M <sub>1</sub> -----	8.0698 .012 119	8.1340 .014 150	----- --- 354	7.4352 .003 --- 125	7.8631 .007 --- 8	7.6587 .005 --- 204	7.7427 .006 --- 235	7.8631 .007 --- 169	7.7472 .006 --- 310	8.0423 .011 --- 310
O <sub>1</sub> -----	7.8631 .007 125	7.7427 .006 ---	7.4352 .003 6	----- --- 131	7.7018 .005 --- 14	7.5483 .004 --- 210	8.1340 .014 --- 241	8.0698 .012 --- 175	7.0956 .001 --- 315	8.3386 .022 --- 315
OO-----	7.4352 .003 354	7.7427 .006 204	7.8631 .007 235	7.7018 .005 229	----- --- 243	8.0443 .011 --- 259	7.7204 .005 --- 290	7.6227 .004 --- 224	7.9496 .009 --- 184	6.6245 .000 --- 184
P <sub>1</sub> -----	8.3392 .022 111	9.0470 .111 141	7.6587 .005 352	7.5483 .004 346	8.0443 .011 117	----- --- 243	7.4186 .003 --- 196	7.7040 .005 --- 227	9.0723 .118 --- 161	7.9254 .008 --- 301
Q <sub>1</sub> -----	7.8244 .007 95	7.8631 .007 125	7.7427 .006 156	8.1340 .014 150	7.7204 .005 101	7.4186 .003 164	----- --- 164	8.1340 .014 --- 210	7.7258 .005 --- 144	9.2882 .194 --- 105
2Q-----	7.6841 .005 64	7.8244 .007 95	7.8631 .007 125	8.0698 .012 119	7.6227 .004 70	7.7040 .005 133	8.1340 .014 150	----- --- 150	7.7948 .006 --- 114	8.3594 .023 --- 75
S <sub>1</sub> -----	8.2809 .019 130	9.0723 .118 161	7.7427 .006 191	7.0956 .001 185	7.9496 .009 136	9.0723 .118 199	7.7258 .005 216	7.7948 .006 246	----- --- 246	7.7844 .006 --- 321
$\rho_1$ -----	7.0934 .001 170	7.5061 .003 20	8.0423 .011 50	8.3386 .022 45	6.6245 .000 176	7.9254 .008 59	9.2882 .194 255	8.3594 .023 285	7.7844 .006 39	----- --- 39

Table 29.—*Elimination factors—Continued*  
SERIES 326 DAYS. SEMIDIURNAL CONSTITUENTS

Constituent sought ( <i>A</i> )	Disturbing constituents ( <i>B</i> , <i>C</i> , etc.)											
	K <sub>2</sub>	L <sub>2</sub>	M <sub>2</sub>	N <sub>2</sub>	2N	R <sub>2</sub>	S <sub>2</sub>	T <sub>2</sub>	λ <sub>2</sub>	μ <sub>2</sub>	ν <sub>2</sub>	2SM
K <sub>2</sub> .....	7.4352	7.7427	7.8631	7.8244	9.0720	9.0470	9.0037	8.3392	7.0934	7.5061	8.0971	
	.003	.006	.007	.007	.118	.111	.101	.022	.001	.003	.013	
	354	204	235	265	199	219	238	249	190	340	53	
L <sub>2</sub> .....	7.4352	8.1340	8.0698	7.9526	8.0858	8.3386	8.4852	9.2882	7.4186	7.5483	7.9254	
	.003	.014	.012	.009	.012	.022	.031	.194	.003	.004	.008	
	6	210	241	271	25	45	64	255	196	346	59	
M <sub>2</sub> .....	7.7427	8.1340	-----	8.1340	8.0698	7.0956	7.5483	7.9190	8.3386	7.5483	8.3386	7.5347
	.006	.014	-----	.014	.012	.001	.004	.008	.022	.004	.022	.003
	156	150	-----	210	241	175	14	34	45	346	315	28
N <sub>2</sub> .....	7.8631	8.0698	8.1340	-----	8.1340	7.7259	7.4186	6.7213	7.5483	8.3386	9.2882	6.3062
	.007	.012	.014	-----	.014	.005	.003	.001	.004	.022	.194	.000
	125	119	150	-----	210	144	164	3	14	315	105	178
2N.....	7.8244	7.9526	8.0698	8.1340	-----	7.7948	7.7040	7.5123	7.4186	9.2882	8.3594	7.4010
	.007	.009	.012	.014	-----	.006	.005	.003	.003	.194	.023	.003
	95	89	119	150	-----	114	133	153	164	105	75	148
R <sub>2</sub> .....	9.0720	8.0858	7.0956	7.7259	7.7948	-----	9.0725	9.0472	8.2809	7.0444	7.7843	7.9190
	.118	.012	.001	.005	.006	-----	.118	.112	.019	.001	.006	.008
	161	335	185	216	246	-----	199	219	230	351	321	34
S <sub>2</sub> .....	9.0470	8.3386	7.5483	7.4186	7.7040	9.0725	-----	9.0725	8.1340	7.5347	7.9254	7.5483
	.111	.022	.004	.003	.005	.118	-----	.118	.014	.003	.008	.004
	141	315	346	196	227	161	-----	199	210	332	301	14
T <sub>2</sub> .....	9.0037	8.4852	7.9190	6.7213	7.5123	9.0472	9.0725	-----	7.7468	7.7359	7.9960	7.0956
	.101	.031	.008	.001	.003	.112	.118	-----	.006	.005	.010	.001
	122	296	326	357	207	141	161	-----	191	312	282	175
λ <sub>2</sub> .....	8.3392	9.2882	8.3386	7.5483	7.4186	8.2809	8.1340	7.7568	-----	7.9254	8.1912	7.4186
	.022	.194	.022	.004	.003	.019	.014	.006	-----	.008	.016	.003
	111	105	315	346	196	130	150	169	-----	301	271	164
μ <sub>2</sub> .....	7.0934	7.4186	7.5483	8.3386	9.2882	7.0444	7.4347	7.7359	7.9254	-----	8.1340	7.5118
	.001	.003	.004	.022	.194	.001	.003	.005	.008	-----	.014	.003
	170	164	14	45	255	9	28	48	.59	-----	150	43
ν <sub>2</sub> .....	7.5061	7.5483	8.3386	9.2882	8.3594	7.7843	7.9254	7.9960	8.1912	8.1340	-----	7.7477
	.003	.004	.022	.194	.023	.006	.008	.010	.016	.014	-----	.006
	20	14	45	255	285	39	59	78	89	210	-----	73
2SM.....	8.0971	7.9254	7.5347	6.3062	7.4010	7.9190	7.5483	7.0956	7.4186	7.5118	7.7477	-----
	.013	.008	.003	.000	.003	.008	.004	.001	.003	.003	.006	-----
	307	301	332	182	212	326	346	185	196	317	287	-----

Table 29.—*Elimination factors—Continued*

## SERIES 355 DAYS. DIURNAL CONSTITUENTS

Constituent sought (A)	Disturbing constituents (B, C, etc.)									
	J <sub>1</sub>	K <sub>1</sub>	M <sub>1</sub>	O <sub>1</sub>	OO	P <sub>1</sub>	Q <sub>1</sub>	2Q	S <sub>1</sub>	$\rho_1$
J <sub>1</sub> -----	7.9464 .009 201	7.9167 .008 222	7.5111 .003 203	7.8888 .008 19	8.0444 .011 211	7.6331 .004 224	7.6506 .004 245	8.0032 .010 206	6.7724 .001 355	
K <sub>1</sub> -----	7.9464 .009 159	7.9464 .009 201	6.7064 .001 182	6.7064 .001 178	8.4581 .029 190	7.5111 .003 203	7.6331 .004 224	8.4598 .029 185	7.5794 .004 334	
M <sub>1</sub> -----	7.9167 .008 138	7.9464 .009 159	----- 341	7.8888 .008 157	7.5111 .003 175	7.7393 .005 8	6.7064 .001 201	7.5111 .003 222	7.8651 .007 164	7.9839 .010 313
O <sub>1</sub> -----	7.5111 .003 157	6.7064 .001 178	7.8888 .008 19	----- 341	6.7060 .001 175	7.2500 .002 8	7.9464 .009 201	7.9167 .008 222	6.7729 .001 3	8.1364 .014 331
OO-----	7.8888 .008 341	6.7064 .001 182	7.5111 .003 203	6.7060 .001 185	----- 341	7.3914 .002 192	7.3284 .002 206	7.4740 .003 227	7.1838 .002 187	7.3105 .002 336
P <sub>1</sub> -----	8.0444 .011 149	8.4581 .029 170	7.7393 .005 191	7.2500 .002 352	7.3914 .002 168	----- 341	7.2957 .002 193	7.5554 .004 214	8.4598 .029 175	7.7296 .005 324
Q <sub>1</sub> -----	7.6331 .004 136	7.5111 .003 157	6.7064 .001 178	7.9464 .009 159	7.3284 .002 154	7.2957 .002 167	----- 341	7.9464 .009 201	7.4212 .003 162	9.1482 .141 130
2Q-----	7.6506 .004 115	7.6331 .004 136	7.5111 .003 157	7.9167 .008 138	7.4740 .003 133	7.5554 .004 146	7.9464 .009 159	----- 341	7.5984 .004 141	8.3129 .021 109
S <sub>1</sub> -----	8.0032 .010 154	8.4598 .029 175	7.8651 .007 196	6.7729 .001 357	7.1838 .002 173	8.4598 .029 185	7.4212 .003 198	7.5984 .004 219	----- 329	7.6607 .005 329
$\rho_1$ -----	6.7724 .001 5	7.5794 .004 26	7.9839 .010 47	8.1364 .014 29	7.3105 .002 24	7.7296 .005 36	9.1482 .141 230	8.3129 .021 251	7.6607 .005 31	----- 31

Table 29.—*Elimination factors—Continued*  
SERIES 355 DAYS. SEMIDIURNAL CONSTITUENTS

Constituent sought ( <i>A</i> )	Disturbing constituents ( <i>B</i> , <i>C</i> , etc.)											
	K <sub>2</sub>	L <sub>2</sub>	M <sub>2</sub>	N <sub>2</sub>	2N	R <sub>2</sub>	S <sub>2</sub>	T <sub>2</sub>	λ <sub>2</sub>	μ <sub>2</sub>	ν <sub>2</sub>	2SM
K <sub>1</sub> -----	7.8888	6.7064	7.5111	7.6331	8.4589	8.4581	8.4553	8.0444	6.7724	7.5794	7.6440	
	.008	.001	.003	.004	.029	.029	.029	.011	.001	.004	.004	
	341	182	203	224	185	190	195	211	355	334	18	
L <sub>1</sub> -----	7.8888	7.9464	7.9167	7.8652	8.0217	8.1364	8.2393	9.1482	7.2957	7.2500	7.7296	
	.008	.009	.008	.007	.011	.014	.017	.141	.002	.002	.005	
	19	201	222	243	24	29	34	230	193	352	36	
M <sub>2</sub> -----	6.7064	7.9464	7.9464	7.9167	6.7712	7.2500	7.4842	8.1364	7.2500	8.1364	7.2458	
	.001	.009	.009	.008	.001	.002	.003	.014	.002	.014	.002	
	178	189	201	222	3	8	13	29	352	331	15	
N <sub>2</sub> -----	7.5111	7.9167	7.9464	7.9464	7.4212	7.2957	7.1008	7.2500	8.1364	9.1482	6.7017	
	.003	.008	.009	---	.009	.003	.002	.001	.002	.014	.141	.001
	157	138	159	201	162	167	172	8	331	130	174	
2N-----	7.6331	7.8652	7.9167	7.9464	7.5985	7.5554	7.5017	7.2957	9.1482	8.3129	7.2840	
	.004	.007	.008	.009	---	.004	.004	.003	.002	.141	.021	.002
	136	117	138	159	---	141	146	151	167	130	109	154
R <sub>2</sub> -----	8.4589	8.0217	6.7712	7.4212	7.5985	7.5554	8.4598	8.4586	8.0034	7.0678	7.6606	7.4842
	.029	.011	.001	.003	.004	---	.029	.029	.010	.001	.005	.003
	175	336	357	198	219	---	185	190	206	350	329	13
S <sub>2</sub> -----	8.4581	8.1364	7.2500	7.2957	7.5554	8.4598	8.4598	7.9464	7.2458	7.7296	7.2500	
	.029	.014	.002	.002	.004	.029	---	.029	.009	.002	.005	.002
	170	331	352	193	214	175	---	185	201	345	324	8
T <sub>2</sub> -----	8.4553	8.2393	7.4842	7.1008	7.5017	8.4586	8.4598	7.8648	7.3738	7.7893	6.7712	
	.029	.017	.003	.001	.003	.029	.029	---	.007	.002	.006	.001
	165	326	347	188	209	170	175	---	196	340	319	3
λ <sub>2</sub> -----	8.0444	9.1482	8.1364	7.2500	7.2957	8.0034	7.9464	7.8648	7.7296	8.0795	7.2957	
	.011	.141	.014	.002	.002	.010	.009	.007	---	.005	.012	.002
	149	130	331	352	193	154	159	164	---	324	303	167
μ <sub>2</sub> -----	6.7724	7.2957	7.2500	8.1364	9.1482	7.0678	7.2458	7.3738	7.7296	7.9464	7.2393	
	.001	.002	.002	.014	.141	.001	.002	.002	.005	---	.009	.002
	5	167	8	29	230	10	15	20	36	---	159	23
ν <sub>2</sub> -----	7.5794	7.2500	8.1364	9.1482	8.3129	7.6606	7.7296	7.7893	8.0795	7.9464	7.5728	
	.004	.002	.014	.141	.021	.005	.005	.006	.012	.009	---	.004
	26	8	29	230	251	31	36	41	57	201	---	44
2SM-----	7.6440	7.7296	7.2458	6.7017	7.2840	7.4842	7.2500	6.7712	7.2957	7.2393	7.5728	
	.004	.005	.002	.001	.002	.003	.002	.001	.002	.002	.004	---
	342	324	345	186	206	347	352	357	193	337	316	---

Table 29.—*Elimination factors—Continued*

SERIES 369 DAYS. DIURNAL CONSTITUENTS

Constituent sought (A)	Disturbing constituents (B, C, etc.)									
	J <sub>1</sub>	K <sub>1</sub>	M <sub>1</sub>	O <sub>1</sub>	OO	P <sub>1</sub>	Q <sub>1</sub>	2Q	S <sub>1</sub>	$\rho_1$
J <sub>1</sub> -----	8.3503	7.8740	7.8760	8.3371	8.2982	7.5509	7.4182	8.3235	5.7099	
	.022	.007	.008	.022	.020	.004	.003	.021	.000	
	290	219	287	112	286	217	326	288	0	
K <sub>1</sub> -----	8.3503	-----	8.3503	6.6332	6.6332	8.0072	7.8760	7.5509	8.0074	7.8892
	.022	-----	.022	.000	.000	.010	.008	.004	.010	.008
	70	-----	290	358	2	356	287	217	358	250
M <sub>1</sub> -----	7.8740	8.3503	-----	8.3371	7.8760	8.4104	6.6332	7.8760	8.3792	7.9045
	.007	.022	-----	.022	.008	.026	.000	.008	.024	.008
	141	70	-----	248	78	67	358	287	69	321
O <sub>1</sub> -----	7.8760	6.6332	8.3371	-----	6.6329	6.5537	8.3503	7.8740	5.7100	8.4169
	.008	.000	.022	-----	.000	.000	.022	.007	.000	.026
	73	2	112	-----	4	178	290	219	0	252
OO-----	8.3371	6.6332	7.8760	6.6329	-----	7.0438	7.6584	7.3523	6.8924	7.6527
	.022	.000	.008	.000	-----	.001	.005	.002	.001	.004
	248	358	287	356	-----	354	285	215	356	248
P <sub>1</sub> -----	8.2982	8.0072	8.4104	6.5537	7.0438	-----	7.8885	7.6024	8.0074	7.9217
	.020	.010	.026	.000	.001	-----	.008	.004	.010	.008
	74	4	293	182	6	-----	291	221	2	254
Q <sub>1</sub> -----	7.5509	7.8760	6.6332	8.3503	7.6584	7.8885	-----	8.3503	7.8824	9.0329
	.004	.008	.000	.022	.005	.008	-----	.022	.008	.108
	143	73	2	70	75	69	-----	290	71	143
2Q-----	7.4182	7.5509	7.8760	7.8740	7.3523	7.6024	8.3503	-----	7.5772	8.0586
	.003	.004	.008	.007	.002	.004	.022	-----	.004	.011
	34	143	73	141	145	139	70	-----	141	33
S <sub>1</sub> -----	8.3235	8.0074	8.3792	5.7100	6.8924	8.0074	7.8824	7.5772	-----	7.9055
	.021	.010	.024	.000	.001	.010	.008	.004	-----	.008
	72	2	291	0	4	358	289	219	-----	252
$\rho_1$ -----	5.7099	7.8892	7.9045	8.4169	7.6527	7.9217	9.0329	8.0586	7.9055	-----
	.000	.008	.008	.026	.004	.008	.108	.011	.008	-----
	0	110	39	108	112	106	217	327	108	-----

Table 29.—*Elimination factors—Continued*  
SERIES 369 DAYS. SEMIDIURNAL CONSTITUENTS

Constituent sought (A)	Disturbing constituents (B, C, etc.)											
	K <sub>2</sub>	L <sub>2</sub>	M <sub>2</sub>	N <sub>2</sub>	2N	R <sub>2</sub>	S <sub>2</sub>	T <sub>2</sub>	λ <sub>2</sub>	μ <sub>2</sub>	ν <sub>2</sub>	2SM
K <sub>2</sub> .....	8.3371	6.6332	7.8760	7.5509	8.0074	8.0072	8.0076	8.2982	5.7099	7.8892	7.1088	
.....	.022	.000	.008	.004	.010	.010	.010	.020	.000	.008	.001	
.....	248	358	287	217	358	356	354	286	0	250	175	
L <sub>2</sub> .....	8.3371	8.3503	7.8740	7.6166	8.3758	8.4169	8.4607	9.0329	7.8885	6.5537	7.9217	
.....	.022	.022	.007	.004	.024	.026	.029	.108	.008	.000	.008	
.....	112	290	219	329	110	108	106	217	291	182	106	
M <sub>2</sub> .....	6.6332	8.3503	-----	8.3503	7.8740	5.7100	6.5537	6.9049	8.4169	6.5537	8.4169	6.5549
.....	.000	.022	-----	.022	.007	.000	.000	.001	.026	.000	.026	.000
.....	2	70	---	290	219	0	178	177	108	182	252	177
N <sub>2</sub> .....	7.8760	7.8740	8.3503	-----	8.3503	7.8824	7.8885	7.8944	6.5537	8.4169	9.0329	7.6658
.....	.008	.007	.022	-----	.022	.008	.008	.008	.000	.026	.026	.005
.....	73	141	70	---	290	71	69	67	178	252	143	67
2N.....	7.5509	7.6166	7.8740	8.3503	-----	7.5772	7.6024	7.6268	7.8885	9.0329	8.0586	7.4452
.....	.004	.004	.007	.022	-----	.004	.004	.004	.008	.108	.011	.003
.....	143	31	141	70	---	141	139	138	69	143	33	138
R <sub>2</sub> .....	8.0074	8.3758	5.7100	7.8824	7.5772	-----	8.0074	8.0072	8.3235	6.1771	7.9055	6.9049
.....	.010	.024	.000	.008	.004	-----	.010	.010	.021	.000	.008	.001
.....	2	250	0	289	219	---	358	356	288	181	252	177
S <sub>2</sub> .....	8.0072	8.4169	6.5537	7.8885	7.6024	8.0074	-----	8.0074	8.3503	6.5549	7.9217	6.5537
.....	.010	.026	.000	.008	.004	.010	-----	.010	.022	.000	.008	.000
.....	4	252	182	291	221	2	---	358	290	183	254	178
T <sub>2</sub> .....	8.0076	8.4607	6.9049	7.8944	7.6268	8.0072	8.0074	-----	8.3792	6.7601	7.9377	5.7100
.....	.010	.029	.001	.008	.004	.010	.010	-----	.024	.001	.009	.000
.....	6	254	183	293	222	4	2	---	291	185	256	0
λ <sub>2</sub> .....	8.2982	9.0329	8.4164	6.5537	7.8885	8.3235	8.3503	8.3792	-----	7.9217	7.9044	7.8885
.....	.020	.108	.026	.000	.008	.021	.022	.024	-----	.008	.008	.008
.....	74	143	252	182	291	72	70	69	-----	254	324	69
μ <sub>2</sub> .....	5.7099	7.8885	6.5537	8.4169	9.0329	6.1771	6.5549	6.7601	7.9217	-----	8.3503	6.5542
.....	.000	.008	.000	.026	.108	.108	.000	.000	.008	-----	.022	.000
.....	0	69	178	108	217	179	177	175	106	-----	70	175
ν <sub>2</sub> .....	7.8892	6.5537	8.4169	9.0329	8.0586	7.9055	7.9217	7.9377	7.9044	8.3503	-----	7.6990
.....	.008	.000	.026	.108	.011	.008	.008	.009	.008	.022	-----	.005
.....	110	178	108	217	327	108	106	104	36	290	-----	105
2SM.....	7.1088	7.9217	6.5549	7.6658	7.4452	6.9049	6.5537	5.7100	7.8885	6.5542	7.6990	-----
.....	.001	.008	.000	.005	.003	.001	.000	.000	.008	.000	.005	-----
.....	185	254	183	293	222	183	182	0	291	185	255	-----

Table 30.—*Products of amplitudes and angular functions for Form 245*

°	1		2		3		4		5		°
	Sin	Cos									
0	0.000	1.000	0.000	2.000	0.000	3.000	0.000	4.000	0.000	5.000	90
1	.017	1.000	.035	2.000	.052	3.000	.070	3.999	.087	4.999	89
2	.035	0.999	.070	1.999	.105	2.998	.140	3.998	.174	4.997	88
3	.052	.999	.105	1.997	.157	2.996	.209	3.995	.262	4.993	87
4	.070	.998	.140	1.995	.209	2.993	.279	3.990	.349	4.988	86
5	.087	.996	.174	1.992	.261	2.989	.349	3.985	.436	4.981	85
6	.105	.995	.209	1.989	.314	2.984	.418	3.978	.523	4.973	84
7	.122	.993	.244	1.985	.366	2.978	.487	3.970	.609	4.963	83
8	.139	.990	.278	1.981	.418	2.971	.557	3.961	.696	4.951	82
9	.156	.988	.313	1.975	.469	2.963	.626	3.951	.782	4.938	81
10	.174	.985	.347	1.970	.521	2.954	.695	3.939	.868	4.924	80
11	.191	.982	.382	1.963	.572	2.945	.763	3.927	.954	4.908	79
12	.208	.978	.416	1.956	.624	2.934	.832	3.913	1.040	4.891	78
13	.225	.974	.450	1.949	.675	2.923	.900	3.897	1.125	4.872	77
14	.242	.970	.484	1.941	.726	2.911	.968	3.881	1.210	4.852	76
15	.259	.966	.518	1.932	.776	2.898	1.035	3.864	1.294	4.830	75
16	.276	.961	.551	1.923	.827	2.884	1.103	3.845	1.378	4.806	74
17	.292	.956	.585	1.913	.877	2.869	1.169	3.825	1.462	4.782	73
18	.309	.951	.618	1.902	.927	2.853	1.236	3.804	1.545	4.755	72
19	.326	.946	.651	1.891	.977	2.837	1.302	3.782	1.628	4.728	71
20	.342	.940	.684	1.870	1.026	2.819	1.368	3.759	1.710	4.698	70
21	.358	.934	.717	1.867	1.075	2.801	1.433	3.734	1.792	4.668	69
22	.375	.927	.749	1.854	1.124	2.782	1.498	3.709	1.873	4.636	68
23	.391	.920	.781	1.841	1.172	2.762	1.563	3.682	1.954	4.602	67
24	.407	.914	.813	1.827	1.220	2.741	1.627	3.654	2.034	4.568	66
25	.423	.906	.845	1.813	1.268	2.719	1.690	3.625	2.113	4.532	65
26	.438	.899	.877	1.798	1.315	2.696	1.753	3.595	2.192	4.494	64
27	.454	.891	.908	1.782	1.362	2.673	1.816	3.564	2.270	4.455	63
28	.469	.883	.939	1.766	1.408	2.649	1.878	3.532	2.347	4.415	62
29	.485	.875	.970	1.749	1.454	2.624	1.939	3.498	2.424	4.373	61
30	.500	.866	1.000	1.732	1.500	2.598	2.000	3.464	2.500	4.330	60
31	.515	.857	1.030	1.714	1.545	2.572	2.060	3.429	2.575	4.286	59
32	.530	.848	1.060	1.696	1.590	2.544	2.120	3.392	2.650	4.240	58
33	.545	.839	1.089	1.677	1.634	2.516	2.179	3.355	2.723	4.193	57
34	.559	.829	1.118	1.658	1.678	2.487	2.237	3.316	2.796	4.145	56
35	.574	.819	1.147	1.638	1.721	2.457	2.294	3.277	2.868	4.096	55
36	.588	.809	1.176	1.618	1.763	2.427	2.351	3.236	2.939	4.045	54
37	.602	.799	1.204	1.597	1.805	2.396	2.407	3.195	3.009	3.993	53
38	.616	.788	1.231	1.576	1.847	2.364	2.463	3.152	3.078	3.940	52
39	.629	.777	1.259	1.554	1.888	2.331	2.517	3.109	3.147	3.886	51
40	.643	.766	1.286	1.532	1.928	2.298	2.571	3.064	3.214	3.830	50
41	.656	.755	1.312	1.509	1.968	2.264	2.624	3.019	3.280	3.774	49
42	.669	.743	1.338	1.486	2.007	2.229	2.677	2.973	3.346	3.716	48
43	.682	.731	1.364	1.463	2.046	2.194	2.728	2.925	3.410	3.657	47
44	.695	.719	1.389	1.439	2.084	2.158	2.779	2.877	3.473	3.597	46
45	.707	0.707	1.414	1.414	2.121	2.121	2.828	2.828	3.536	3.536	45
	Cos	Sin									
	1		2		3		4		5		

Table 30.—*Products of amplitudes and angular functions for Form 245—Continued*

o	6		7		8		9		o
	Sin	Cos	Sin	Cos	Sin	Cos	Sin	Cos	
0	0.000	6.000	0.000	7.000	0.000	8.000	0.000	9.000	90
1	.105	5.999	.122	6.999	.140	7.999	.157	8.999	89
2	.209	5.996	.244	6.996	.279	7.995	.314	8.995	88
3	.314	5.992	.366	6.990	.419	7.989	.471	8.988	87
4	.419	5.985	.488	6.983	.558	7.980	.628	8.978	86
5	.523	5.977	.610	6.973	.697	7.970	.784	8.966	85
6	.627	5.967	.732	6.962	.836	7.956	.941	8.951	84
7	.731	5.955	.853	6.948	.975	7.940	1.097	8.933	83
8	.835	5.942	.974	6.932	1.113	7.922	1.253	8.912	82
9	.939	5.926	1.095	6.914	1.251	7.902	1.408	8.899	81
10	1.042	5.909	1.216	6.894	1.389	7.878	1.563	8.863	80
11	1.145	5.890	1.336	6.871	1.526	7.853	1.717	8.835	79
12	1.247	5.869	1.455	6.847	1.663	7.825	1.871	8.803	78
13	1.350	5.846	1.575	6.821	1.800	7.795	2.025	8.769	77
14	1.452	5.822	1.693	6.792	1.935	7.762	2.177	8.733	76
15	1.553	5.796	1.812	6.762	2.071	7.727	2.329	8.693	75
16	1.654	5.768	1.929	6.729	2.205	7.690	2.481	8.651	74
17	1.754	5.738	2.047	6.694	2.339	7.650	2.631	8.607	73
18	1.854	5.706	2.163	6.657	2.472	7.608	2.781	8.560	72
19	1.953	5.673	2.279	6.619	2.605	7.564	2.930	8.510	71
20	2.052	5.638	2.394	6.578	2.736	7.518	3.078	8.457	70
21	2.150	5.601	2.509	6.535	2.867	7.469	3.225	8.402	69
22	2.248	5.563	2.622	6.490	2.997	7.417	3.371	8.345	68
23	2.344	5.523	2.735	6.444	3.126	7.364	3.517	8.284	67
24	2.440	5.481	2.847	6.395	3.254	7.308	3.661	8.222	66
25	2.536	5.438	2.958	6.344	3.381	7.250	3.804	8.157	65
26	2.630	5.393	3.069	6.292	3.507	7.190	3.945	8.089	64
27	2.724	5.346	3.178	6.237	3.632	7.128	4.086	8.019	63
28	2.817	5.298	3.286	6.181	3.756	7.064	4.225	7.947	62
29	2.909	5.248	3.394	6.122	3.878	6.997	4.363	7.872	61
30	3.000	5.196	3.500	6.062	4.000	6.928	4.500	7.794	60
31	3.090	5.143	3.605	6.000	4.120	6.857	4.635	7.715	59
32	3.180	5.088	3.709	5.936	4.239	6.784	4.769	7.632	58
33	3.268	5.032	3.812	5.871	4.357	6.709	4.902	7.548	57
34	3.355	4.974	3.914	5.803	4.474	6.632	5.033	7.461	56
35	3.441	4.915	4.015	5.734	4.589	6.553	5.162	7.372	55
36	3.527	4.854	4.115	5.663	4.702	6.472	5.290	7.281	54
37	3.611	4.792	4.213	5.590	4.815	6.389	5.416	7.188	53
38	3.694	4.728	4.310	5.516	4.925	6.304	5.541	7.092	52
39	3.776	4.663	4.405	5.440	5.035	6.217	5.664	6.994	51
40	3.857	4.596	4.500	5.362	5.142	6.128	5.785	6.894	50
41	3.936	4.528	4.592	5.283	5.248	6.038	5.905	6.792	49
42	4.015	4.459	4.684	5.202	5.353	5.945	6.022	6.688	48
43	4.092	4.388	4.774	5.119	5.456	5.851	6.138	6.582	47
44	4.168	4.316	4.863	5.035	5.557	5.755	6.252	6.474	46
45	4.243	4.243	4.950	4.950	5.657	5.657	6.364	6.364	45
	Cos	Sin	Cos	Sin	Cos	Sin	Cos	Sin	
	6		7		8		9		

Table 31.—For construction of primary stencils

Difference	Constituent 2Q											
Hour	d.	h.	d.	h.	d.	h.	d.	h.	d.	h.	d.	h.
0	1	0	7	21	14	21	21	20*	28	20*	35	20
+23	-1	4	8	4	15	4	22	3*	29	3*	36	3
+22	-2	11	11	11	10*	10*	10*	10*	17	17	17	10*
+21	-3	18	18	18	17*	17*	17*	17*	17	17	16*	16*
+20	-4	2	1	9	1	16	1	23	0*	30	0*	37
+19	-5	8	8	8	7*	7*	7*	7*	7	7	51	6*
+18	-6	15	15	15	14*	14*	14	14	14	13*	13*	13*
+17	-7	22	22	21*	21*	21*	21	21	21	20*	20*	20*
+16	-8	3	5	10	5	17	4*	24	4*	31	4*	38
+15	-9	12	12	11*	11*	11*	11*	11*	11	11	10*	10*
+14	-10	19	19	18*	18*	18*	18*	18*	18	18	17*	17*
+13	-11	4	2	11	2	18	1*	25	1*	32	1*	39
+12	-12	9	9	8*	8*	8*	8	8	8	7*	7*	7*
+11	-13	16	16	15*	15*	15*	15	15	15	14*	14*	14*
+10	-14	23	23	22*	22*	22*	22	22	22	21*	21*	21*
+9	-15	5	6	12	6	19	5*	26	5*	33	5	40
+8	-16	13	13	12*	12*	12*	12	12	12	11*	11*	11*
+7	-17	20	20	19*	19*	19*	19	19	19	18*	18*	18*
+6	-18	6	3	13	3	20	2*	27	2*	34	2	41
+5	-19	10	10	9*	9*	9*	9	9	9	8*	8*	8*
+4	-20	17	17	16*	16*	16*	16	16	16	15*	15*	15*
+3	-21	7	0	14	0	23*	23*	23*	23	23	22*	22*
+2	-22	7	7	21	6*	28	6*	35	6	42	6	49
+1	-23	14	14	13*	13*	13*	13	13	13	12*	12*	12*

Difference	Constituent 2Q												
Hour	d.	h.	d.	h.	d.	h.	d.	h.	d.	h.	d.	h.	
0	70	19	77	19	84	18*	91	18*	98	18	105	18	
+23	-1	71	2	78	2	85	1*	92	1*	99	1	105	1
+22	-2	9	9	8*	8*	8*	8	8	8	7*	7*	7	
+21	-3	16	16	15*	15*	15*	15	15	15	14*	14*	14	
+20	-4	23	23	22*	22*	22*	22	22	22	21*	21*	21*	
+19	-5	72	6	79	5*	86	5*	93	5*	100	5	107	5
+18	-6	13	12*	12*	12*	12*	12	12	12	11*	11*	11	
+17	-7	20	19*	19*	19*	19*	19	19	19	18*	18*	18	
+16	-8	73	3	80	2*	87	2*	94	2*	101	2	108	2
+15	-9	10	9*	9*	9*	9*	9	9	9	8*	8*	8	
+14	-10	17	16*	16*	16*	16*	16	16	16	15*	15*	15	
+13	-11	74	0	23*	23*	23*	23	23	23	22*	22*	22	
+12	-12	7	81	6*	88	6*	95	6	102	6	109	6	
+11	-13	14	13*	13*	13*	13	13	13	13	12*	12*	12	
+10	-14	21	20*	20*	20*	20	20	20	20	19*	19*	19	
+9	-15	75	4	82	3*	89	3*	96	3	103	3	110	3
+8	-16	11	10*	10*	10*	10	10	10	10	9*	9*	9	
+7	-17	18	17*	17*	17*	17	17	17	16*	16*	16*	16	
+6	-18	76	1	83	0*	90	0*	97	0	104	0	23*	23*
+5	-19	8	7*	7*	7*	7	7	7	111	6*	118	6*	
+4	-20	15	14*	14*	14*	14	14	14	13*	13*	13*	13	
+3	-21	22	21*	21*	21*	21	21	21	20*	20*	20*	20	
+2	-22	77	5	84	4*	91	4*	98	4	105	4	112	3*
+1	-23	12	11*	11*	11*	11	11	11	10*	10*	10	10	

TABLE 31.—*For construction of primary stencils—Continued*

Difference		Constituent 2Q													
Hour		d.	h.	d.	h.	d.	h.	d.	h.	d.	h.	d.	h.		
0	140	17	147	16*	154	16*	161	16	168	16	175	15*	182	15*	
+23	-1	141	0	23*	23*	23*	23	23	23	22*	22*	22*	22	22	
+22	-2	7	148	6*	155	6*	162	6	169	6	176	5*	183	5*	
+21	-3	14	13*	13*	13*	13	13	13	13	12*	12*	190	5	197	5
+20	-4	20*	20*	20*	20*	20	20	20	20	19*	19*	19	19	19	19
+19	-5	142	3*	149	3*	156	3*	163	3	170	3	177	2*	184	2*
+18	-6	10*	10*	10*	10*	10	10	10	10	9*	9*	9	9	9	8
+17	-7	17*	17*	17*	17*	17	17	17	17	16*	16*	16	16	16	15*
+16	-8	143	0*	150	0*	157	0*	164	0	171	0	23*	23*	23	23
+15	-9	7*	7*	7*	7*	7	7	7	7	178	6*	185	6*	192	6
+14	-10	14*	14*	14	14	14	14	14	14	13*	13*	13	13	13	12*
+13	-11	21*	21*	21	21	21	21	21	21	20*	20*	20	20	20	19*
+12	-12	144	4*	151	4*	158	4	165	4	172	4	179	3*	186	3*
+11	-13	11*	11*	11*	11	11	11	11	11	10*	10*	10	10	10	9
+10	-14	18*	18*	18*	18	18	18	18	18	17*	17*	17	17	17	16*
+9	-15	145	1*	152	1*	159	1	166	1	173	0*	180	0*	187	0*
+8	-16	8*	8*	8*	8	8	8	8	8	7*	7*	7	7	7	6*
+7	-17	15*	15*	15	15	15	15	15	14*	14*	14*	14	14	14	13*
+6	-18	22*	22*	22	22	22	22	22	21*	21*	21*	21	21	21	20*
+5	-19	146	5*	153	5*	160	5	167	5	174	4*	181	4*	188	4*
+4	-20	12*	12*	12	12	12	12	12	11*	11*	11*	11	11	11	10*
+3	-21	19*	19*	19	19	19	19	19	18*	18*	18*	18	18	18	17*
+2	-22	147	2*	154	2*	161	2	168	2	175	1*	182	1*	189	1
+1	-23	9*	9*	9	9	9	9	9	8*	8*	8*	8	8	8	7*

Table 31.—*For construction of primary stencils—Continued*

Difference	Constituent 2Q													
	Hour	d.	h.	d.	h.	d.	h.	d.	h.	d.	h.	d.	h.	
0	280	12*	287	12	294	12	301	11*	308	11*	315	11	322	11
+23	-1	19*	19	19	19	18*	18*	18*	18	18	18	17*	17*	
+22	-2	281	2*	288	2	295	2	302	1*	309	1*	316	1	
+21	-3	9*	9	9	9	8*	8*	8*	8	8	8	7*	7*	
+20	-4	16*	16	16	16	15*	15*	15*	15	15	15	14*	14*	
+19	-5	23*	23	23	23	22*	22*	22*	22	22	22	21*	21	
+18	-6	282	6*	289	6	296	6	303	5*	303	5*	317	5	
+17	-7	13	13	13	13	12*	12*	12*	12	12	12	11*	11*	
+16	-8	20	20	20	20	19*	19*	19*	19	19	19	18*	18*	
+15	-9	283	3	290	3	297	3	304	2*	311	2*	318	2	
+14	-10	10	10	10	10	9*	9*	9*	9	9	9	8*	8*	
+13	-11	17	17	17	17	16*	16*	16*	16	16	16	15*	15*	
+12	-12	284	0	291	0	23*	23*	23*	23	23	23	22*	22*	
+11	-13	7	7	298	6*	305	6*	312	6*	319	6	326	6	
+10	-14	14	14	13*	13*	13*	13*	13*	13	13	13	12*	12*	
+9	-15	21	21	20*	20*	20*	20*	20*	20	20	20	19*	19*	
+8	-16	285	4	292	4	299	3*	306	3*	313	3*	320	3	
+7	-17	11	11	10*	10*	10*	10*	10*	10	10	10	9*	9*	
+6	-18	18	18	17*	17*	17*	17	17	17	17	17	16*	16*	
+5	-19	286	1	293	1	300	0*	307	0*	314	0	321	0	
+4	-20	8	8	7*	7*	7*	7	7	7	7	7	335	6*	
+3	-21	15	15	14*	14*	14*	14	14	14	14	14	13*	13*	
+2	-22	22	22	21*	21*	21*	21	21	21	21	21	20*	20*	
+1	-23	287	5	294	5	301	4*	308	4*	315	4	322	4	
Difference	Constituent 2Q						Constituent Q							
Hour	d.	h.	d.	h.	d.	h.	d.	h.	d.	h.	d.	h.	d.	
0	350	10	357	10	364	9*	1	0	10	5	19	13*	28	22*
+23	-1	17	17	17	16*	15*	14	14	23	29	8	16*	48	1*
+22	-2	351	0	358	0	23*	15	23*	20	8*	17	39	2	11
+21	-3	7	7	365	6*	2	0	11	9	18	30	2*	11*	20
+20	-4	14	14	13*	13*	9*	9*	18*	21	3	12	21	49	5*
+19	-5	21	21	20*	20*	19	12	3*	12*	21*	21*	40	6	15
+18	-6	352	4	359	4	366	3*	3	4*	13	22	31	6*	15*
+17	-7	11	11	10*	10*	13*	22*	22*	22	7*	16	41	1	9*
+16	-8	18	18	17*	17*	23	13	8	16*	32	1*	10*	19	60
+15	-9	353	1	360	1	367	0*	4	8*	17	23	2	11	19*
+14	-10	8	8	7*	7*	17*	14	2*	11*	20	20	42	5	51
+13	-11	15	14*	14*	14*	5	3	12	20*	33	5*	14*	23	61
+12	-12	22	22	21*	21*	21*	12*	21*	24	6	15	23*	52	8*
+11	-13	354	5	361	4*	368	4*	22	15	6*	15*	34	0*	18
+10	-14	12	11*	11*	11*	6	7	16	25	1	9*	18*	53	3*
+9	-15	19	18*	18*	18*	16*	16*	16	1*	10	19	44	4	12*
+8	-16	355	2	362	1*	369	1*	7	2	11	19*	35	4*	13
+7	-17	9	8*	8*	8*	8*	11*	20	26	5	13*	22*	54	7*
+6	-18	16	15*	15*	15*	20*	17	5*	14*	23	45	8	16*	64
+5	-19	23	22*	22*	22*	8	6	15	23*	36	8*	17*	55	2
+4	-20	356	6	363	5*	370	5*	15*	18	0	27	9	46	2*
+3	-21	13	12*	12*	12*	9	1	9*	18*	37	3	12	21	65
+2	-22	20	19*	19*	19*	10	19	28	4	12*	21*	56	6	15
+1	-23	357	3	364	2*		19*	19	4*	13	22	47	7	15*

TABLE 31.—*For construction of primary stencils—Continued*

Difference	Constituent Q															
Hour	d.	h.	d.	h.	d.	h.	d.	h.	d.	h.	d.	h.				
0	66	9*	75	18*	85	3*	94	12	103	21	113	6				
+23	-1	19	76	4	12*		104	6*	15		122	14*				
+22	-2	67	4*	13*	22	95	7	15*	114	0*	123	0				
+21	-3	14	22*	86	7*	16*	105	1	10	18*	133	3*				
+20	-4	23	77	8	17	96	1*	10*	19*	124	4	13				
+19	-5	68	8*	17*	87	2	11	20	115	4*	13*	22*				
+18	-6	18	78	3	11*	20*	106	5	14		134	7*				
+17	-7	69	3*	12	21	97	6	14*	23*	125	8	17*				
+16	-8	12*	21*	88	6*	15	107	0	116	8*	135	2*				
+15	-9	22	79	7	15*	98	0*	9*	18	126	3	11*				
+14	-10	70	7*	16	89	1	10	18*	117	3*	12*	21				
+13	-11	17	80	1*	10*		19	108	4	13	21*	136	6*			
+12	-12	71	2	11	20	99	4*	13*	22	127	7	16	146	0*		
+11	-13	11*	20*	90	5	14		23	118	7*	16*	137	1			
+10	-14	21	81	5*	14*	23*	109	8	17	128	2	10*	17*	156	4	
+9	-15	72	6*	15	91	0	100	8*	17*	119	2*	11	20	147	5	
+8	-16	15*	82	0*	9	18	110	3	11*	20*	138	5*	14	23		
+7	-17	73	1	10	18*	101	3*	12	21	129	6	14*	23*	157	8*	
+6	-18	10*	19	92	4	13	21*	120	6*	15	139	0	148	9	17*	
+5	-19	19*	83	4*	13*	22	111	7	16	130	0*	9*	18	158	3	
+4	-20	74	5	14	22*	102	7*	16*	121	1	10	19	149	3*	12*	
+3	-21	14*	23*	93	8	17	112	1*	10*	19*	140	4	13	22		
+2	-22	75	0	84	8*	17*	103	2*	11	20	131	4*	13*	22*	159	7
+1	-23	9	18	94	3	11*	20*	122	5*	14	23	150	7*	16*		
Difference	Constituent Q															
Hour	d.	h.	d.	h.	d.	h.	d.	h.	d.	h.	d.	h.				
0	160	2	169	10*	178	19*	188	4*	197	13	206	22				
+23	-1	11	20	179	5	13*		22*	207	7*	16	225	15*			
+22	-2	20*	170	5*	14	23	198	8	16*	217	1*	10*	19			
+21	-3	161	6	15	23*	189	8*	17	208	2	11	19*	236	4*		
+20	-4	15*	171	0	180	9	18	199	2*	11*	20	227	5			
+19	-5	162	0*	9*	18*	190	3	12	21	218	5*	14*	23	246	8	
+18	-6	10	19	181	3*	12*	21*	209	6	15	228	0	237	8*	17*	
+17	-7	19*	172	4	13	22	200	6*	15*	219	0*	9	18	247	2*	
+16	-8	163	5	13*	22*	191	7	16	210	1	9*	18*	238	3*	12	
+15	-9	14	23	182	8	16*	201	1*	10	19	229	4	12*	21*		
+14	-10	23*	173	8*	17	192	2	11	19	220	4*	13	22	248	7	
+13	-11	164	9	17*	183	2*	11*	20	211	5	14	22*	239	7*	16	
+12	-12	18*	174	3	12	20*	202	5*	14*	23	230	8	17	249	1*	
+11	-13	165	3*	12*	21*	193	6	15	23*	221	8*	17*	240	2	11	
+10	-14	13	22	184	6*	15*	203	0*	212	9	18	231	2*	11*	20*	
+9	-15	22*	175	7	16	194	1	9*	18*	222	3	12	21	250	5*	
+8	-16	166	7*	16*	185	1*	10	19	213	4	12*	21*	241	6	15	
+7	-17	17	176	2	10*	19*	204	4*	13	22	232	7	15*	251	0*	
+6	-18	167	2*	11*	20	195	5	13*	22*	223	7*	16	242	1	10	
+5	-19	12	20*	186	5*	14*	23	214	8	16*	233	1*	10*	19		
+4	-20	21	177	6	15	23*	205	8*	17*	224	2	11	19*	252	4*	
+3	-21	168	6*	15*	187	0	196	9	18	215	2*	11*	20*	243	5	
+2	-22	16	178	1	9*	18*	206	3	12	21	234	5*	14*	23*		
+1	-23	169	1*	10	19	197	3*	12*	21*	225	6	15	244	0	253	8*

TABLE 31.—*For construction of primary stencils—Continued*

Difference	Constituent Q															
Hour	d.	h.	d.	h.	d.	h.	d.	h.	d.	h.	d.	h.				
0	253	18	263	3	272	11*	281	20*	291	5	300	14				
+23	-1	254	3*	12	21	282	6	14*	23*	310	8	17	329	2		
+22	-2	12*	21*	273	6*	15	292	0	301	9	17*	320	2*			
+21	-3	22	264	7	15*	283	0*	9*	18	311	3	12	20*	339	5*	
+20	-4	255	7*	16*	274	1	10	18*	302	3*	12*	21	330	6	15	
+19	-5	17	265	1*	10*	19*	293	4	13	21*	321	6*	15*	340	0	
+18	-6	256	2	11	20	284	4*	13*	22	312	7	16	331	0*	9*	
+17	-7	11*	20*	275	5	14	23	303	7*	16*	322	1	10	349	19	
+16	-8	21	266	5*	14*	23*	294	8	17	313	2	10*	19*	341	4	
+15	-9	257	6*	15	276	0	285	8*	17*	304	2*	11	20	332	5	13*
+14	-10	15*	267	0*	9*	18	295	3	11*	20*	323	5*	14	23		
+13	-11	258	1	10	18*	286	3*	12*	21	314	6	14*	23*	342	8*	
+12	-12	10*	19	277	4	13	21*	305	6*	15*	324	0	333	9	17*	
+11	-13	20	268	4*	13*	22	296	7	16	315	0*	9*	18*	343	3	
+10	-14	259	5	14	23	287	7*	16*	306	1	10	19	334	3*	12*	
+9	-15	14*	23*	278	8	17	297	1*	10*	319	1*	19*	325	4	13	
+8	-16	260	0	269	8*	17*	288	2*	11	20	318	4*	13*	22*	344	7
+7	-17	9	18	279	3	11*	20*	307	5*	14	23	335	7*	16*		
+6	-18	18*	270	3*	12	21	298	6	14*	23*	326	8*	17	345	2	
+5	-19	261	4	13	21*	289	6*	15	308	0	317	9	17*	336	2*	11*
+4	-20	13*	22	280	7	16	299	0*	9	18	327	3	12	20*		
+3	-21	22*	271	7*	16*	290	1	10	19*	318	3*	12*	21	346	6	
+2	-22	262	8	17	281	1*	10*	19*	309	4	13	21*	337	6*	15*	
+1	-23	17*	272	2	11	20	300	4*	13*	22*	328	7	16	347	0*	
Difference	Constituent Q						Constituent $\rho$									
Hour	d.	h.	d.	h.	d.	h.	d.	h.	d.	h.	d.	h.				
0	347	10	356	19	366	3*	1	0	10	15*	20	11	30	6*	40	2
+23	-1	19*	357	4*	13	5*	11	1	20*	16	12	12	50	7*	60	3
+22	-2	348	5	13*	22*	15*	11	21	6*	31	2	21*	17	12*		
+21	-3	14	23	367	8	2	1*	21	16*	12	41	7*	51	3	22*	
+20	-4	23*	358	8	17	11	12	6*	22	2	21*	17	12*	61	8*	
+19	-5	349	9	18	368	2*	21	16*	12	32	7*	42	3	22*	18	
+18	-6	18*	359	3	12	3	6*	13	2*	22	17*	13	52	8*	62	4
+17	-7	350	3*	12*	21*	16*	12	23	7*	33	3	22*	18	13*		
+16	-8	13	22	369	6*	4	2*	22	17*	13	43	8*	53	4	23*	
+15	-9	22*	360	7	16	12	14	7*	24	3	22*	18*	14	63	9*	
+14	-10	351	8	16*	370	1*	22	17*	13	34	8*	44	4	23*	19	
+13	-11	17	361	2	-----	5	8	15	3*	23	18*	14	54	9*	64	5
+12	-12	352	2*	11*	-----	17*	13	25	8*	35	4	23*	19	15		
+11	-13	12	20*	-----	6	3*	23	18*	14	45	9*	55	5	65	0*	
+10	-14	21	362	6	-----	13	16	9	26	4*	36	0	19*	15	10*	
+9	-15	353	6*	15*	-----	23	18*	14	9*	46	5	56	0*	20		
+8	-16	16	363	1	-----	7	9	17	4*	27	0	19*	15	10*	66	6
+7	-17	354	1*	10	-----	18*	14	9*	37	5*	47	1	20*	16		
+6	-18	10*	19*	-----	8	4*	18	0	19*	15	10*	57	6	67	1*	
+5	-19	20	364	5	-----	14*	10	28	5*	38	1	20*	16	11*		
+4	-20	355	5*	14*	-----	9	0	19*	15	10*	48	6	58	2	21*	
+3	-21	15	23*	-----	10	19	5*	29	1	20*	16	11*	68	7		
+2	-22	356	0	365	9	-----	19*	15*	11	39	6*	49	2	21*	17	
+1	-23	9*	18*	-----	10	5*	20	1	20*	16	11*	59	7	69	2*	

Table 31.—*For construction of primary stencils—Continued*

Difference		Constituent $\rho$											
Hour		d. h.	d. h.	d. h.	d. h.	d. h.	d. h.	d. h.	d. h.	d. h.	d. h.	d. h.	d. h.
+23	-1	69 12*	79 8	89 3*	98 23	108 18*	118 14	128 9*	138 5	148 1	157 20*		
+22	-2	22*	18	13*	99 9	109 4*	119 0	19*	15	10*	158 6		
+21	-3	70 8	80 3*	23	18*	14*	10	129 5*	139 1	20*	16		
+20	-4	18	13*	90 9	100 4*	110 0	19*	15	10*	149 6	159 1*		
+19	-5	71 4	23*	19	14*	10	120 5*	130 1	20*	16	11*		
+18	-6	13*	81 9	91 4*	101 0	19*	15	11	140 6*	150 2	21*		
+17	-7	72 9	82 5	92 0*	20	15*	11	131 6*	141 2	21*	17		
+16	-8	19	14*	10	102 5*	112 1	20*	16	11*	151 7*	161 3		
+15	-9	73 5	83 0*	20	15*	11	122 6*	132 2	21*	17	12*		
+14	-10	14*	10	93 5*	103 1*	21	16*	12	142 7*	152 3	22*		
+13	-11	74 0*	20	15*	11	113 6	123 2	21*	17	12*	162 8		
+12	-12	10*	84 6	94 1*	21	16*	12	133 7*	143 3	22*	18		
+11	-13	20	15*	11	104 6*	114 2	21*	17*	13	153 8*	163 4		
+10	-14	75 6	85 1*	21	16*	12	124 7*	134 3	22*	18	13*		
+9	-15	15*	11*	95 7	105 2*	22	17*	13	141 8*	154 4	23*		
+8	-16	76 1*	21	16*	12	115 7*	125 3	22*	18	14	164 9*		
+7	-17	11*	86 7	96 2*	22	17*	13	135 8*	145 4	23*	19		
+6	-18	21	16*	12	106 8	116 3*	23	18*	14	155 9*	165 5		
+5	-19	77 7	87 2*	22	17*	13	126 8*	136 4	23*	19	14*		
+4	-20	17	12*	97 8	107 3*	23	18*	14	146 9*	156 5	166 0*		
+3	-21	78 2*	22	17*	13	117 8	127 4*	137 0	19*	15	10*		
+2	-22	12*	88 8	98 3*	23	18*	14	9*	147 5	157 0*	20		
+1	-23	22	18	13*	108 9	118 4*	128 0	19*	15	10*	167 6		
Difference		Constituent $\rho$											
Hour		d. h.	d. h.	d. h.	d. h.	d. h.	d. h.	d. h.	d. h.	d. h.	d. h.	d. h.	d. h.
+23	-1	167 16	177 11*	187 7	197 2*	206 22	216 17*	226 13	236 8*	246 4	255 23*		
+22	-2	168 1*	21	16*	12	207 7*	217 3*	23	18*	14	256 9*		
+21	-3	21	11*	178 7	188 2*	22	17*	13	227 8*	237 4	23*	19	
+20	-4	169 7	179 2*	22	198 8	208 3*	23	18*	14	247 9*	257 5		
+19	-5	17	12*	189 8	199 3*	23	18*	14	9*	248 5	258 0*		
+18	-6	170 2*	22	17*	13*	209 9	219 4*	229 0	19*	15	10*		
+17	-7	12*	180 8	190 3*	23	18*	14	9*	239 5	249 0*	20		
+16	-8	22*	18	13*	200 9	210 4*	220 0	19*	15	10*	259 6		
+15	-9	171 8	181 3*	23	18*	14	10	230 5*	240 1	20*	16		
+14	-10	18	13*	191 9	201 4*	211 0	19*	15	10*	250 6	260 1*		
+13	-11	172 4	23*	19	14*	10	221 5*	231 1	20*	16	11*		
+12	-12	13*	182 9	192 4*	202 0	19*	15	10*	241 6*	251 2	21*		
+11	-13	23*	19	14*	10	212 5*	222 1	20*	16	11*	261 7		
+10	-14	173 9	183 4*	193 0*	20	15*	11	232 6*	242 2	21*	17		
+9	-15	19	14*	10	203 5*	213 1	20*	16	11*	252 7	262 3		
+8	-16	174 5	184 0*	20	15*	11	223 6*	233 2	21*	17	12*		
+7	-17	14*	10	194 5*	204 1	20*	16*	12	243 7*	253 3	22*		
+6	-18	175 0*	20	15*	11	214 6*	224 2	21*	17	12*	263 8		
+5	-19	10*	185 6	195 1*	21	16*	12	234 7*	244 3	22*	18		
+4	-20	20	15*	11	205 6*	215 2	21*	17	13	254 8*	264 4		
+3	-21	176 6	186 1*	21	16*	12	225 7*	235 3	22*	18	13*		
+2	-22	15*	11	196 7	206 2*	22	17*	13	245 8*	255 4	23*		
+1	-23	177 1*	21	16*	12	216 7*	226 3	22*	18	13*	265 9*		

Table 31.—*For construction of primary stencils—Continued*

Difference	Constituent $\rho$																			
Hour	d.	h.	d.	h.	d.	h.	d.	h.	d.	h.	d.	h.								
0	265	19	275	14*	285	10	295	5*	305	1	314	20*	324	16						
+23	-1	266	5	276	0*	20	15*	11	315	6*	325	2	21*	17						
+22	-2	14*	10	286	6	296	1*	21	16*	12	355	7*	345	3						
+21	-3	267	0*	20	15*	11	306	6*	316	2	21*	17	12*	355	8*					
+20	-4	10*	277	6	287	1*	21	16*	12	326	7*	336	3	22*	18					
+19	-5	20	15*	11	297	6*	307	2*	22	17*	13	346	8*	356	4					
+18	-6	268	6	278	1*	21	16*	12	317	7*	327	3	22*	18	13*					
+17	-7	16	11*	288	7	298	2*	22	17*	13	337	8*	347	4	23*					
+16	-8	269	1*	21	16*	12	308	7*	318	3	23	18*	14	357	9*					
+15	-9	11*	279	7	289	2*	22	17*	13	328	8*	338	4	23*	19					
+14	-10	21	16*	12*	299	8	309	3*	23	18*	14	348	9*	358	5					
+13	-11	270	7	280	2*	22	17*	13	319	8*	329	4	23*	19	15					
+12	-12	17	12*	290	8	300	3*	23	18*	14	339	9*	349	5	359	0*				
+11	-13	271	2*	22	17*	13	310	9	320	4*	330	0	19*	15	10*					
+10	-14	12*	281	8	291	3*	23	18*	14	9*	340	5	350	0*	20					
+9	-15	22*	18	13*	301	9	311	4*	321	0	19*	15	10*	360	6					
+8	-16	272	8	282	3*	23	18*	14	9*	331	5*	341	1	20*	16					
+7	-17	18	13*	292	9	302	4*	312	0	19*	15	10*	351	6	361	1*				
+6	-18	273	3*	23*	19	14*	10	322	5*	332	1	20*	16	11*						
+5	-19	13*	283	9	293	4*	303	0	19*	15	10*	342	6	352	2					
+4	-20	23*	19	14*	10	313	5*	323	1	20*	16	11*	362	7						
+3	-21	274	9	284	4*	294	0	19*	15*	11	333	6*	343	2	21*					
+2	-22	19	14*	10	304	5*	314	1	20*	16	11*	353	7	363	2*					
+1	-23	275	5	285	0*	20	15*	11	324	6*	334	2	21*	17	12*					
Difference	$\rho$	Constituent O																		
Hour	d.	h.	d.	h.	d.	h.	d.	h.	d.	h.	d.	h.	d.	h.						
0	363	22*	1	0	14	22*	29	3	43	7*	57	12	71	16*	85	21	100	2	114	6*
+23	-1	364	8	8	15	12*	17	21*	58	2	72	7	86	11*	16	20*				
+22	-2	18	22	16	2*	30	7	44	12	16*	21	87	1*	101	6	115	11			
+21	-3	365	4	2	12	17	21*	45	2	59	6*	73	11	16	20*	116	1			
+20	-4	13*	3	2*	17	7	31	11*	16	21	74	1*	88	6	102	10*	15			
+19	-5	23*	16*	21	32	2	46	6*	60	11	15*	20	103	1	117	5*				
+18	-6	366	9	4	7	18	11*	16	20*	61	1	75	6	89	10*	15	19*			
+17	-7	19	21	19	1*	33	6	47	11	15*	20	90	0*	104	5	118	10			
+16	-8	367	5	5	11	16	20*	48	1	62	5*	76	10	15	19*	119	0			
+15	-9	14*	6	1*	20	6	34	10*	15	20	77	0*	91	5	105	9*	14			
+14	-10	368	0*	15*	20	35	1	49	5*	63	10	14*	19	106	0	120	4*			
+13	-11	10*	7	6	21	10*	15	19*	64	0	78	5	92	9*	14	18*				
+12	-12	20	20	22	0*	36	5	50	9*	14*	19	23*	107	4	121	8*				
+11	-13	369	6	8	10	14*	19*	51	0	65	4*	79	9	93	13*	18*	23			
+10	-14	15*	9	0*	23	5	37	9*	14	18*	23*	94	4	108	8*	122	13			
+9	-15	370	1*	14*	19	23*	52	4*	66	9	80	13*	18	22*	123	3*				
+8	-16	-----	10	4*	24	9*	38	14	18*	23	81	3*	95	8*	109	13	17*			
+7	-17	-----	19	23*	39	4	53	8*	67	13*	18	22*	110	3	124	7*				
+6	-18	-----	11	9	25	13*	18*	23	68	3*	82	8	96	12*	17*	22				
+5	-19	-----	23*	26	4	40	8*	54	13	17*	22*	97	3	111	7*	125	12			
+4	-20	-----	12	13*	18	22*	55	3*	69	8	83	12*	17	21*	126	2*				
+3	-21	-----	13	3*	27	8*	41	13	17*	22	84	2*	98	7*	112	12	16*			
+2	-22	-----	18	22*	42	3	56	7*	70	12*	17	21*	113	2	127	6*				
+1	-23	-----	14	8	28	12*	17*	22	71	2*	85	7	99	11*	16	21				

Table 31.—*For construction of primary stencils—Continued*

Difference	Constituent O																
Hours	d.	h.	d.	h.	d.	h.	d.	h.	d.	h.	d.	h.	d.	h.			
0	128	11	142	15*	156	20	171	1	185	5*	199	10	213	14*			
+23	-1	129	1	143	6	157	10*	15	19*	200	5	214	5	228	9*		
+22	-2	15*	20	158	0*	172	5	186	10	14*	19	23*	243	4	257	9	
+21	-3	130	5*	144	10	15	19*	187	0	201	4*	215	9	229	14	18*	23
+20	-4	20	145	0*	159	5	173	9*	14	19	23*	230	4	244	8*	258	13
+19	-5	131	10	14*	19	174	0	188	4*	202	9	216	13*	18	23	259	3*
+18	-6	132	0	146	5	160	9*	14	18*	23	217	4	231	8*	245	13	17*
+17	-7	14*	19	23*	175	4	189	9	203	13*	18	22*	246	3	260	8	
+16	-8	133	4*	147	9	161	14	18*	23	204	3*	218	8	232	13	17*	22
+15	-9	19	23*	162	4	176	8*	190	13	18	22*	233	3	247	7*	261	12
+14	-10	134	9	148	13*	18	22*	191	3*	205	8	219	12*	17	21*	262	2*
+13	-11	23	149	3*	163	8*	177	13	17*	22	220	2*	234	7*	248	12	16*
+12	-12	135	13*	18	22*	178	3	192	7*	206	12*	17	21*	249	2	263	6*
+11	-13	136	3*	150	8	164	12*	17*	22	207	2*	221	7	235	11*	16*	21
+10	-14	17*	22*	165	3	179	7*	193	12	16*	21*	236	2	250	6*	264	11
+9	-15	137	8	151	12*	17	21*	194	2*	208	7	222	11*	16	20*	265	1*
+8	-16	22	152	2*	166	7*	180	12	16*	21	223	1*	237	6*	251	11	15*
+7	-17	138	12*	17	21*	181	2	195	6*	209	11*	16	20*	252	1	266	5*
+6	-18	139	2*	153	7	167	11*	16*	21	210	1*	224	6	238	10*	15*	20
+5	-19	16*	21*	168	2	182	6*	196	11	15*	20*	239	1	253	5*	267	10
+4	-20	140	7	154	11*	16	20*	197	1*	211	6	225	10*	15	19*	268	0*
+3	-21	21	155	1*	169	6*	183	11	15*	20	226	0*	240	5	254	10	14*
+2	-22	141	11*	16	20*	184	1	198	5*	212	10	15	19*	255	0	269	4*
+1	-23	142	1*	156	6	170	10*	15	20	213	0*	227	5	241	9*	14	19

Difference	Constituent O												Component 2N					
Hour	d.	h.	d.	h.	d.	h.	d.	h.	d.	h.	d.	h.	d.	h.				
0	270	9	284	13*	298	18	312	23	327	3*	341	8	355	12*	369	17	1	0
+23	-1	23	285	4	299	8*	313	13	17*	22	356	3	370	7*	8	15	14	23*
+22	-2	271	13*	18	22*	314	3	328	8	342	12*	17	-----	-----	22	16	4	-----
+21	-3	272	3*	286	8	300	13	17*	22	343	2*	357	7	-----	2	12*	18*	-----
+20	-4	18	22*	301	3	315	7*	329	12	17	21*	-----	3	2*	17	8*	-----	17
+19	-5	273	8	287	12*	17	22	330	2*	344	7	358	11*	-----	17	23	-----	-----
+18	-6	22	288	3	302	7*	316	12	16*	21	359	2	-----	4	7	18	13*	-----
+17	-7	274	12*	17	21*	317	2	331	7	345	11*	16	-----	21*	19	3*	-----	-----
+16	-8	275	2*	289	7	303	11*	16*	21	346	1*	360	6	-----	5	11*	18	-----
+15	-9	16*	21*	304	2	318	6*	332	11	15*	20*	-----	6	2	20	8	-----	-----
+14	-10	276	7	290	11*	16	20*	333	1*	347	6*	361	10*	-----	6	16	22*	-----
+13	-11	21	291	1*	305	6*	319	11	15*	20	362	0*	-----	7	6*	21	12*	-----
+12	-12	277	11*	16	20*	320	1	334	5*	348	10*	15	-----	20*	22	3	-----	-----
+11	-13	278	1*	292	6	306	10*	15*	20	349	0*	363	5	8	11	17	-----	-----
+10	-14	15*	20*	307	1	321	5*	335	10	14*	19*	-----	9	1	23	7*	-----	-----
+9	-15	279	6	293	10*	15	19*	336	0*	350	5	364	9*	15*	21*	-----	-----	-----
+8	-16	20	294	0*	308	5*	322	10	14*	19	23*	-----	10	5*	24	12	-----	-----
+7	-17	280	10*	15	19*	323	0	337	4*	351	9*	365	14	20	25	2	-----	-----
+6	-18	281	0*	295	5	309	9*	14*	19	23*	366	4	-----	11	10	16*	-----	-----
+5	-19	14*	19*	310	0	324	4*	338	9	352	13*	18	-----	12	0*	26	6*	-----
+4	-20	282	5	296	9*	14	18*	23	353	4	367	8*	-----	14*	21	-----	-----	-----
+3	-21	19	23*	311	4	325	9	339	13*	18	22*	-----	13	5	27	11	-----	-----
+2	-22	283	9	297	14	18*	23	340	3*	354	8	368	13	19	28	1*	-----	-----
+1	-23	23*	298	4	312	8*	326	13	18	22*	369	3	-----	14	9*	15*	-----	-----

Table 31.—*For construction of primary stencils—Continued*

Difference	Constituent 2N															
Hour	d.	h.	d.	h.	d.	h.	d.	h.	d.	h.	d.	h.				
0	29	6	43	12	57	18	72	0	86	6	100	12*				
+23	-1	20	44	2	58	8*	14*	20*	101	2*	115	8*				
+22	-2	30	10*	16*	22*	73	4*	87	10*	17	23	130	5			
+21	-3	31	0*	45	6*	59	13	19	88	1	102	7				
+20	-4	15	21	60	3	74	9	15	21*	117	3*	131	9*			
+19	-5	32	5	46	11	17	23*	89	5	103	11*	17*	132	0		
+18	-6	19*	47	1*	61	7°	75	13*	20	104	2	118	8			
+17	-7	33	9	15*	22	76	4	90	10	16	22	133	4*			
+16	-8	34	0	48	6	62	12	18	91	0*	105	6*				
+15	-9	14	20	63	2*	77	8*	14*	20*	120	2*	134	9			
+14	-10	35	4*	49	10*	16*	22*	92	5	106	11	17	23			
+13	-11	18*	50	0*	64	7	78	13	19	107	1	121	7			
+12	-12	36	9	15	21	79	3	93	9*	15*	21*	136	3*			
+11	-13	23	51	5	65	11*	17*	23*	108	5*	122	11*				
+10	-14	37	13*	19*	66	1*	80	7*	94	14	20	123	2			
+9	-15	38	3*	52	9*	16	22	95	4	109	10	16	22*			
+8	-16	18	53	0	67	6	81	12	18*	110	0*	124	6*			
+7	-17	39	8	14	20*	82	2*	96	8*	14*	20*	139	3			
+6	-18	22*	54	4*	68	10*	16*	23	111	5	125	11				
+5	-19	40	12*	19	69	1	83	7	97	13	126	1*				
+4	-20	41	3	55	9	15	21	98	3*	112	9	15*	21*			
+3	-21	17	23*	70	5*	84	11*	17*	23	127	6	141	12			
+2	-22	42	7*	56	13*	19*	85	1*	99	8	113	14	20	142	2	
+1	-23	21*	57	4	71	10	16	22	114	4	128	10*	16*	22*	171	4*
Difference	Constituent 2N															
Hour	d.	h.	d.	h.	d.	h.	d.	h.	d.	h.	d.	h.				
0	171	19	186	1	200	7	214	13	228	19	243	1*				
+23	-1	172	9	15	21*	215	3*	229	9*	15*	21*	271	13*			
+22	-2	23*	187	5*	201	11*	17*	23*	244	6	258	12				
+21	-3	173	13*	19*	202	2	216	8	230	14	20	259	2			
+20	-4	174	4	188	10	16	22	231	4	245	10*	16*	22*			
+19	-5	18	189	0	203	6*	217	12*	18*	246	0*	260	6*			
+18	-6	175	8*	14*	20*	218	2*	232	9	15	21	275	3			
+17	-7	22*	190	4*	204	11	17	23	247	5	261	11				
+16	-8	176	13	19	205	1	219	7	233	13*	262	1*				
+15	-9	177	3	191	9	15*	21*	234	3*	248	9*	15*	22			
+14	-10	17*	23*	206	5*	220	11*	18	249	0	263	6				
+13	-11	178	7*	192	13*	20	221	2	235	8	14	20	278	2*		
+12	-12	22	193	4	207	10	16	22*	250	4*	264	10*				
+11	-13	179	12	18	208	0*	222	6*	236	12*	18*	265	0*			
+10	-14	180	2*	194	8*	14*	20*	237	3	251	9	15	21			
+9	-15	16*	22*	209	5	223	11	17	23	266	5	280	11*			
+8	-16	181	7	195	13	19	224	1	238	7*	252	13*				
+7	-17	21	196	3	210	9*	15*	21*	253	3*	267	10				
+6	-18	182	11*	17*	23*	225	5*	239	12	18	268	0				
+5	-19	183	1*	197	8	211	14	20	240	2	254	8				
+4	-20	16	22	212	4	226	10	16*	22*	269	4*	283	10*			
+3	-21	184	6	198	12*	18*	227	0*	241	6*	255	12*				
+2	-22	20*	199	2*	213	8*	14*	21	256	3	270	9				
+1	-23	185	10*	17	23	228	5	242	11	17	23*	285	5*			

Table 31.—*For construction of primary stencils—Continued*

Difference	Constituent 2N					Constituent $\mu$					
<i>Hour</i>	<i>d. h.</i>	<i>d. h.</i>	<i>d. h.</i>	<i>d. h.</i>	<i>d. h.</i>	<i>d. h.</i>	<i>d. h.</i>	<i>d. h.</i>	<i>d. h.</i>	<i>d. h.</i>	<i>d. h.</i>
0	314 8	328 14	342 20	357 2	1 0	15 11*	30 6	45 0*	59 19	74 13	
+23	-1	22	329 4	343 10*	16*	8	16 2*	21	15	60 9*	75 4
+22	-2	315 12*	18*	344 0*	358 6*	23	17*	31 11*	46 6	61 0*	18*
+21	-3	316 2*	330 8*	15	21	2 13*	17 8	32 2*	21	15	76 9*
+20	-4	17	23	345 5	359 11	3 4*	23	17	47 11*	62 6	77 0*
+19	-5	317 7	331 13	19*	360 1*	19	18 13*	33 8	48 2*	20*	15
+18	-6	21*	332 3*	346 9*	15*	4 10	19 4*	22*	17	63 11*	78 6
+17	-7	318 11	17*	347 0	361 6	5 0*	19	34 13*	49 8	64 2	20*
+16	-8	319 2	333 8	14	20	15*	20 10	35 4	22*	17	79 11*
+15	-9	16	22	348 4*	362 10*	6 6*	21 0*	19	50 13*	65 7*	80 2*
+14	-10	320 6*	334 12*	18*	363 0*	21	15*	36 10	51 4	22*	17
+13	-11	20*	335 2*	349 9	15	7 12	22 6	37 0*	19	66 13*	81 7*
+12	-12	321 11	17	23	364 5	8 2*	21	15*	52 9*	67 4	22*
+11	-13	322 1	336 7	350 13*	19*	17*	23 11*	38 6	53 0*	19	82 13
+10	-14	15*	21*	351 3*	365 9	9 8	24 2*	21	15	68 9*	83 4
+9	-15	323 5*	337 11*	18	366 0	23	17	39 11*	54 6	69 0*	18*
+8	-16	20	338 2	352 8	14	10 13*	25 8	40 2*	20*	15	84 9
+7	-17	324 10	16*	22*	367 4*	11 4*	22*	17	55 11*	70 6	85 0
+6	-18	325 0*	339 6*	353 12*	18*	19	26 13*	41 8	56 2	20*	15
+5	-19	14*	21	354 3	368 9	12 10	27 4*	22*	17	71 11*	86 5*
+4	-20	326 5	340 11	17	23	13 0*	19	42 13*	57 8	72 2	20*
+3	-21	19	341 1*	355 7*	369 13*	15*	28 10	43 4	22*	17	87 11*
+2	-22	327 9*	15*	21*	370 3*	14 6	29 0*	19	58 13*	73 7*	88 2
+1	-23	23*	342 6	356 12	-----	21	15*	44 9*	59 4	22*	17
<i>Difference</i>	Constituent $\mu$										
<i>Hour</i>	<i>d. h.</i>	<i>d. h.</i>	<i>d. h.</i>	<i>d. h.</i>	<i>d. h.</i>	<i>d. h.</i>	<i>d. h.</i>	<i>d. h.</i>	<i>d. h.</i>	<i>d. h.</i>	<i>d. h.</i>
0	89 7*	104 2	118 20*	133 14*	148 9	163 3*	177 22*	192 16	207 10*	222 5	
+23	-1	22*	16*	119 11	134 5*	149 0	18	178 12*	193 7	208 1*	19*
+22	-2	90 13	105 7	120 2	20	14*	164 9	179 3*	21*	16	223 10*
+21	-3	91 4	22	16*	135 11	150 5*	23*	18	194 12*	209 7	224 1
+20	-4	18*	106 13	121 7*	136 1*	20	165 14*	180 9	195 3	21*	16
+19	-5	92 9*	107 4	22	16*	151 11	166 5	23*	18	210 12*	225 6*
+18	-6	93 0	18*	122 13	137 7*	152 1*	20	181 14*	196 8*	211 3	21*
+17	-7	15	108 9*	123 3*	22	16*	167 11	182 5	23*	18	226 12
+16	-8	94 5*	109 0	18*	138 13	153 7	168 1*	20	197 14*	212 8*	227 3
+15	-9	20*	15	124 9	139 3*	22	16*	183 10*	198 5	23*	18
+14	-10	95 11	110 5*	125 0	18*	154 12*	169 7	184 1*	20	213 14	228 8*
+13	-11	96 2	20*	14*	140 9	155 3*	22	16	199 10*	214 5	23*
+12	-12	17	111 11	126 5*	141 0	18	170 12*	185 7	200 1*	19*	229 14
+11	-13	97 7*	112 2	20*	14*	156 9	171 3*	21*	16	215 10*	230 5
+10	-14	22*	16*	127 11	142 5*	157 0	18	186 12*	201 7	216 1	19*
+9	-15	98 13	113 7*	128 2	20	14*	172 9	187 3	21*	16	231 10*
+8	-16	99 4	22	16*	143 11	158 5*	23*	18	202 12*	217 7	232 1
+7	-17	18*	114 13	129 7*	144 1*	20	173 14*	188 9	203 3	21*	16
+6	-18	100 9*	115 3*	22	16*	159 11	174 5	23*	18	218 12*	233 6*
+5	-19	101 0	18*	130 13	145 7	160 1*	20	189 14*	204 8*	219 3	21*
+4	-20	15	116 9	131 3*	22	16*	175 10*	190 5	23*	18	234 12
+3	-21	102 5*	117 0	18*	146 12*	161 7	176 1*	20	205 14	220 8*	235 3
+2	-22	20*	15	132 9*	147 3*	22	16	191 10*	206 5	23*	17*
+1	-23	103 11	118 5*	133 0	18*	162 12*	177 7	192 1*	19*	221 14	236 8*

Table 31.—*For construction of primary stencils—Continued*

Difference	Constituent $\mu$																				
Hour	d.	h.	d.	h.	d.	h.	d.	h.	d.	h.	d.	h.									
0	236	23	251	17*	266	12	281	6*	296	0*	310	19									
+23	-1	237	14	252	8*	267	2*	21	15*	311	10	326	4								
+22	-2	238	5	23		17*	282	12	297	6	312	0*	19								
+21	-3		19*	253	14	268	8*	283	2*	21	15*	327	9*								
+20	-4	239	10*	254	4*	23		17*	298	12	313	6	328	0*							
+19	-5	240	1	19		269	14	284	8	299	2*	21	15*								
+18	-6	16	255	10	270	4*	23		17*	314	11*	329	6								
+17	-7	241	6*	256	1	19*	285	13*	300	8	315	2*	21								
+16	-8	21*		15*	271	10	286	4*	23	17	330	11*	345	6							
+15	-9	242	12	257	6*	272	1	19	301	13*	316	8	331	2*							
+14	-10	243	3	21*		15*	287	10	302	4*	22*	17	346	11*							
+13	-11	17*	258	12	273	6*	288	1	19	317	13*	332	8								
+12	-12	244	8*	259	3	21		15*	303	10	318	4*	22								
+11	-13	23	17*	274	12	289	6*	304	0*	19	333	13*	348	8							
+10	-14	245	14	260	8*	275	2*	21	15*	319	10	334	4	22*							
+9	-15	246	4*	23		17*	290	12	305	6	320	0*	19	349	13*						
+8	-16	19*	261	14	276	8	291	2*	21	15*	335	9*	350	4	22*						
+7	-17	247	10	262	4*	23		17*	306	11*	321	6	336	0*	19						
+6	-18	248	1	19*	277	13*	292	8	307	2*	21	15	351	9*	366	4					
+5	-19	16	263	10	278	4*	23		17	322	11*	337	6	352	0*	18*					
+4	-20	249	6*	264	1	19*	293	13*	308	8	323	2*	20*	15	367	9*					
+3	-21	21*		15*	279	10	294	4*	23	17	338	11*	353	6	368	0					
+2	-22	250	12	265	6*	280	1	19	309	13*	324	8	339	2*	20*						
+1	-23	251	3	21		15*	295	10	310	4*	22*	17	354	11*	369	6					
Difference	Constituent N																				
Hour	d.	h.	d.	h.	d.	h.	d.	h.	d.	h.	d.	h.									
0	1	0	19	20*	39	2	58	7*	77	13	96	18*	116	0	135	5*	154	11	173	16*	
+23	-1	10*	20	16	21*		59	2*	78	8	97	13*	19	136	0*	155	6	174	11*		
+22	-2	2	5*	21	11	40	16*	22	79	3*	98	9	117	14*	20	156	1*	175	6*		
+21	-3	3	1	22	6*	41	11*	60	17	22*	99	4	118	9*	137	15	20*	176	2		
+20	-4	20	23	1*	42	7	61	12*	80	18	23*	119	5	138	10*	157	15*	171	21		
+19	-5	4	15*		20*	43	2	62	7*	81	13	100	18*	120	0	139	5*	158	11	177	16*
+18	-6	5	10*	24	16	21*		63	3	82	8*	101	14	19*	140	0*	159	6	178	11*	
+17	-7	6	5*	25	11	44	16*	22	83	3*	102	9	121	14*	20	160	1*	179	7		
+16	-8	7	1	26	6*	45	12	64	17*	23	103	4	122	9*	141	15	20*	180	2		
+15	-9	20	27	1*	46	7	65	12*	84	18	23*	123	5	142	10*	161	16	181	21*		
+14	-10	8	15*	21		47	2*	66	8	85	13	104	18*	124	0	143	5*	162	11	181	16
+13	-11	9	10*	28	16	21*		67	3	86	8*	105	14	19*	144	1	163	6*	182	12	
+12	-12	10	6	29	11*	48	17	22	87	3*	106	9	125	14*	20	164	1*	183	7		
+11	-13	11	1	30	6*	49	12	68	17*	23	107	4*	126	10	145	15*	21	184	2		
+10	-14	20*	31	2	50	7	69	12*	88	18	23*	127	5	146	10*	165	16	21*			
+9	-15	12	15*	21	51	2*	70	8	89	13*	108	19	128	0*	147	6	166	11	185	16*	
+8	-16	13	11	32	16	21*		71	3	90	8*	109	14	19*	148	1	167	6*	186	12	
+7	-17	14	6	33	11*	52	17	22*	91	4	110	9*	129	15	20	168	1*	187	7		
+6	-18	15	1	34	6*	53	12	72	17	23	111	4*	130	10	149	15*	21	188	2*		
+5	-19	20*	35	2	54	7*	73	13	92	18*	23*	131	5	150	10*	169	16	21*			
+4	-20	16	15*	21	55	2*	74	8	93	13*	112	19	132	0*	151	6*	170	11*	189	17	
+3	-21	17	11	36	16*	22	75	3*	94	8*	113	14	19*	152	1	171	6*	190	12		
+2	-22	18	6	37	11*	56	17	22*	95	4	114	9*	133	15	20*	172	2	191	7*		
+1	-23	19	1*	38	7	57	12*	76	17*	23	115	4*	134	10	153	15*	21	192	2*		

Table 31.—*For construction of primary stencils—Continued*

Difference	Constituent N											
Hour	d.	h.	d.	h.	d.	h.	d.	h.	d.	h.	d.	h.
0	192	21*	212	3	231	8*	250	14	269	19*	289	1
+23	-1	193	17	22*	232	4	251	9*	270	15	290	20
+22	-2	194	12	213	17*	23	252	4*	271	10	290	15*
+21	-3	195	7*	214	13	233	18*	253	0	272	5	
+20	-4	196	2*	215	8	234	13*	254	19	273	0*	
+19	-5	22	216	3*	235	9	254	14	274	19*	293	1
+18	-6	197	17	22*	236	4	255	9*	274	15	295	20*
+17	-7	198	12*	217	18	23	256	4*	275	10	294	15*
+16	-8	199	7*	218	13	237	18*	257	0	276	5*	
+15	-9	200	3	219	8	238	13*	258	19	277	0*	
+14	-10	202	22	220	3*	239	9	258	14*	278	15	
+13	-11	201	17	22*	240	4	259	9*	278	15	297	1*
+12	-12	202	12*	221	18	23*	260	5	279	10*	298	15*
+11	-13	203	7*	222	13	241	18*	261	0	280	5*	
+10	-14	204	3	223	8*	242	14	262	19*	281	0*	
+9	-15	205	22	224	3*	243	9	262	14*	282	15*	
+8	-16	205	17*	23	244	4*	263	9*	282	15	297	10*
+7	-17	206	12*	225	18	23*	264	5	283	10*	302	16
+6	-18	207	8	226	13*	245	18*	265	0	284	5*	
+5	-19	208	3	227	8*	246	14	266	14*	285	1	
+4	-20	209	22*	228	3*	247	9	267	10	286	15*	
+3	-21	210	17*	23	248	4*	268	5	287	10*		
+2	-22	210	12*	229	18	23*	269	0*	288	6	306	16
+1	-23	211	8	230	13*	249	19	269	0*	288	6	

Difference	Constituent ν												
Hour	d.	h.	d.	h.	d.	h.	d.	h.	d.	h.	d.	h.	
0	1	0	20	18*	40	23	61	3	81	7	101	11	
+23	-1	11	21	15	41	19	23	28	3	102	7	121	15
+22	-2	2	27	22	11	42	15	62	19	23	103	3*	
+21	-3	3	3	23	7	43	11*	63	15*	83	19*	123	7*
+20	-4	23*	24	3*	44	7*	64	11*	84	15*	104	19*	
+19	-5	4	19*	23*	45	3*	65	7*	85	12	105	16	
+18	-6	5	15*	25	19*	46	0	66	4	86	8	106	12
+17	-7	6	12	26	16	47	0	67	0	87	4	107	8
+16	-8	7	8	27	12	47	16	20	88	0*	108	4*	
+15	-9	8	4	28	8	48	12*	68	16*	20*	109	0*	
+14	-10	9	0*	29	4*	49	8*	69	12*	89	16*	20*	
+13	-11	20*	30	0*	50	4*	70	8*	90	13	110	17	
+12	-12	10	16*	21	51	1	71	5	91	9	111	13	
+11	-13	11	13	31	17	21	72	1	92	5	112	9	
+10	-14	12	9	32	13	52	17	21	93	1*	113	5*	
+9	-15	13	5	33	9*	53	13*	73	17*	21*	114	1*	
+8	-16	14	1*	34	5*	54	9*	74	13*	94	17*	21*	
+7	-17	21*	35	1*	55	5*	75	9*	95	14	115	18	
+6	-18	15	17*	22	56	2	76	6	96	10	116	14	
+5	-19	16	14	36	18	22	77	2	97	6	117	10	
+4	-20	17	10	37	14	57	18	22	98	2*	118	6*	
+3	-21	18	6	38	10*	58	14*	78	18*	22*	119	2*	
+2	-22	19	2*	39	6*	59	10*	79	14*	99	18*	22*	
+1	-23	22*	40	2*	60	6*	80	10*	100	15	120	19	

Table 31.—*For construction of primary stencils—Continued*

Difference	Constituent ν												
Hour	d.	h.	d.	h.	d.	h.	d.	h.	d.	h.	d.	h.	
+23 0	202	7*	222	11*	242	15*	262	19*	282	23*	303	3	
+23 -1	203	3*	223	7*	243	11*	263	15*	283	19*	303	7*	
+22 -2		23*	224	3*	244	7*	264	12	284	16	304	20	
+21 -3	204	20	225	0	245	4	265	8	285	12	305	16	
+20 -4	205	16	226	20	246	0	266	4	286	8	306	12*	
+19 -5	206	12	226	16	20	267	0*	287	4*	307	8*	326	16*
+18 -6	207	8*	227	12*	247	16*	20*	288	0*	308	4*	328	8*
+17 -7	208	4*	228	8*	248	12*	268	16*	20*	309	1	348	12*
+16 -8	209	0*	229	4*	249	8*	269	13	289	17	309	1	
+15 -9		21	230	1	250	5	270	9	290	13	310	17	
+14 -10	210	17	21	251	1	271	5	291	9	311	13*		
+13 -11	211	13	231	17	21*	272	1*	292	5*	312	9*	331	13*
+12 -12	212	9*	232	13*	252	17*	21*	293	1*	313	5*	333	9*
+11 -13	213	5*	233	9*	253	13*	273	17*	21*	314	2	353	13*
+10 -14	214	1*	234	5*	254	10	274	14	294	18	314	2	
+9 -15		22	235	2	255	6	275	10	295	14	315	18	
+8 -16	215	18	22	256	2	276	6	296	10	316	14*		
+7 -17	216	14	236	18	22*	277	2*	297	6*	317	10*	336	18*
+6 -18	217	10*	237	14*	257	18*	22*	298	2*	318	6*	338	10*
+5 -19	218	6*	238	10*	258	14*	278	18*	22*	319	3	358	14*
+4 -20	219	2*	239	6*	259	11	279	15	299	19	320	3	
+3 -21		23	240	3	260	7	280	11	300	15	320	19	
+2 -22	220	19	23	261	3	281	7	301	11	321	15*		
+1 -23	221	15	241	19	23*	282	3	302	7*	322	11*	342	15*
Difference	Constituent 2 MK												
Hour	d.	h.	d.	h.	d.	h.	d.	h.	d.	h.	d.	h.	
+23 0	1	0	22	7	44	0	65	17	87	10	109	3	
+23 -1		11*	23	4*	44	21*	66	14*	88	7*	110	0*	
+22 -2	2	9*	24	2*	45	19*	67	12*	89	5*	110	22*	
+21 -3	3	7	25	0	46	17	68	10	90	3	111	20	
+20 -4	4	4*	22	22	47	15	69	8	91	1	112	18	
+19 -5	5	2*	26	19*	48	12*	70	5*	22*	113	15*	135	
+18 -6	6	0	27	17	49	10	71	3	92	20	114	13	
+17 -7	22	28	15	50	8	72	1	93	18	115	11		
+16 -8	7	19*	29	12*	51	5*	22*	94	15*	116	8*		
+15 -9	8	17*	30	10*	52	3*	73	20*	95	13*	117	6*	
+14 -10	9	15	31	8	53	1	74	18	96	11	118	4	
+13 -11	10	12*	32	5*	22*	75	16	97	9	119	2		
+12 -12	11	10*	33	3*	54	20*	76	13*	98	6*	23*	141	
+11 -13	12	8	34	1	55	18	77	11	99	4	120	21	
+10 -14	13	6	23	23	56	16	78	9	100	2	121	19	
+9 -15	14	3*	35	20*	57	13*	79	6*	23*	122	16*		
+8 -16	15	1	36	18*	58	11*	80	4*	101	21*	123	14*	
+7 -17	23	37	16	59	9	81	2	102	19	124	12		
+6 -18	16	20*	38	13*	60	6*	23*	103	16*	125	10		
+5 -19	17	18*	39	11*	61	4*	82	21*	104	14*	126	7*	
+4 -20	18	16	40	9	62	2	83	19	105	12	127	5	
+3 -21	19	14	41	7	63	0	84	17	106	10	128	3	
+2 -22	20	11*	42	4*	21*	85	14*	107	7*	129	0*		
+1 -23	21	9	43	2	64	19	86	12*	108	5*	22*	151	

Table 31.—*For construction of primary stencils—Continued*

Difference	Constituent 2MK												Constituent MN							
Hour	d.	h.	d.	h.	d.	h.	d.	h.	d.	h.	d.	h.	d.	h.						
0	217	16	239	9	261	2	282	19	304	12	326	5	347	22	369	15				
+23	-1	218	14	240	7	262	0	283	17	305	10	327	3	348	20	370	13			
+22	-2	219	11*	241	4*	21*	284	14*	306	7*	328	0*	349	17*	-----	2 11	25 18*			
+21	-3	220	9	242	2	263	19*	285	12*	307	5*	322	22*	350	15*	-----	3 11	26 18		
+20	-4	221	7	243	0	264	17*	286	10	308	3	329	20	351	13	-----	4 10*	27 17		
+19	-5	222	4*	244	21*	265	14*	287	7*	309	0*	330	17*	352	10*	-----	5 9*	28 16*		
+18	-6	223	2*	244	19*	266	12*	288	5*	322	15*	353	8*	-----	6 9	29 16				
+17	-7	224	0	245	17	267	10	289	3	310	20	332	13	354	6	-----	7 8	30 15		
+16	-8	22	22	246	15	268	8	290	1	311	18	333	11	355	4	-----	8 7*	31 14*		
+15	-9	225	19*	247	12*	269	5*	22*	312	15*	334	8*	356	1*	-----	9 6*	32 13*			
+14	-10	226	17	248	10	270	3	291	20	313	13*	335	6*	357	23	-----	10 6	33 13		
+13	-11	227	15	249	8	271	1	292	18	314	11	336	4	357	21	-----	11 5*	34 12*		
+12	-12	228	12*	250	5*	22*	293	15*	315	8*	337	1*	358	18*	-----	12 4*	35 11*			
+11	-13	229	10*	251	3*	272	20*	294	13*	316	6*	339	16*	-----	13 4	36 11				
+10	-14	230	8	252	1	273	18	295	11	317	4	338	21	360	14	-----	14 3	37 10		
+9	-15	231	5*	222*	274	16	296	9	318	2	339	19	361	12	-----	15 2*	38 9*			
+8	-16	232	3*	253	20*	275	13*	297	6*	323	23*	340	16*	362	9	-----	16 2	39 8*		
+7	-17	233	1	254	18	276	11	298	4	319	21	341	14	363	7*	-----	17 1	40 8		
+6	-18	23	255	16	277	9	299	2	320	19	342	12	364	5	-----	18 0*	41 7*			
+5	-19	234	20*	256	13*	278	6*	300	21*	321	16*	343	9*	365	2*	-----	23*	42 6*		
+4	-20	235	18*	257	11*	279	4*	301	21*	322	14*	344	7*	366	0*	-----	19 23	43 6		
+3	-21	236	16	258	9	280	2	301	19	323	12	345	5	367	22	-----	20 22	44 5		
+2	-22	237	13*	259	6*	281	23*	302	16*	324	10	346	3	367	20	-----	21 21*	45 4*		
+1	-23	238	11*	260	4*	281	21*	303	14*	325	7*	347	0*	368	17*	-----	22 21	46 4		
Difference	Constituent MN																			
Hour	d.	h.	d.	h.	d.	h.	d.	h.	d.	h.	d.	h.	d.	h.						
0	47	3	70	10	93	17	117	0	140	7	163	14	186	21	210	4	233	11	256 18	
+23	-1	48	2*	71	9*	94	16*	23	141	6	164	13	187	20	211	3	234	10	257 17	
+22	-2	49	1*	72	8*	95	15*	118	22*	142	5*	165	12*	188	19*	212	2*	235	9*	258 16*
+21	-3	50	1	73	8	96	15	119	22	143	5	166	12	189	18*	213	1*	236	8*	259 15*
+20	-4	51	0	74	7	97	14	120	21	144	4	167	11	190	18	214	1	237	8	260 15
+19	-5	23*	75	6*	98	13*	121	20*	145	3*	168	10*	191	17*	215	0*	238	7*	261 14	
+18	-6	52	23	76	6	99	12*	122	19*	146	2*	169	9*	192	16*	23*	2	239	6*	262 13*
+17	-7	53	22	77	5	100	12	123	19	147	2	170	9	193	16	216	23	240	6	263 13
+16	-8	54	21*	78	4*	101	11*	124	18*	148	1	171	8	194	15	217	22	241	5	264 12
+15	-9	55	20*	79	3*	102	10*	125	17*	149	0*	172	7*	195	14*	218	21*	242	4*	265 11*
+14	-10	56	20	80	3	103	10	126	17	150	0	173	7	196	14	219	20*	243	3*	266 10*
+13	-11	57	19	81	2	104	9	127	16	23	174	6	197	13	220	20	244	3	267 10	
+12	-12	58	18*	82	1*	105	8*	128	15*	151	5*	175	2*	198	12*	221	19*	245	2*	268 9*
+11	-13	59	18	83	1	106	8	129	14*	152	21*	176	4*	199	11*	222	18*	246	1*	269 8*
+10	-14	60	17	84	0	107	7	130	14	153	21	177	4	200	11	223	18	247	1	270 8
+9	-15	61	16*	85	23*	108	6*	131	13*	154	20*	178	3	201	10	224	17	248	0	271 7
+8	-16	62	15*	85	22*	109	5*	132	12*	155	19*	179	2*	202	9*	225	16*	249	23*	272 6*
+7	-17	63	15	86	22	110	5	133	12	156	19	180	2	203	9	226	16	249	22*	273 5*
+6	-18	64	14*	87	21	111	4	134	11	157	18	181	1	204	8	227	15	250	22	274 5
+5	-19	65	13*	88	20*	112	3*	135	10*	158	17*	182	0*	205	7*	228	14*	251	21*	275 4*
+4	-20	66	13	89	20	113	3	136	10	159	16*	23*	206	6*	229	13*	252	20*	276 3*	
+3	-21	67	12	90	19	114	2	137	9	160	16	183	23	207	6	230	13	253	20	277 3
+2	-22	68	11*	91	18*	115	1*	138	8*	161	15*	184	22*	208	5*	231	12	254	19	278 2
+1	-23	69	10*	92	17	116	0*	139	7*	162	14*	185	21*	209	4*	232	11*	255	18*	279 1*

Table 31.—*For construction of primary stencils—Continued*

Difference	Constituent MN					Constituent M															
Hour	d.	h.	d.	h.	d.	d.	h.	d.	h.	d.	d.	h.	d.								
0	280	0*	303	7*	326	14*	349	21*	1	0	29	22*	59	11*	89	0	118	13	148	1*	
+23	-1	281	0	304	7	327	14	350	21	15*	31	4	60	17	90	5*	119	18*	149	7	
+22	-2	282	23*	305	6*	328	13*	351	20	2	21	32	10	61	22*	91	11	121	0	150	12*
+21	-3	282	22*	306	5*	329	12*	352	19*	4	2*	33	15*	63	4	92	17	122	5*	151	18
+20	-4	283	22	307	5	330	12	353	19	5	8	34	21	64	9*	93	22*	123	11	153	0
+19	-5	284	21	308	4	331	11	354	18	6	13*	36	2*	65	15	95	4	124	16*	154	5*
+18	-6	285	20*	309	3*	332	10*	355	17*	7	19	37	8	66	20*	96	9*	125	22	155	11
+17	-7	286	20	310	2*	333	9*	356	16*	9	0*	38	13*	68	2	97	15	127	3*	156	16*
+16	-8	287	19	311	2	334	9	357	16	10	6	39	19	69	7*	98	20*	128	9	157	22
+15	-9	288	18*	312	1*	335	8*	358	15*	11	12	41	0*	70	13	100	2	129	14*	159	3*
+14	-10	289	17	313	0*	336	7*	359	14*	12	17*	42	6	71	19	101	7*	130	20	160	9
+13	-11	290	17	314	0	337	7	360	14	13	23	43	11*	73	0*	102	13	132	2	161	14*
+12	-12	291	16	323	6	338	6	361	13	15	4*	44	17	74	6	103	18*	133	7*	162	20
+11	-13	292	15*	315	22*	339	5*	362	12*	16	10	45	22*	75	11*	105	0	134	13	164	1*
+10	-14	293	15	316	22	340	5	363	11*	17	15*	47	4	76	17	106	5*	135	18*	165	7
+9	-15	294	14	317	21	341	4	364	11	18	21	48	9*	77	22*	107	11	137	0	166	12*
+8	-16	295	13*	318	20*	342	3*	365	10*	20	2*	49	15	79	4	108	16*	138	5*	167	18
+7	-17	296	12*	319	19*	343	2*	366	9*	21	8	50	20*	80	9*	109	22	139	11	168	23*
+6	-18	297	12	320	19	344	2	367	9	22	13*	52	2*	81	15	111	3*	140	16*	170	5
+5	-19	298	11*	321	18	345	1	368	8	23	19	53	8	82	20*	112	9*	141	22	171	10*
+4	-20	299	10*	322	17*	346	0*	369	7*	25	0*	54	13*	84	2	113	15	143	3*	172	16*
+3	-21	300	10	323	17	347	0	370	7	26	6	55	19	85	7*	114	20*	144	9	173	22
+2	-22	301	9	324	16	23	-----	-----	27	11*	57	0*	86	13	116	2	145	14*	175	3*	
+1	-23	302	8	325	15*	348	22*	-----	28	17	58	6	87	18*	117	7*	146	20	176	9	
Difference	Constituent M					Constituent MK															
Hour	d.	h.	d.	h.	d.	d.	h.	d.	h.	d.	d.	h.	d.								
0	177	14*	207	3	236	16	266	4*	295	17*	325	6	354	19	1	0	46	5*	92	9*	
+23	-1	178	20	208	8*	237	21*	267	10	296	23	326	11*	356	0*	2	0	48	3*	94	7*
+22	-2	180	1*	209	14	239	3	268	15*	298	4*	327	17	357	6	3	22	50	2	96	6
+21	-3	181	7	210	19*	240	8*	269	21	299	10	328	22*	358	11*	5	20	52	0	98	4
+20	-4	182	12*	212	1	241	14	271	2*	300	15*	330	4	359	17	7	18*	53	22	100	2
+19	-5	183	18	213	7	242	19*	272	8	301	21	331	9*	360	22*	9	16*	55	20*	102	0*
+18	-6	184	23*	214	12*	244	1	273	14	303	2*	332	15	362	4	11	14*	57	18*	103	22*
+17	-7	186	5	215	18	245	6*	274	19*	304	8	333	21	363	9*	13	13	59	16*	105	20*
+16	-8	187	10*	216	23*	246	12	276	1	305	13*	335	2	364	15	15	11	61	15	107	18*
+15	-9	188	16	218	5	247	17*	277	6*	306	19	336	8	365	20*	17	9	63	13	109	17
+14	-10	189	21*	219	10*	248	23	278	12	308	0*	337	13*	367	2	19	7*	63	11	111	15
+13	-11	191	3	220	16	250	4*	279	17*	309	6	338	19	368	7*	21	5*	67	9*	113	13
+12	-12	192	9	221	21*	251	10	280	23	310	11*	340	0*	369	13	23	3*	69	7*	115	11*
+11	-13	193	14*	223	3	252	16	282	4*	311	17	341	6	370	18*	25	2	71	5*	117	9*
+10	-14	194	20	224	8*	253	21*	283	10	312	23	342	11*	-----	-----	27	0	73	4	119	7*
+9	-15	196	1*	225	14	255	3	284	15*	314	4*	343	17	-----	-----	28	22	75	2	121	6
+8	-16	197	7	226	19*	256	8*	285	21	315	10	344	22*	-----	-----	30	20*	77	0	123	4
+7	-17	198	12*	228	1	257	14	287	2*	316	15*	346	4	-----	-----	32	18*	78	22*	125	2
+6	-18	199	18	229	6*	258	19*	288	8	317	21	347	9*	-----	-----	34	16*	80	20*	127	0*
+5	-19	200	23*	230	12	260	1	289	13*	319	2*	348	15	-----	-----	36	14*	82	18*	128	22*
+4	-20	202	5	231	17*	261	6*	290	19	320	8	349	20*	-----	-----	38	13	84	17	130	20*
+3	-21	203	10*	232	23*	262	12	292	0*	321	13*	351	2	-----	-----	40	11	86	15	132	19
+2	-22	204	16	234	5	263	17*	293	6*	322	19	352	7*	-----	-----	42	9	88	13	134	17
+1	-23	205	21*	235	10*	264	23	294	12	324	0*	353	13*	-----	-----	44	7*	90	11*	136	15

Table 31.—*For construction of primary stencils—Continued*

Difference	Constituent MK								Constituent λ								
Hour	d.	h.	d.	h.	d.	h.	d.	h.	d.	h.	d.	h.	d.	h.	d.	h.	
0	138	13*	184	17	230	21	277	1	323	5	369	9	1	0	55	0	
+23	-1	140	11*	186	15*	232	19*	278	23	325	3	371	7	2	4*	57	7
+22	-2	142	9*	188	13*	234	17*	280	21*	327	1	-----	4	11*	59	14	
+21	-3	144	8	190	11*	236	15*	282	19*	328	23	-----	6	18*	61	21	
+20	-4	146	6	192	10	238	14	284	21*	330	21	-----	9	1*	64	4*	
+19	-5	148	4	194	8	240	12	286	16	332	19*	-----	11	8*	66	11*	
+18	-6	150	2*	196	6	242	10	288	14	334	18	-----	13	16	68	18*	
+17	-7	152	0*	198	4*	244	8	290	12	336	16	-----	15	23	71	1*	
+16	-8	153	22*	200	2*	246	6*	292	10*	338	14	-----	18	6	73	8*	
+15	-9	155	21	202	0*	248	4*	294	8*	340	12*	-----	20	13	75	16	
+14	-10	157	19	203	23	250	2*	296	6*	342	10*	-----	22	20*	77	23	
+13	-11	159	17	205	21	252	1	298	5	344	8*	-----	25	3*	80	6	
+12	-12	161	15*	207	19	253	23	300	3	346	7	-----	27	10*	82	13	
+11	-13	163	13*	209	17*	255	21	302	1	348	5	-----	28	17*	84	20	
+10	-14	165	11*	211	15*	257	19*	303	23*	350	3	-----	32	0*	87	3*	
+9	-15	167	10	213	13*	259	17*	305	21*	352	1*	-----	34	8	89	10*	
+8	-16	169	8	215	12	261	15*	307	19*	353	23*	-----	36	15	91	17*	
+7	-17	171	6	217	10	263	14	309	18	355	21*	-----	38	22	94	0*	
+6	-18	173	4*	219	8	265	12	311	16	357	20	-----	41	5	96	8	
+5	-19	175	2*	221	6*	267	10	313	14	359	18	-----	43	12*	98	15	
+4	-20	177	0*	223	4*	269	8*	315	12	361	16	-----	45	19*	100	22	
+3	-21	178	22*	225	2*	271	6*	317	10*	363	14*	-----	48	2*	103	5	
+2	-22	180	21	227	1	273	4*	319	8*	365	12*	-----	50	9*	105	12	
+1	-23	182	19	228	23	275	3	321	6*	367	10*	-----	52	16*	107	19*	
Difference	Constituent λ								Constituent MS								
Hour	d.	h.	d.	h.	d.	h.	d.	h.	d.	h.	d.	h.	d.	h.	d.	h.	
0	220	7*	275	10*	330	13	1	0	58	20*	117	22	176	23*	236	1	
+23	-1	222	15	277	17*	332	20	2	6*	61	7*	120	9	179	10*	238	12
+22	-2	224	22	280	0*	335	3	4	17*	63	19	122	20*	181	21*	240	23
+21	-3	227	5	282	7*	337	10	7	4*	66	6	125	7*	184	9	243	10*
+20	-4	229	12	284	14*	339	17*	9	15*	68	17	127	18*	186	20	245	21*
+19	-5	231	19	286	22	342	0*	12	2*	71	4	130	5*	189	7	248	8*
+18	-6	234	2*	289	5	344	7*	14	13*	73	15	132	16*	191	18	250	19*
+17	-7	236	9*	291	12	346	14*	17	0*	76	2	135	3*	194	5	253	6*
+16	-8	238	16*	293	19	348	22	19	11*	78	13	137	14*	196	16	255	17*
+15	-9	240	23*	296	2*	351	5	21	23	81	0	149	-1*	199	3	258	4*
+14	-10	243	7	298	9*	353	12	24	10	83	11*	142	13	201	14	260	15*
+13	-11	245	14	300	16*	355	19	26	21	85	22*	145	0	204	1*	263	3
+12	-12	247	21	302	23*	358	2	29	8	88	9*	147	11	206	12*	265	14
+11	-13	250	4	305	6*	360	9*	31	19	90	20*	149	22	208	23*	268	1
+10	-14	252	11	307	14	362	16*	34	6	93	7*	152	9	211	10*	270	12
+9	-15	254	18*	309	21	364	23*	36	17	95	18*	154	20	213	21*	272	23
+8	-16	257	1*	312	4	367	6*	39	4	98	5*	157	7	216	8*	275	10
+7	-17	259	8*	314	11	369	14	41	15*	100	16*	159	18	218	19*	277	21
+6	-18	261	15*	316	18*	371	21	44	2*	103	4	162	5*	221	6*	280	8
+5	-19	263	22*	319	1*	-----	-----	46	13*	105	15	164	16*	223	18	282	19*
+4	-20	266	6	321	8*	-----	-----	49	0*	108	2	167	3*	226	5	285	6*
+3	-21	268	13	323	15*	-----	-----	51	11*	110	13	169	14*	228	16	287	17*
+2	-22	270	20	325	22*	-----	-----	53	22*	113	0	172	1*	231	3	290	4*
+1	-23	273	3	328	6	-----	-----	56	9*	115	11	174	12*	233	14	292	15*

Table 31.—*For construction of primary stencils—Continued*

Difference		Constituent L							Constituent P		Constituent T
Hour		d. h.	d. h.	d. h.	d. h.	d. h.	d. h.	d. h.	d. h.	d. h.	d. h.
+23	-1	1 0	63 8	126 23	190 14	254 5	317 20	1 0	358 16	1 0	1 0
+22	-2	2 8*	65 23*	129 14*	193 5*	256 20*	320 11*	8 15*	373 21	18 6	46 16*
+21	-3	5 0	68 15	132 6	195 21	259 12	323 3	23 20*		77 3	
+20	-4	7 16	71 7	134 22	198 12*	262 3*	325 18*	39 2		107 13*	
+19	-5	10 7*	73 22*	137 13*	201 4*	264 19*	328 10*	54 7		138 0	
+18	-6	15 14*	79 5*	142 20*	206 11*	270 2*	333 17*	84 17*		168 10*	
+17	-7	18 6*	81 21*	145 12*	209 3	272 18	336 9	99 23		198 21	
+16	-8	20 22	84 13	148 4	211 19	275 10	339 1	115 4		229 7*	
+15	-9	23 13*	87 4*	150 19*	214 10*	278 1*	341 16*	130 9*		259 18	
+14	-10	26 5	89 20	153 11	217 2	280 17	344 8	145 14*		290 4*	
+13	-11	28 21	92 12	156 2*	219 17*	283 8*	346 23*	160 20		320 15	
+12	-12	31 12*	95 3*	158 18*	222 9*	286 0*	349 15*	176 1		351 1*	
+11	-13	34 4	97 19	161 10	225 1	288 16	352 7	191 6*		381 12	
+10	-14	36 19*	100 10*	164 1*	227 16*	291 7*	354 22*	206 11*			
+9	-15	39 11*	103 2*	166 17	230 8	293 23	357 14	221 17			
+8	-16	42 3	105 18	169 9	233 0	296 15	360 6	236 22			
+7	-17	44 18*	108 9*	172 0*	235 15*	299 6*	362 21*	252 3*			
+6	-18	47 10	111 1	174 16	238 7	301 22	365 13	267 8*			
+5	-19	50 2	113 17	177 7*	240 22*	304 13*	368 4*	282 13*			
+4	-20	52 17*	116 8*	179 23*	243 14*	307 5*	370 20*	297 19			
+3	-21	55 9	119 0	182 15	246 6	309 21			313 0		
+2	-22	58 0*	121 15*	185 6*	248 21*	312 12*			328 5*		
+1	-23	60 16*	124 7*	187 22	251 13	315 4			343 10*		
Difference		Constituent R	Constituent K	Constituent 2SM							
Hour		d. h.	d. h.	d. h.	d. h.	d. h.	d. h.	d. h.	d. h.	d. h.	d. h.
0		1 0	1 0	358 16	1 0	29 22*	59 11*	89 0	118 13	148 1*	177 14*
+1	-23	16 6	8 15*	373 21	15*	31 4	60 17	90 5*	119 18*	149 7	178 20
+2	-22	46 16*	23 20*		2 21	32 10	61 22*	91 11	121 0	150 12*	180 1*
+3	-21	77 3	39 2		4 2*	33 15*	63 4	92 17	122 5*	151 18	181 7
+4	-20	107 13*	54 7		5 8	34 21	64 9*	93 22*	123 11	153 0	182 12*
+5	-19	138 0	69 12*		6 13*	36 2*	65 15	95 4	124 16*	154 5*	183 18
+6	-18	168 10*	84 17*		7 19	37 8	66 20*	96 9*	125 22	155 11	184 23*
+7	-17	198 21	99 23		9 0*	38 13*	68 2	97 15	127 3*	156 16*	186 5
+8	-16	229 7*	115 4		10 6	39 19	69 7*	98 20*	128 9	157 22	187 10*
+9	-15	259 18	130 9*		11 12	41 0*	70 13	100 2	129 14*	159 3*	188 16
+10	-14	290 4*	145 14*		12 17*	42 6	71 19	101 7*	130 20	160 9	189 21*
+11	-13	320 15	160 20		13 23	43 11*	73 0*	102 13	132 2	161 14*	191 3
+12	-12	351 1*	176 1		15 4*	44 17	74 6	103 18*	133 7*	162 20	192 9
+13	-11	381 12	191 6*		16 10	45 22*	75 11*	105 0	134 13	164 1*	193 14*
+14	-10		206 11*		17 15*	47 4	76 17	106 5*	135 18*	165 7	194 20
+15	-9		221 17		18 21	48 9*	77 22*	107 11	137 0	166 12*	196 1*
+16	-8		236 22		20 2*	49 15	79 4	108 16*	138 5*	167 18	197 7
+17	-7		252 3*		21 8	50 20*	80 9*	109 22	139 11	168 23*	198 12*
+18	-6		267 8*		22 13*	52 2*	81 15	111 3*	140 16*	170 5	199 18
+19	-5		282 13*		23 19	53 8	82 20*	112 9*	141 22	171 10*	200 23*
+20	-4		297 19		25 0*	54 13*	84 2	113 15	143 3*	172 16*	202 5
+21	-3		313 0		26 6	55 19	85 7*	114 20*	144 9	173 22	203 10*
+22	-2		328 5*		27 11*	57 0*	86 13	116 2	145 14*	175 3*	204 16
+23	-1		343 10*		28 17	58 6	87 18*	117 7*	146 20	176 9	205 21*

Table 31.—*For construction of primary stencils—Continued*

Difference	Constituent 2SM							Constituent J							
	Hour	d.	h.	d.	h.	d.	h.	d.	h.	d.	h.	d.	h.	d.	h.
+1 -23	207 3	236 16	266 4*	295 17*	325 6	354 19	1 0	26 3	51 18	77 8*					
+1 -22	208 8*	237 21*	267 10	296 23	326 11*	356 0*	13*	27 4*	52 19*	78 10*					
+3 -21	209 14	239 3	268 15*	298 4*	327 17	357 6	2 15	28 6	53 21	79 12					
+4 -20	210 19*	240 8*	269 21	299 10	328 22*	358 11*	3 17	29 7*	54 22*	80 13*					
+5 -19	212 1	241 14	271 2*	300 15*	330 4	359 17	4 18*	30 9*	56 0*	81 15					
+6 -18	214 12*	244 1	273 14	303 2*	332 15	362 4	6 21*	32 12*	58 3*	83 18*					
+7 -17	215 18	245 6*	274 19*	304 8	333 21	363 9*	7 23*	33 14	59 5	84 20					
+8 -16	216 23*	246 12	276 1	305 13*	335 2*	364 15	9 1	34 16	60 6*	85 21*					
+9 -15	218 5	247 17*	277 6*	306 19	336 8	365 20*	10 2*	35 17*	61 8*	86 23*					
+10 -14	219 10*	248 23	278 12	308 0*	337 13*	367 2	11 4	36 19	62 10	88 1					
+11 -13	220 16	250 4*	279 17*	309 6	338 19	368 7*	12 6	37 20*	63 11*	89 2*					
+12 -12	221 21*	251 10	280 23	310 11*	340 0*	369 13	13 7*	38 22*	64 13	90 4					
+13 -11	223 3	252 16	282 4*	311 17	341 6	370 18*	14 9	40 0	65 15	91 6					
+14 -10	224 8*	253 21*	283 10	312 23	342 11*	-----	15 10*	41 1*	66 16*	92 7*					
+15 -9	225 14	255 3	284 15*	314 4*	343 17	-----	16 12*	42 3	67 18	93 9					
+16 -8	226 19*	256 8*	285 21	315 10	344 22*	-----	17 14	43 5	68 19*	94 10*					
+17 -7	228 1	257 14	287 2*	316 15*	346 4	-----	18 15*	44 6*	69 21*	95 12*					
+18 -6	229 6*	258 19*	288 8	317 21	347 9*	-----	19 17	45 8	70 23	96 14					
+19 -5	230 12	260 1	289 13*	319 2*	348 15	-----	20 18*	46 9*	72 0*	97 15*					
+20 -4	231 17*	261 6*	290 19	320 8	349 20*	-----	21 20*	47 11*	73 2	98 17					
+21 -3	232 23*	262 12	292 0*	321 13*	351 2	-----	22 22	48 13	74 4	99 18*					
+22 -2	234 5	263 17*	293 6*	322 19	352 7*	-----	23 23*	49 14*	75 5*	100 20*					
+23 -1	235 10*	264 23	294 12	324 0*	353 13*	-----	25 1	50 16	76 7	101 22					
Difference	Constituent J														
Hour	d.	h.	d.	h.	d.	h.	d.	h.	d.	h.	d.	h.	d.	h.	
0	102 23*	128 14*	154 5*	179 20*	205 11*	231 2	256 17	282 8	307 23	333 14					
+1 -23	104 1	129 16	155 7	180 22	206 13	232 4	257 18*	283 9*	309 0*	334 15*					
+2 -22	105 3	130 18	156 8*	181 23*	207 14*	233 5*	258 20*	284 11*	310 2	335 17					
+3 -21	106 4*	131 19*	157 10*	183 1	208 16	234 7	259 22	285 13	311 4	336 18*					
+4 -20	107 6	132 21	158 12	184 3	209 18	235 8*	260 23*	286 14*	312 5*	337 20*					
+5 -19	108 7*	133 22*	159 13*	185 4*	210 19*	236 10*	262 1	287 16	313 7	338 22					
+6 -18	109 9*	135 0*	160 15	186 6	211 21	237 12	263 3	288 18	314 8*	339 23*					
+7 -17	110 11	136 2	161 17	187 7*	212 22*	238 13*	264 4*	289 19*	315 10*	341 1					
+8 -16	111 12*	137 3*	162 18*	188 9*	214 0*	239 15	265 6	290 21	316 12	342 3					
+9 -15	112 14	138 5	163 20	189 11	215 2	240 17	266 7*	291 22*	317 13*	343 4*					
+10 -14	113 16	139 6*	164 21*	190 12*	216 3*	241 18*	267 9*	293 0*	318 15	344 6					
+11 -13	114 17*	140 8*	165 23*	191 14	217 5	242 20	268 11	294 2	319 17	345 7*					
+12 -12	115 19	141 10	167 1	192 16	218 6*	243 21*	269 12*	295 3*	320 18*	346 9*					
+13 -11	116 20*	142 11*	168 2*	193 17*	219 8*	244 23*	270 14	296 5	321 20	347 11					
+14 -10	117 22*	143 13	169 4	194 19	220 10	246 1	271 16	297 6*	322 21*	348 12*					
+15 -9	119 0	144 15	170 6	195 20*	221 11*	247 2*	272 17*	298 8*	323 23*	349 14					
+16 -8	120 1*	145 16*	171 7*	196 22*	222 13	248 4	273 19	299 10	325 1	350 16					
+17 -7	121 3	146 18	172 9	198 0	223 15	249 6	274 20*	3 0 11*	326 2*	351 17*					
+18 -6	122 5	147 19*	173 10*	199 1*	224 16*	250 7*	275 22*	301 13	327 4	352 19					
+19 -5	123 6*	148 21*	174 12*	200 3	225 18	251 9	277 0	302 15	328 6	353 20*					
+20 -4	124 8	149 23	175 14	201 5	226 19*	252 10*	278 1*	303 16*	329 7*	354 22*					
+21 -3	125 9*	151 0*	176 15*	202 6*	227 21*	253 12*	279 3	304 18	330 9	356 0					
+22 -2	126 11*	152 2	177 17	203 8	228 23	254 14	280 5	305 19*	331 10*	357 1*					
+23 -1	127 13	153 4	178 18*	204 9*	230 0*	255 15*	281 6*	306 21*	332 12*	358 3					

Table 31.—*For construction of primary stencils—Continued*

Difference	Con. J	Constituent OO												
<i>Hour</i>		<i>d.</i>	<i>h.</i>	<i>d.</i>	<i>h.</i>	<i>d.</i>	<i>h.</i>	<i>d.</i>	<i>h.</i>	<i>d.</i>	<i>h.</i>	<i>d.</i>	<i>h.</i>	
0	359 5	1 0	13 22	27 2	40 6*	53 10*	66 14*	79 18*	92 22*	106 2*				
+1	-23	360 6*	7*	14 11*	15*	19*	23*	67 3*	80 7*	93 11*	15*			
+2	-22	361 8	20*	15 0*	28 4*	41 8*	54 12*	16*	20*	94 1	107 5			
+3	-21	362 9*	2 9*	13*	17*	22	55 2	68 6	81 10	14	18			
+4	-20	363 11*	23	16 3	29 7	42 11	15	19	23	95 3	108 7			
+5	-19	364 13	3 12	16	20	43 0	56 4	69 8	82 12	16*	20*			
+6	-18	365 14*	4 1	17 5	30 9*	13*	17*	21*	83 1*	96 5*	109 9*			
+7	-17	366 16	14*	18*	22*	44 2*	57 6*	70 10*	14*	18*	22*			
+8	-16	367 18	5 3*	18 7*	31 11*	15*	19*	23*	84 3*	97 8	110 12			
+9	-15	368 19*	16*	20*	32 1	45 5	58 9	71 13	17	21	111 1			
+10	-14	369 21	6 6	19 10	14	18	22	72 2	85 6	98 10	14			
+11	-13	370 22*	19	23	33 3	46 7	59 11	15	19	23*	112 3*			
+12	-12	-----	7 8	20 12	16*	20*	60 0*	73 4*	86 8*	99 12*	16*			
+13	-11	-----	21*	21 1*	34 5*	47 9*	13*	17*	21*	100 1*	113 5*			
+14	-10	-----	8 10*	14*	18*	22*	61 2*	74 6*	87 11	15	19			
+15	-9	-----	23*	22 3*	35 8	48 12	16	20	88 0	101 4	114 8			
+16	-8	-----	9 13	17	21	49 1	62 5	75 9	13	17	21			
+17	-7	-----	10 2	23 6	36 10	14	18	22	89 2*	102 6*	115 10*			
+18	-6	-----	15	19	23*	50 3*	63 7*	76 11*	15*	19*	23*			
+19	-5	-----	11 4*	24 8*	37 12*	16*	20*	77 0*	90 4*	103 8*	116 12*			
+20	-4	-----	17*	21*	38 1*	51 5*	64 9*	13*	18	22	117 2			
+21	-3	-----	12 6*	25 10*	15	19	23	78 3	91 7	104 11	115			
+22	-2	-----	20	26 0	39 4	52 8	65 12	16	20	105 0	118 4			
+23	-1	-----	13 9	13	17	21	66 1	79 5	92 9*	13*	17*			

Difference		Constituent OO												
<i>Hour</i>		<i>d.</i>	<i>h.</i>	<i>d.</i>	<i>h.</i>	<i>d.</i>	<i>h.</i>	<i>d.</i>	<i>h.</i>	<i>d.</i>	<i>h.</i>	<i>d.</i>	<i>h.</i>	
0	119 6*	132 10*	145 14*	158 18*	171 22*	185 2*	198 6*	211 11	224 15	237 19				
+1	-23	119 19*	23*	146 4	159 8	172 12	16	20	212 0	225 4	238 8			
+2	-22	120 9	133 13	17	21	173 1	186 5	199 9	13	17	21			
+3	-21	121 22	134 2	147 6	160 10	14	18	22	213 2*	226 6*	239 10*			
+4	-20	121 11	15	19*	23*	174 3*	187 7*	200 11*	15*	19*	23*			
+5	-19	122 0*	135 4*	148 8*	161 12*	16*	20*	201 0*	214 4*	227 8*	240 12*			
+6	-18	13*	17*	21*	162 1*	175 5*	188 9*	14	18	22	241 2			
+7	-17	123 2*	136 6*	149 11	15	19	23	202 3	215 7	228 11	15			
+8	-16	16	20	150 0	163 4	176 8	189 12	16	20	229 0	242 4			
+9	-15	124 5	137 9	13	17	21	190 1	203 5*	216 9*	13*	17*			
+10	-14	18	22	151 2*	164 6*	177 10*	14*	18*	22*	230 2*	243 6*			
+11	-13	125 7*	138 11*	15*	19*	23*	191 3*	204 7*	217 11*	15*	19*			
+12	-12	20*	139 0*	152 4*	165 8*	178 12*	16*	21	218 1	231 5	244 9			
+13	-11	126 9*	13*	18	22	179 2	192 6	205 10	14	18	22			
+14	-10	23	140 3	153 7	166 11	15	19	23	219 3	232 7	245 11			
+15	-9	127 12	16	20	167 0	180 4	193 8	206 12*	16*	20*	246 0*			
+16	-8	128 1	141 5	154 9*	13*	17*	21*	207 1*	220 5*	233 9*	13*			
+17	-7	14*	18*	22*	168 2*	181 6*	194 10*	14*	18*	22*	247 2*			
+18	-6	129 3*	142 7*	155 11*	15*	19*	23*	208 4	221 8	234 12	16			
+19	-5	16*	20*	156 1	169 5	182 9	195 13	17	21	235 1	248 5			
+20	-4	130 6	143 10	14	18	22	196 2	209 6	222 10	14	18			
+21	-3	19	23	157 3	170 7	183 11	15	19*	23*	236 3*	249 7*			
+22	-2	131 8	144 12*	16*	20*	184 0*	197 4*	210 8*	223 12*	16*	20*			
+23	-1	21*	145 1*	158 5*	171 9*	13*	17*	21*	224 1*	237 5*	250 9*			

Table 31.—*For construction of primary stencils—Continued*

Difference	Constituent OO												
Hour	d.	h.	d.	h.	d.	h.	d.	h.	d.	h.	d.	h.	
0	250	23	264	3	277	7	290	11	303	15	316	19	
+1	-23	251	12	16	20	291	0	304	4	317	8*	330	12*
+2	-22	252	1	265	5*	278	9*	292	13*	305	17*	311	21*
+3	-21		14*	18*		22*	2*	292	6*	318	10*	331	14*
+4	-20	253	3*	266	7*	279	11*	292	15*	305	19*	319	0
+5	-19		16*		21	280	1	293	5	306	9	312	13
+6	-18	254	6	267	10		14		18	22	320	2	
+7	-17	19		23	281	3	294	7	307	11	315	15*	
+8	-16	255	8	268	12*		16*	20*	208	0*	321	4*	
+9	-15		21*	269	1*	282	5*	295	9*	313	13*	334	17*
+10	-14	256	10*	14*		18*	22*	299	2*	322	7	335	11
+11	-13		23*	270	4	283	8	296	12	316	20	336	0
+12	-12	257	13		17		21	297	1	310	5	323	9
+13	-11	258	2	271	6	284	10	298	14	311	18	327	22*
+14	-10		15*	19*		23*	298	3*	311	7*	324	11*	
+15	-9	259	4*	272	8*	285	12*	298	16*	316	20*	325	0*
+16	-8		17*	21*		286	1*	299	5*	312	9*	338	4*
+17	-7	260	7	273	11		15		19	23	326	3	
+18	-6	20	274	0	287	4	300	8	313	12	316	16	
+19	-5	261	9	13		17		21	314	1	327	5*	
+20	-4		22*	275	2*	288	6*	301	10*	314	14*	340	9*
+21	-3	262	11*	15*		19*		23*	315	3*	328	7*	
+22	-2	263	0*	276	4*	289	8*	302	12*	317	21	341	11*
+23	-1		14		18		22	303	2	316	6	329	10

Table 32.—*Divisors for primary stencil sums*

## CONSTITUENT J

Series	29	58	87	105	134	163	192	221	250	279	297	326	355	369
<i>Hour</i>														
0-----	30	59	87	106	134	164	192	221	250	279	298	326	355	370
1-----	31	59	89	106	135	164	193	222	250	280	298	327	356	369
2-----	28	58	86	104	134	162	192	220	250	278	296	326	354	369
3-----	30	59	88	106	135	165	192	222	251	280	299	326	356	370
4-----	29	59	88	104	135	163	193	222	250	280	297	327	355	369
5-----	28	59	87	105	134	163	193	221	251	278	297	326	355	370
6-----	30	57	88	106	134	165	192	222	250	280	298	326	356	369
7-----	28	58	87	104	134	163	193	221	250	279	297	327	354	369
8-----	29	58	88	106	134	164	193	222	251	279	298	326	356	371
9-----	29	57	87	105	134	163	192	222	250	280	297	326	355	369
10-----	28	58	86	104	134	162	193	220	250	278	297	326	354	368
11-----	30	59	88	107	134	164	193	223	251	280	299	327	357	370
12-----	29	57	87	104	134	162	191	221	250	279	296	326	354	368
13-----	28	58	85	104	133	162	191	220	250	278	297	325	354	368
14-----	30	58	88	106	134	164	192	223	250	280	297	327	356	369
15-----	29	58	87	105	135	162	192	220	251	279	296	327	355	369
16-----	28	58	86	105	133	163	191	220	250	279	297	325	355	369
17-----	30	57	87	105	134	163	192	221	250	280	296	326	355	368
18-----	28	58	86	104	134	162	192	220	250	278	296	325	355	369
19-----	29	58	87	106	133	163	191	221	249	280	297	325	356	369
20-----	29	57	87	104	134	162	191	220	249	279	296	326	354	368
21-----	28	58	85	104	133	162	191	219	249	277	296	325	354	369
22-----	30	58	88	106	134	164	192	222	249	279	298	326	356	369
23-----	28	57	86	104	134	161	191	219	249	277	295	325	353	368

## CONSTITUENT K

Series	14	29	58	87	105	134	163	192	221	250	279	297	326	355	369
<i>Hour</i>															
0-----	15	30	59	88	106	135	164	193	221	250	279	297	326	355	369
1-----	14	30	59	88	106	135	164	193	222	251	279	297	326	355	369
2-----	14	29	59	88	106	135	164	193	222	251	280	298	327	355	369
3-----	14	29	59	88	106	135	164	193	222	251	280	298	327	356	370
4-----	14	29	57	87	105	134	163	192	221	250	279	297	326	355	369
5-----	14	29	58	88	105	134	163	192	221	250	279	297	326	355	369
6-----	14	29	58	87	106	135	163	192	221	250	279	297	326	355	369
7-----	14	29	58	87	105	135	164	193	221	250	279	296	325	354	368
8-----	14	29	58	87	105	135	164	193	222	251	280	298	327	355	369
9-----	14	29	58	87	105	134	164	193	222	251	280	298	327	356	370
10-----	14	29	57	86	104	133	163	192	221	250	279	297	326	355	369
11-----	14	29	58	87	105	133	162	192	221	250	279	297	326	355	369
12-----	14	29	58	87	105	134	163	192	221	250	279	297	326	355	369
13-----	14	29	58	87	105	134	163	192	222	250	279	297	326	355	369
14-----	14	29	58	87	105	134	163	192	222	251	280	297	326	355	369
15-----	13	28	57	86	104	133	162	191	220	250	279	297	326	355	368
16-----	14	29	58	86	104	133	162	191	220	249	279	297	326	355	369
17-----	14	29	58	87	105	133	162	191	220	249	279	297	326	355	369
18-----	14	29	58	87	105	134	163	191	220	249	278	297	326	355	369
19-----	14	29	58	87	105	134	163	192	221	250	278	297	326	355	369
20-----	14	29	58	87	105	134	163	192	221	250	279	297	326	355	369
21-----	14	28	57	86	104	133	162	191	220	249	278	296	325	355	369
22-----	14	29	58	86	104	133	162	191	220	249	278	296	325	355	369
23-----	14	29	58	87	105	134	162	191	220	249	278	296	325	354	369

Table 32.—*Divisors for primary stencil sums—Continued*

## CONSTITUENT L

Series	29	58	87	105	134	163	192	221	250	279	297	326	355	369
<i>Hour</i>														
0	29	59	87	105	133	163	191	221	250	279	297	326	355	369
1	29	59	87	106	134	164	192	222	251	279	297	326	355	369
2	29	58	87	106	134	163	192	221	250	280	298	326	356	370
3	30	58	87	105	134	163	192	221	250	279	298	326	356	370
4	30	58	88	106	135	164	192	222	250	279	297	326	355	370
5	29	58	88	106	134	164	192	222	250	280	298	327	356	369
6	29	57	86	105	133	163	191	221	249	279	297	325	355	368
7	30	59	88	106	135	164	193	222	250	279	298	326	356	369
8	30	58	88	105	135	164	193	222	251	280	298	327	357	370
9	29	57	87	104	133	163	191	221	250	279	296	326	355	369
10	30	58	87	105	134	164	192	221	249	279	296	326	354	368
11	29	58	87	105	134	162	192	222	250	280	297	326	355	369
12	29	58	87	104	134	162	192	221	250	279	297	326	355	369
13	29	58	88	105	135	163	192	220	250	279	296	326	354	368
14	29	58	88	105	134	163	193	221	250	280	297	327	355	370
15	28	58	86	105	134	163	192	221	250	279	297	327	355	369
16	28	58	86	104	134	162	191	220	249	278	296	325	353	367
17	28	57	86	104	134	162	192	220	250	278	297	326	355	369
18	29	58	87	105	134	162	192	220	250	278	296	326	355	369
19	29	58	87	105	135	163	192	221	250	279	297	326	354	369
20	28	58	86	105	134	163	192	221	250	279	297	327	355	369
21	28	58	86	104	132	162	191	219	249	277	296	324	354	368
22	29	58	87	105	134	163	193	221	251	279	297	325	355	369
23	29	58	87	105	134	163	193	221	251	279	298	326	355	370

## CONSTITUENT M

Series	15	29	58	87	105	134	163	192	221	250	279	297	326	355	369
<i>Hour</i>															
0	15	29	59	87	105	135	164	192	222	250	279	297	325	355	369
1	15	29	57	87	105	134	163	192	221	250	279	296	326	354	369
2	15	28	58	86	105	134	162	192	221	250	279	296	325	354	369
3	16	29	59	88	107	135	165	193	222	251	281	299	328	357	371
4	16	30	58	87	106	135	164	193	222	251	280	297	326	355	370
5	15	28	57	86	104	134	163	192	221	250	278	296	325	354	368
6	15	29	58	87	106	134	163	192	222	250	280	297	326	355	369
7	16	29	58	87	105	134	163	192	221	250	279	296	326	354	369
8	16	29	59	87	106	135	164	193	221	251	280	298	326	355	370
9	15	29	58	87	106	135	165	193	223	251	280	298	327	357	371
10	15	29	57	87	105	134	163	192	221	250	279	296	326	354	368
11	15	28	57	86	104	133	162	192	221	250	278	296	325	354	369
12	15	29	58	87	105	133	162	191	220	250	280	297	326	355	368
13	15	30	59	88	105	134	163	192	221	250	279	298	327	355	369
14	15	29	58	87	105	134	163	192	220	250	278	297	326	356	369
15	14	29	58	87	104	134	163	192	222	250	279	298	326	356	369
16	15	29	57	87	104	133	162	191	220	249	278	296	326	354	368
17	15	29	59	87	105	134	162	192	220	250	279	298	326	355	369
18	14	29	58	87	105	133	163	191	220	249	278	297	326	355	368
19	15	30	58	88	105	135	163	192	221	250	279	297	326	356	369
20	14	28	57	86	103	133	162	191	220	249	277	296	325	354	368
21	14	29	58	87	105	133	162	192	221	250	280	298	327	356	369
22	15	30	59	88	105	134	163	192	221	249	279	298	327	355	369
23	15	29	58	87	105	134	163	192	220	250	278	296	325	355	369

Table 32.—*Divisors for primary stencil sums—Continued*

## CONSTITUENT N

Series	15	29	58	87	105	134	163	192	221	250	279	297	326	355	369
<i>Hour</i>															
0-----	16	29	58	87	105	134	163	191	220	250	279	297	327	356	370
1-----	16	29	58	88	106	135	165	194	223	252	281	299	327	357	370
2-----	15	29	57	87	105	133	162	191	220	248	278	296	324	354	367
3-----	16	30	58	88	106	134	163	192	221	249	279	297	326	355	370
4-----	16	30	58	87	105	135	164	193	223	252	282	299	328	357	371
5-----	15	30	59	88	106	134	164	192	222	250	279	297	326	355	369
6-----	15	29	58	87	105	133	163	191	221	249	278	296	324	354	367
7-----	15	29	58	87	105	133	163	191	220	250	279	298	326	357	370
8-----	14	29	58	88	107	135	164	194	223	251	281	299	327	356	370
9-----	15	30	58	88	105	134	163	192	221	249	279	297	325	354	368
10-----	15	30	58	88	105	134	163	191	221	249	279	297	326	356	370
11-----	15	30	58	86	106	135	165	193	224	252	281	299	328	357	371
12-----	15	28	59	87	106	134	164	192	220	250	278	297	325	355	368
13-----	15	28	58	86	104	133	161	191	219	249	277	295	324	354	368
14-----	14	28	57	86	104	133	161	191	220	250	279	297	326	354	369
15-----	14	29	58	88	105	135	164	194	222	251	280	298	327	355	370
16-----	15	29	58	86	104	134	162	191	220	249	277	295	325	353	368
17-----	15	28	58	86	104	134	162	191	220	249	278	296	327	355	370
18-----	15	28	58	87	105	134	164	193	222	252	280	298	327	356	371
19-----	15	29	59	87	105	134	163	192	220	250	278	296	325	354	367
20-----	14	28	57	86	104	133	161	191	219	249	277	295	325	353	367
21-----	14	28	57	86	103	133	161	192	220	249	279	297	326	354	368
22-----	16	30	59	88	106	137	165	194	223	252	281	298	328	356	370
23-----	15	29	58	86	104	133	162	191	220	249	277	295	325	353	367

## CONSTITUENT 2N

Series	29	58	87	105	134	163	192	221	250	279	297	326	355	369
<i>Hour</i>														
0-----	28	58	86	105	135	163	193	222	251	280	299	327	357	371
1-----	30	58	88	106	135	165	194	223	252	281	299	329	357	371
2-----	28	58	87	105	134	164	193	222	250	279	297	325	353	368
3-----	30	59	88	106	136	164	193	221	251	280	298	326	356	370
4-----	29	57	86	104	132	161	190	220	249	278	295	325	353	368
5-----	28	58	86	105	134	163	192	222	251	280	298	326	356	369
6-----	30	58	88	106	135	164	194	222	252	281	298	328	356	370
7-----	29	59	88	106	135	165	193	223	251	280	297	325	354	368
8-----	29	59	88	106	135	163	192	220	249	279	296	325	365	368
9-----	29	57	86	104	133	162	191	220	250	278	296	326	354	369
10-----	29	58	87	106	135	164	193	223	251	280	298	327	357	370
11-----	29	58	87	106	135	164	194	222	251	280	298	326	355	369
12-----	29	58	88	106	135	165	193	221	249	277	295	325	354	368
13-----	29	59	87	105	134	162	190	219	248	278	296	325	354	368
14-----	29	57	86	104	133	161	191	220	250	278	297	326	355	370
15-----	29	58	87	105	133	163	192	222	250	280	298	327	356	370
16-----	29	58	87	104	134	163	192	221	251	279	297	325	354	368
17-----	29	58	88	105	134	163	192	220	249	278	296	326	355	369
18-----	29	59	87	104	132	161	189	219	248	278	296	325	354	368
19-----	29	57	86	103	133	161	191	220	249	278	297	326	355	369
20-----	30	59	88	106	134	164	193	222	251	280	299	328	357	371
21-----	28	58	87	104	134	163	192	221	250	279	297	325	354	367
22-----	30	58	87	106	135	163	192	220	249	278	296	326	355	369
23-----	28	56	85	103	131	161	189	219	248	277	295	325	354	368

Table 32.—*Divisors for primary stencil sums—Continued*  
CONSTITUENT O

Series	14	29	58	87	105	134	163	192	221	250	279	297	326	355	369
<i>Hours</i>															
0.....	13	29	58	87	106	135	164	192	222	251	279	298	327	355	369
1.....	14	29	59	88	105	134	164	192	221	251	280	298	327	355	369
2.....	14	28	57	86	105	133	162	192	221	250	279	296	325	354	368
3.....	14	30	57	87	105	134	164	193	221	251	280	297	326	356	370
4.....	14	29	58	87	106	135	163	193	222	250	280	297	325	354	369
5.....	14	29	59	87	105	135	163	192	222	251	280	297	326	355	369
6.....	14	29	58	87	105	134	164	193	222	251	279	297	326	355	369
7.....	14	28	58	87	105	135	164	192	222	251	280	297	327	355	369
8.....	14	29	58	88	106	134	164	193	221	250	278	296	325	355	368
9.....	15	30	58	87	106	134	163	193	221	250	279	297	326	355	369
10.....	14	29	58	87	105	135	164	193	221	249	279	297	326	356	370
11.....	14	30	59	88	107	136	164	192	222	251	280	299	327	356	370
12.....	14	29	59	87	105	135	163	192	221	250	279	297	327	355	369
13.....	13	28	57	87	104	132	161	189	219	248	276	295	324	353	367
14.....	14	29	58	87	105	133	163	192	220	250	279	297	327	356	369
15.....	14	29	58	87	105	133	162	192	221	250	280	297	326	355	370
16.....	14	30	58	87	104	134	163	192	221	250	279	298	325	355	369
17.....	14	29	58	86	104	133	161	191	220	248	277	296	325	354	369
18.....	13	28	58	87	104	134	163	191	221	250	279	297	327	355	369
19.....	14	29	58	88	104	133	163	192	221	250	278	297	326	355	368
20.....	15	29	58	87	105	134	163	192	220	250	279	296	326	355	369
21.....	14	29	58	87	105	133	163	192	220	249	279	297	326	356	370
22.....	15	30	57	86	105	134	162	191	221	250	279	298	326	355	369
23.....	14	28	58	86	104	134	162	192	221	249	279	297	326	355	369

## CONSTITUENT OO

Series	29	58	87	105	134	163	192	221	250	279	297	326	355	369
<i>Hours</i>														
0.....	29	58	86	104	134	163	192	221	250	280	298	326	355	369
1.....	30	60	88	107	136	164	193	221	250	279	297	326	355	369
2.....	29	57	86	103	133	162	192	220	250	280	297	327	355	369
3.....	31	60	89	107	137	166	194	223	251	281	298	327	355	370
4.....	30	58	87	104	132	162	191	220	249	278	297	326	355	369
5.....	29	59	88	106	135	166	194	223	251	280	298	326	355	369
6.....	28	58	86	104	132	161	190	219	249	277	297	325	355	368
7.....	29	58	88	105	135	165	194	223	251	280	298	327	355	369
8.....	29	59	88	105	134	163	191	220	249	278	297	326	355	369
9.....	29	58	87	105	134	163	193	223	251	280	298	326	355	369
10.....	28	58	87	105	133	162	191	219	248	277	296	325	355	368
11.....	28	57	87	104	134	162	193	222	251	280	299	327	355	369
12.....	29	58	88	106	135	163	193	221	250	278	296	326	355	369
13.....	29	57	87	104	133	162	192	221	250	280	297	325	356	370
14.....	30	59	88	107	135	164	192	222	250	278	296	325	354	369
15.....	28	57	85	104	132	162	191	220	250	279	297	327	356	369
16.....	29	58	87	106	135	164	193	222	251	279	297	326	354	369
17.....	29	57	86	104	133	161	190	220	249	278	296	326	355	369
18.....	30	58	88	106	135	165	193	224	252	281	298	327	356	370
19.....	28	57	85	104	132	161	189	218	248	277	295	323	354	368
20.....	29	58	87	106	135	164	193	222	252	280	298	326	356	369
21.....	28	58	86	104	133	161	190	218	248	277	295	324	354	368
22.....	29	57	87	105	135	163	193	222	251	281	298	327	356	370
23.....	29	58	87	105	134	163	191	220	249	278	295	325	354	369

Table 32.—*Divisors for primary stencil sums—Continued*

## CONSTITUENT P

Series	29	58	87	105	134	163	192	221	250	279	297	326	355	369
<i>Hour</i>														
0-----	29	58	87	105	135	164	193	222	251	280	298	327	356	369
1-----	29	58	87	105	134	163	192	221	250	279	297	297	354	368
2-----	29	58	87	105	134	163	192	222	251	280	298	327	355	369
3-----	29	59	88	106	135	164	193	222	251	280	298	327	356	370
4-----	29	58	87	105	134	164	193	222	251	280	297	326	355	369
5-----	29	58	87	105	134	163	192	221	250	279	296	325	354	368
6-----	29	58	87	105	134	163	192	221	251	279	297	326	355	369
7-----	29	58	88	106	135	164	193	222	251	279	297	326	356	370
8-----	29	58	87	105	134	163	192	221	249	278	296	325	354	368
9-----	29	58	87	105	134	164	193	221	250	279	297	326	355	369
10-----	29	58	87	105	134	163	192	220	249	279	297	326	355	369
11-----	29	58	88	106	135	164	192	221	250	279	297	326	356	370
12-----	29	58	87	105	134	163	191	220	249	278	296	325	354	368
13-----	29	58	87	105	134	162	192	221	250	279	297	326	355	369
14-----	30	59	88	106	135	163	192	221	250	280	298	327	356	370
15-----	29	58	87	105	134	164	193	221	250	278	296	325	354	368
16-----	29	58	87	105	133	162	191	220	249	278	296	325	354	368
17-----	29	58	87	106	134	163	192	221	250	279	297	326	355	369
18-----	30	59	87	105	134	163	192	221	250	279	298	327	356	370
19-----	29	58	86	104	133	162	191	220	249	278	296	325	354	368
20-----	29	57	86	104	134	163	192	221	250	279	297	326	355	369
21-----	29	57	86	104	133	162	191	221	250	279	297	326	355	369
22-----	28	57	86	104	133	162	191	220	249	278	296	325	354	368
23-----	28	58	87	105	134	163	192	221	250	279	298	327	356	370

## CONSTITUENT Q

Series	29	58	87	105	134	163	192	221	250	279	297	326	355	369
<i>Hour</i>														
0-----	29	59	88	106	136	164	194	222	250	280	297	326	355	368
1-----	29	58	86	104	133	162	191	221	250	280	298	327	357	370
2-----	29	59	88	106	135	165	193	222	251	280	298	326	354	368
3-----	28	56	86	103	132	161	190	220	249	278	297	326	354	369
4-----	30	59	89	107	136	166	195	225	253	282	299	328	356	370
5-----	29	58	87	105	133	162	191	219	249	277	296	325	354	369
6-----	30	59	88	107	136	165	195	224	254	281	300	328	356	371
7-----	30	58	87	104	133	162	191	219	248	277	295	324	354	369
8-----	28	58	87	105	135	164	194	223	251	280	298	326	356	369
9-----	29	59	88	106	135	163	192	221	248	278	296	325	355	369
10-----	28	58	86	104	134	163	192	221	250	280	298	327	355	370
11-----	29	58	88	106	134	164	192	220	249	277	295	324	353	368
12-----	29	57	86	104	133	163	191	220	250	279	297	327	356	370
13-----	30	59	89	107	136	165	192	221	250	278	296	325	354	369
14-----	29	58	87	104	133	161	191	220	250	279	297	327	356	371
15-----	30	59	88	107	136	164	194	222	251	280	297	326	355	368
16-----	29	58	86	104	133	161	190	219	248	278	296	325	355	368
17-----	29	59	87	106	135	164	193	223	251	280	298	326	355	368
18-----	28	57	85	103	131	160	188	218	247	277	295	324	354	367
19-----	29	58	87	105	134	164	193	223	252	281	299	328	356	370
20-----	29	57	86	104	132	161	190	219	247	276	294	324	353	367
21-----	30	57	87	105	134	164	193	222	252	281	299	328	356	370
22-----	29	58	87	105	134	162	191	220	249	277	295	325	354	368
23-----	27	56	85	103	133	162	192	221	251	280	298	327	357	370

Table 32.—*Divisors for primary stencil sums—Continued*

## CONSTITUENT 2Q

Series	29	58	87	105	134	163	192	221	250	279	297	326	355	369
<i>Hour</i>														
0-----	25	50	83	113	142	167	192	217	242	279	309	334	359	371
1-----	25	59	101	116	141	166	191	216	255	293	308	333	358	370
2-----	36	77	102	117	142	167	192	233	269	293	309	334	359	371
3-----	39	64	89	104	129	154	196	230	255	279	295	320	345	366
4-----	25	50	75	90	115	159	192	217	242	266	282	307	335	370
5-----	25	50	75	90	136	167	192	217	242	266	282	332	358	370
6-----	25	50	83	113	142	167	192	217	241	277	309	334	358	370
7-----	25	60	102	117	142	167	192	217	254	293	309	334	358	370
8-----	36	76	101	116	141	166	191	232	267	292	308	333	357	369
9-----	39	64	89	104	129	154	197	230	255	280	296	320	345	365
10-----	25	50	75	90	115	159	191	215	240	265	281	305	333	369
11-----	25	50	75	90	136	167	192	216	241	266	283	331	358	370
12-----	25	50	83	113	142	167	192	216	241	277	307	332	357	369
13-----	25	60	101	117	142	167	191	216	254	293	308	333	358	370
14-----	37	77	101	117	142	167	191	231	268	293	308	333	358	370
15-----	38	63	87	103	128	153	194	229	254	279	294	319	344	364
16-----	25	50	74	90	115	159	191	216	241	266	281	306	334	370
17-----	25	49	74	90	136	165	190	215	240	265	280	330	357	369
18-----	25	49	81	113	142	166	191	216	241	278	308	333	358	370
19-----	25	58	101	117	142	166	191	216	254	292	307	332	357	369
20-----	36	76	101	117	141	166	191	231	268	293	308	333	358	370
21-----	37	62	87	103	127	152	194	229	254	279	294	319	344	364
22-----	24	49	74	90	114	158	191	216	241	266	281	306	334	370
23-----	24	49	74	90	135	166	191	216	241	266	281	331	358	370

## CONSTITUENT R

Series	29	58	87	105	134	163	192	221	250	279	297	326	355	369
<i>Hour</i>														
0-----	30	59	88	106	135	164	193	222	251	280	298	327	356	370
1-----	29	59	88	106	135	164	193	222	251	280	298	326	355	369
2-----	29	58	88	106	135	164	193	222	251	279	297	326	355	369
3-----	29	58	87	105	135	164	193	221	250	279	297	326	355	369
4-----	29	58	87	105	134	163	192	221	250	279	297	326	355	369
5-----	29	58	86	104	133	162	192	221	250	279	297	326	355	369
6-----	28	57	86	104	133	162	191	221	250	279	297	326	355	369
7-----	29	58	87	105	134	163	192	221	251	280	298	327	356	370
8-----	29	58	87	105	134	163	192	221	250	280	298	327	356	370
9-----	29	58	87	105	134	163	192	221	250	279	298	327	356	370
10-----	29	58	87	105	134	163	192	221	250	279	297	327	356	370
11-----	29	58	87	105	134	163	192	221	250	279	297	326	356	370
12-----	29	58	87	105	134	163	192	221	250	279	297	326	354	368
13-----	29	58	87	105	134	163	192	221	250	279	296	325	354	368
14-----	29	58	87	105	134	163	192	221	249	278	296	325	354	368
15-----	29	58	87	105	134	163	191	220	249	278	296	325	354	368
16-----	29	58	87	105	133	162	191	220	249	278	296	325	354	368
17-----	29	57	86	104	133	162	191	220	249	278	296	325	354	368
18-----	29	58	87	105	134	163	192	221	250	279	297	326	355	369
19-----	29	58	87	105	134	163	192	221	250	279	297	326	355	369
20-----	29	58	87	105	134	163	192	221	250	279	297	326	355	369
21-----	29	58	87	105	134	163	192	221	250	279	297	326	355	369
22-----	29	58	87	105	134	163	192	221	250	279	297	326	355	369
23-----	29	58	87	105	134	163	192	221	250	279	297	326	355	369

Table 32.—*Divisors for primary stencil sums—Continued*

## CONSTITUENT T

Series	29	58	87	105	134	163	192	221	250	279	297	326	355	369
<i>Hour</i>														
0	29	58	88	106	135	164	193	222	251	280	298	327	356	370
1	29	58	87	105	134	163	192	221	250	279	297	326	355	369
2	29	58	87	105	134	163	192	221	250	279	297	326	355	369
3	29	58	87	105	134	163	192	221	250	279	297	326	355	369
4	29	58	87	105	134	163	193	222	251	280	298	328	357	371
5	30	59	88	106	135	164	193	222	251	280	298	327	356	370
6	29	58	87	105	134	163	192	221	250	279	297	326	355	369
7	29	58	87	105	134	163	192	221	250	279	297	326	355	369
8	29	58	87	105	134	163	192	221	250	279	297	326	355	369
9	29	58	87	105	135	164	193	222	251	281	299	328	357	371
10	29	58	87	105	134	163	192	221	250	279	297	326	355	369
11	29	58	87	105	134	163	192	221	250	279	297	326	355	369
12	29	58	87	105	134	163	192	221	250	279	297	326	354	368
13	29	58	87	105	134	163	192	221	250	279	297	325	355	369
14	29	59	88	106	135	164	193	223	252	281	298	327	356	370
15	29	58	87	105	134	163	192	221	250	278	296	325	354	368
16	29	58	87	105	134	163	192	221	249	278	296	325	354	368
17	29	58	87	105	134	163	192	220	249	278	296	325	354	368
18	29	58	87	105	134	163	191	220	249	278	297	326	355	369
19	29	58	87	105	134	163	192	221	250	279	297	326	355	369
20	29	58	87	105	133	162	191	220	249	278	296	325	354	368
21	29	58	86	104	133	162	191	220	249	278	296	325	354	368
22	29	57	86	104	133	162	191	220	249	278	296	325	354	368
23	28	57	86	104	133	162	191	220	250	279	297	326	355	369

## CONSTITUENT λ

Series	29	58	87	105	134	163	192	221	250	279	297	326	355	369
<i>Hour</i>														
0	29	58	89	107	135	164	194	223	252	280	298	330	358	372
1	29	57	87	106	134	162	191	221	250	278	296	325	355	369
2	29	57	86	104	134	162	191	219	250	278	296	324	354	369
3	31	59	88	105	136	165	194	222	252	282	300	328	357	371
4	31	59	88	105	134	164	193	221	250	279	298	326	355	369
5	29	59	88	105	134	162	193	221	250	278	297	326	355	369
6	29	58	88	105	134	162	193	221	250	278	296	326	355	369
7	29	57	88	106	135	163	192	223	252	280	298	326	358	371
8	29	57	86	104	134	162	191	219	250	278	296	324	354	368
9	30	58	87	104	135	163	192	220	250	279	297	325	354	367
10	31	60	88	106	135	166	195	223	252	282	301	329	358	371
11	28	59	87	105	134	162	193	221	249	278	297	326	355	368
12	28	57	87	105	134	162	191	221	249	278	296	326	355	368
13	28	57	87	105	134	162	190	221	249	278	296	324	354	368
14	28	57	85	105	134	163	191	220	251	280	298	326	355	371
15	29	58	86	104	134	163	191	220	249	279	296	325	353	367
16	30	59	87	105	133	164	192	221	249	280	297	326	354	368
17	28	60	88	106	134	164	194	223	251	280	299	329	357	371
18	28	57	87	105	133	162	191	221	249	278	295	326	354	368
19	28	57	86	105	133	162	190	221	249	278	295	324	354	368
20	28	57	85	104	133	162	190	219	249	278	295	324	353	368
21	29	58	86	104	134	164	192	221	250	281	298	327	355	370
22	30	59	87	105	133	164	192	221	249	279	297	326	354	368
23	28	58	87	105	133	163	192	221	249	277	296	326	354	368

Table 32.—*Divisors for primary stencil sums—Continued*CONSTITUENT  $\mu$ 

Series	29	58	87	105	134	163	192	221	250	279	297	326	355	369
<i>Hour</i>														
0-----	29	59	89	105	135	163	192	223	252	280	299	326	356	369
1-----	30	61	88	107	135	164	194	223	252	282	298	327	356	369
2-----	30	57	88	105	134	164	193	221	250	280	296	326	355	369
3-----	27	57	87	104	134	163	190	219	249	276	296	325	353	368
4-----	30	60	87	106	135	162	192	222	249	279	298	326	356	369
5-----	30	57	86	106	133	163	193	220	250	280	297	327	356	370
6-----	27	56	86	103	133	163	190	220	250	278	297	326	354	369
7-----	29	59	88	106	136	163	193	223	251	280	299	326	355	369
8-----	30	59	87	107	134	164	194	222	250	279	296	326	356	369
9-----	29	57	87	104	134	163	191	219	249	279	296	326	354	369
10-----	28	58	88	105	134	163	191	221	251	278	298	326	355	369
11-----	30	59	86	105	134	162	192	222	249	279	297	326	356	369
12-----	29	57	86	105	133	163	193	220	250	280	297	327	356	369
13-----	28	57	87	104	134	164	191	221	251	279	297	326	353	369
14-----	29	59	88	106	136	163	193	222	250	278	298	325	355	368
15-----	30	59	87	107	134	164	193	222	250	280	297	327	357	370
16-----	29	57	87	104	133	162	191	219	249	278	296	326	354	369
17-----	28	57	86	103	133	162	190	220	249	277	297	325	354	368
18-----	29	59	86	106	135	163	193	222	250	280	298	327	356	369
19-----	30	57	87	106	134	164	193	221	251	281	298	327	356	370
20-----	27	57	87	104	134	163	191	221	250	277	296	325	353	369
21-----	30	60	88	106	136	164	193	222	250	279	297	326	356	369
22-----	30	58	87	105	132	162	192	220	249	279	295	325	356	369
23-----	28	56	85	101	131	161	190	219	249	278	295	325	352	369

CONSTITUENT  $\nu$ 

Series	29	58	87	105	134	163	192	221	250	279	297	326	355	369
<i>Hour</i>														
0-----	31	59	86	103	135	165	193	221	249	283	300	327	355	368
1-----	28	56	83	103	134	161	189	216	250	278	295	322	351	367
2-----	28	56	89	107	135	162	190	224	252	280	297	324	358	371
3-----	28	60	89	106	134	161	195	222	250	278	295	329	357	370
4-----	33	62	90	107	135	167	196	223	251	280	302	329	357	370
5-----	30	57	85	102	135	164	192	219	249	281	299	326	354	367
6-----	28	55	83	104	134	161	189	217	250	277	295	322	352	369
7-----	28	55	90	107	135	162	191	224	252	279	296	325	358	371
8-----	28	61	89	106	134	161	195	222	250	277	295	329	357	370
9-----	34	62	90	107	134	169	197	224	252	282	302	330	358	371
10-----	29	56	84	101	134	162	190	217	248	279	296	324	351	365
11-----	28	55	85	107	134	162	190	219	251	278	295	323	353	371
12-----	28	56	89	106	133	161	190	223	251	278	295	326	356	370
13-----	29	62	90	107	134	164	195	223	251	278	298	330	357	371
14-----	32	60	88	105	134	167	194	222	250	281	300	328	355	369
15-----	27	55	83	101	133	161	188	216	247	278	295	323	350	365
16-----	27	55	86	107	134	162	189	221	250	278	295	323	355	371
17-----	27	58	88	106	133	161	192	223	250	278	295	327	356	369
18-----	30	62	89	107	134	165	195	223	250	278	299	330	357	370
19-----	31	59	86	103	134	166	193	221	248	282	299	327	354	367
20-----	27	55	82	101	133	161	188	216	249	278	295	322	350	366
21-----	27	55	87	106	134	162	189	222	250	278	295	322	356	369
22-----	27	59	88	105	133	160	193	223	250	278	295	328	357	370
23-----	31	62	89	106	134	165	195	223	250	279	300	328	356	369

Table 32.—*Divisors for primary stencil sums—Continued*CONSTITUENT  $\rho$ 

Series	29	58	87	105	134	163	192	221	250	279	297	326	355	369
<i>Hour</i>														
0-----	30	59	89	107	135	164	193	222	251	279	297	326	355	369
1-----	29	59	88	105	134	164	193	222	251	280	299	328	356	372
2-----	28	57	87	105	134	162	191	220	249	278	295	324	353	367
3-----	29	58	87	106	135	165	194	224	252	281	299	328	357	371
4-----	30	58	87	106	135	163	192	221	250	279	297	326	355	370
5-----	29	58	87	106	134	163	192	220	250	278	296	325	354	368
6-----	28	58	87	106	136	165	193	223	252	282	299	328	357	371
7-----	29	58	87	104	134	163	192	221	249	278	296	325	354	369
8-----	29	58	87	105	135	165	194	222	251	280	298	326	355	370
9-----	29	58	87	105	133	162	192	222	251	280	298	327	357	371
10-----	29	57	86	104	133	162	191	220	249	278	296	325	353	366
11-----	28	58	87	105	134	163	193	223	252	280	298	327	356	369
12-----	29	58	86	104	133	162	190	219	249	279	297	326	355	369
13-----	30	59	88	106	135	163	192	221	250	280	297	326	355	368
14-----	29	58	87	104	134	163	193	221	250	280	299	328	357	370
15-----	28	57	86	104	132	162	190	219	248	276	295	324	354	367
16-----	30	59	88	106	135	164	193	222	250	279	298	327	355	369
17-----	28	57	86	104	133	161	191	220	250	279	298	327	356	370
18-----	29	58	87	104	133	162	191	220	248	277	295	325	354	367
19-----	30	58	87	106	135	164	193	222	251	280	297	326	356	369
20-----	28	57	86	104	132	161	190	219	249	277	295	325	354	368
21-----	30	59	87	105	134	163	191	220	249	278	296	324	353	368
22-----	29	58	88	105	135	164	193	222	251	281	298	328	357	371
23-----	29	58	86	104	133	162	191	219	248	277	295	323	352	367

## CONSTITUENT MK

Series	29	58	87	105	134	163	192	221	250	279	297	326	355	369
<i>Hour</i>														
0-----	30	59	88	105	135	164	192	222	251	279	297	325	355	368
1-----	29	58	88	106	135	163	192	222	251	280	298	327	356	369
2-----	29	58	88	105	134	164	192	221	249	278	297	326	355	369
3-----	30	58	86	104	133	163	191	221	250	278	296	325	355	368
4-----	29	58	87	106	134	163	192	222	251	280	298	326	356	369
5-----	29	58	87	105	134	164	192	220	250	279	298	327	356	369
6-----	30	58	86	105	134	164	193	221	250	279	297	326	356	369
7-----	30	58	87	105	133	163	192	221	251	279	297	326	355	369
8-----	30	59	88	106	135	164	193	221	251	280	299	327	355	369
9-----	28	57	86	105	134	164	192	220	249	278	296	325	354	369
10-----	29	58	86	104	133	163	192	221	251	279	297	326	355	369
11-----	29	59	88	106	134	162	192	221	250	279	298	326	354	369
12-----	28	58	86	105	134	163	192	220	250	279	297	326	355	369
13-----	29	58	86	105	133	162	192	221	250	278	297	326	355	369
14-----	29	59	88	106	134	163	193	222	250	280	297	326	355	370
15-----	29	59	87	106	135	163	192	221	250	280	297	327	355	369
16-----	28	57	86	105	133	162	192	220	249	278	296	326	355	369
17-----	29	59	87	104	134	163	193	222	250	279	296	325	354	369
18-----	29	58	88	105	134	162	191	220	249	279	296	326	354	368
19-----	28	57	87	105	135	163	193	221	250	279	297	327	356	370
20-----	29	57	86	103	133	162	191	221	249	278	296	325	354	369
21-----	29	58	88	105	134	162	191	221	250	280	297	326	355	369
22-----	28	57	87	105	135	163	191	221	250	280	297	327	355	369
23-----	29	57	87	104	134	163	192	221	249	278	297	325	355	370

Table 32.—*Divisors for primary stencil sums—Continued*

## CONSTITUENT 2MK

Series	29	58	87	105	134	163	192	221	250	279	297	326	355	369
<i>Hour</i>														
0	30	59	87	106	134	164	193	221	251	280	298	326	355	369
1	29	59	87	106	134	164	193	221	250	280	298	327	356	370
2	29	59	87	106	134	163	192	221	250	279	297	325	355	368
3	29	58	86	105	133	163	193	222	251	280	298	326	355	369
4	30	59	87	105	134	164	192	221	250	279	297	326	355	369
5	29	59	88	106	135	164	192	222	251	279	298	327	357	370
6	29	58	86	104	134	163	191	221	250	278	297	325	354	368
7	30	59	87	105	135	164	192	221	251	280	299	327	356	370
8	30	58	87	105	134	163	192	221	250	279	296	326	355	368
9	29	57	87	104	134	164	192	222	251	279	297	326	356	369
10	30	58	88	105	135	164	192	221	251	279	297	326	355	369
11	30	59	88	106	135	164	193	222	251	280	297	327	356	370
12	29	57	87	105	134	162	192	221	249	278	296	326	356	369
13	29	57	87	104	134	162	191	221	249	278	296	325	354	368
14	29	57	86	104	133	162	191	220	249	279	297	326	355	369
15	29	58	87	105	134	162	192	221	249	279	296	326	355	369
16	28	57	87	105	135	163	192	222	250	279	297	326	354	369
17	28	57	86	104	133	162	191	220	249	278	296	325	353	368
18	29	59	88	106	135	163	193	222	250	280	298	328	356	371
19	28	58	87	105	134	162	192	221	249	278	296	325	354	368
20	28	57	87	104	133	162	191	220	250	279	297	326	355	369
21	29	58	87	105	133	163	192	220	250	279	297	326	354	369
22	28	58	87	106	134	163	193	221	250	279	297	326	355	370
23	28	57	87	104	133	162	191	219	249	278	296	325	354	368

## CONSTITUENT MN

Series	29	58	87	105	134	163	192	221	250	279	297	326	355	269
<i>Hour</i>														
0	28	56	85	104	134	163	193	223	253	283	301	328	356	370
1	30	60	90	109	139	166	194	222	250	277	296	325	355	370
2	28	56	84	101	129	158	188	218	248	278	297	326	355	369
3	30	60	91	109	139	168	198	226	254	281	299	326	355	370
4	30	59	87	104	132	159	187	217	248	277	296	325	355	369
5	28	57	88	106	136	165	195	225	256	283	301	328	356	369
6	30	61	90	109	136	164	192	220	247	277	295	325	355	369
7	28	56	83	101	130	160	190	221	250	280	298	327	355	368
8	30	61	90	109	138	168	196	224	251	279	296	325	355	369
9	29	57	84	102	129	157	187	218	247	277	295	325	355	369
10	30	60	89	108	137	167	198	228	255	283	300	328	356	369
11	31	61	89	107	134	162	190	218	247	277	295	325	356	370
12	28	55	84	102	132	162	193	222	252	282	299	327	355	368
13	30	59	89	107	137	165	193	220	248	276	294	324	355	369
14	28	55	83	100	128	159	189	218	248	278	296	327	355	368
15	30	59	89	107	137	168	198	225	253	281	298	326	356	370
16	30	58	86	103	131	159	187	216	246	276	294	325	355	369
17	28	56	86	104	135	165	195	224	254	282	299	327	355	368
18	29	59	89	107	135	163	190	218	246	276	295	325	354	369
19	27	55	83	100	131	161	190	220	250	280	299	328	355	369
20	29	59	89	108	138	168	195	223	251	279	296	325	354	369
21	28	56	84	101	129	157	186	216	246	276	295	325	354	369
22	28	58	88	107	137	167	196	226	254	282	299	327	354	368
23	29	59	88	105	133	161	188	216	246	276	295	325	354	369

Table 32.—*Divisors for primary stencil sums—Continued*

## CONSTITUENT MS

Series	29	58	87	105	134	163	192	221	250	279	297	326	355	369
<i>Hour</i>														
0-----	30	59	88	106	135	164	192	222	250	279	297	326	354	369
1-----	30	58	88	106	134	164	192	221	250	279	296	326	355	369
2-----	29	58	87	105	134	163	192	222	250	280	298	327	355	369
3-----	30	58	88	107	135	165	194	223	252	281	298	328	356	370
4-----	29	58	87	105	134	163	191	221	249	279	296	325	354	368
5-----	30	58	88	105	134	164	192	221	250	280	297	327	356	370
6-----	29	58	88	105	135	164	194	223	252	281	299	328	356	371
7-----	30	58	87	105	134	162	192	220	250	278	296	326	354	368
8-----	29	58	87	104	134	162	191	220	249	278	296	326	355	369
9-----	29	57	87	104	133	162	192	220	250	279	297	326	355	369
10-----	30	59	88	106	136	164	193	222	251	279	298	327	355	370
11-----	30	58	88	105	134	163	192	220	250	278	296	326	354	368
12-----	28	58	87	104	134	162	192	221	250	279	298	326	356	370
13-----	28	57	86	105	134	163	193	221	251	279	297	325	355	369
14-----	29	59	87	105	135	163	192	221	250	278	297	325	354	369
15-----	29	58	87	105	134	163	192	220	250	278	296	325	355	369
16-----	28	58	86	104	134	163	193	222	251	280	299	327	356	371
17-----	29	58	87	106	135	163	193	221	251	279	297	326	355	369
18-----	28	58	86	104	133	162	190	220	248	278	296	324	354	367
19-----	28	57	86	104	132	162	191	220	249	279	297	326	356	369
20-----	29	59	87	106	134	164	192	222	250	279	298	326	355	369
21-----	29	58	86	105	133	162	191	220	249	278	296	325	354	367
22-----	28	58	86	104	133	162	190	220	248	278	296	325	355	368
23-----	28	57	86	105	133	163	192	221	250	280	297	326	356	369

## CONSTITUENT 2SM

Series	29	58	87	105	134	163	192	221	250	279	297	326	355	369
<i>Hour</i>														
0-----	28	57	87	106	136	164	192	220	250	280	299	329	356	369
1-----	30	60	88	106	133	163	193	223	252	280	297	325	355	370
2-----	28	56	86	105	135	165	193	220	249	279	297	327	356	370
3-----	30	60	89	107	135	163	193	223	253	281	298	326	355	370
4-----	28	55	84	103	133	163	191	219	247	276	294	324	353	367
5-----	30	60	90	107	135	163	193	223	253	282	299	326	355	370
6-----	28	56	84	103	133	163	192	220	248	277	295	325	355	370
7-----	30	60	90	109	137	164	193	223	253	283	300	328	356	370
8-----	29	57	85	103	133	163	193	221	249	277	295	325	355	370
9-----	29	59	89	108	137	165	193	222	252	282	300	328	356	370
10-----	30	59	86	104	133	163	193	223	251	279	295	325	355	370
11-----	28	57	87	106	136	164	192	220	249	279	298	327	355	369
12-----	30	59	87	104	132	161	191	221	249	277	295	323	353	367
13-----	28	57	87	105	135	164	191	219	249	279	298	328	356	369
14-----	30	60	88	105	133	162	192	222	251	279	297	325	355	369
15-----	27	55	85	103	133	163	191	219	247	277	296	326	355	368
16-----	30	60	89	106	134	162	192	222	252	281	298	326	355	369
17-----	28	56	84	102	132	162	191	219	247	276	295	325	355	368
18-----	30	60	90	108	135	163	192	222	252	282	300	328	355	369
19-----	29	57	85	102	132	162	192	220	248	276	295	325	355	369
20-----	29	59	89	107	135	163	190	220	250	280	298	326	354	367
21-----	29	57	85	102	132	162	192	221	249	276	295	325	355	369
22-----	28	58	88	106	135	163	191	220	250	280	299	327	355	368
23-----	30	58	86	103	132	162	192	222	250	278	295	325	355	369

Table 33.—*For construction of secondary stencils*

Constituent A.	J				S				L			
	OO		2SM		K and P		R and T		MS		λ	
Page	J hours	Difference, hours	J hours	Difference, hours	S hours	Difference, hours	S hours	Difference, hours	L hours	Difference, hours	L hours	Difference, hours
1.....	0-23	3	0-23	-0	0-23	±0	0-23	±0	0-23	-0	0-23	-0
2.....	10- 3	9	0-23	1	0-23	1	0-23	0	0-23	0	0-23	1
3.....	16- 4	15	0-23	2	0-23	1	0-23	1	17-21	0	0-23	1
4.....	23- 5	21	0-23	3	0-23	2	0-23	1	0-23	1	0-23	1
5.....	5- 6	3	0-23	4	0-23	2	0-23	1	0-23	1	0-23	2
6.....	0-23	10	0-23	5	0- 1	2	0-23	1	0-23	1	0-23	2
7.....	19-12	16	0-23	6	0-23	3	0-15	1	0-23	1	0-23	3
8.....	1-12	22	1-11	6	0-23	3	0-23	2	0-23	2	0-23	3
9.....	8-13	4	0-23	7	0-23	4	0-23	2	0-23	2	0-23	3
10.....	14	10	0-23	8	0-23	4	0-23	2	0-23	2	0-23	4
11.....	0-23	17	0-23	9	0-23	5	0-23	2	0-23	2	0-23	4
12.....	3-20	23	0-23	10	0-23	5	0-23	3	0-23	2	0-23	5
13.....	10-21	5	0-23	11	0-23	6	0-23	3	0-23	3	0-23	5
14.....	16-22	11	0-23	12	0-23	6	0-23	3	0-23	3	12-20	5
15.....	23	17	0-23	13	0-23	7	0-23	3	0-23	3	0-23	6
16.....	6- 4	0	6- 3	13	0-23	7	0-23	4	0-23	3	0-23	6
17.....	12- 5	6	0-23	14	0-23	8	0-23	4	0-23	3	0-23	7
18.....	19- 6	12	0-23	15	0-23	8	0-23	4	0-23	4	0-23	7
19.....	1- 7	18	0-23	16	0- 8	8	0-23	4	0-23	4	0-23	8
20.....	8	0	0-23	17	0-23	9	0-23	4	0-23	4	0-23	8
21.....	0-23	7	0-23	18	0-23	9	0-23	5	0-23	4	0-23	8
22.....	21-14	13	0-23	19	0-23	10	0-23	5	0-23	4	0-23	9
23.....	4-14	19	4- 9	19	0-23	10	0-23	5	0-23	5	0-23	9
24.....	10-15	1	0-23	20	0-23	11	0-23	5	0-23	5	0-23	10
25.....	0-23	8	0-23	21	0-23	11	0-23	6	0-23	5	0-23	10
26.....	23-21	14	0-23	22	0-23	12	0-23	6	0-23	5	0-23	10
27.....	6-22	20	0-23	23	0-23	12	0-23	6	0-23	5	0-23	11
28.....	12-23	2	0-23	0	0-23	13	0-23	6	0-23	6	0-23	11
29.....	19- 0	8	0-23	1	0-23	13	0-23	7	0-23	6	0-23	12
30.....	1	14	0-23	2	0-23	14	0-23	7	0-23	6	0-23	12
31.....	8- 6	21	8- 1	2	0-23	14	0-23	7	0-23	6	0-23	12
32.....	15- 7	3	0-23	3	0-15	14	0-23	7	0-23	6	0-23	13
33.....	21- 8	9	0-23	4	0-23	15	0-23	7	0-23	7	0-23	13
34.....	4- 9	15	0-23	5	0-23	15	0-23	8	0-23	7	0-23	14
35.....	0-23	22	0-23	6	0-23	16	0-23	8	0-23	7	0-23	14
36.....	17-15	4	0-23	7	0-23	16	0-23	8	0-23	7	2-21	14
37.....	23-15	10	0-23	8	0-23	17	0-23	8	0-23	7	0-23	15
38.....	6-16	16	6- 8	8	0-23	17	0-23	9	0-23	8	0-23	15
39.....	12-17	22	0-23	9	0-23	18	0-23	9	0-23	8	0-23	16
40.....	0-23	5	0-23	10	0-23	18	0-23	9	0-23	8	0-23	16
41.....	2-23	11	0-23	11	0-23	19	0-23	9	0-23	8	13-15	16
42.....	8- 0	17	0-23	12	0-23	19	0-23	10	0-23	8	0-23	17
43.....	15- 1	23	0-23	13	0-23	20	0-23	10	0-23	9	0-23	17
44.....	21- 2	5	0-23	14	0-23	20	0-23	10	0-23	9	0-23	18
45.....	0-23	12	0-23	15	0-23	20	0-23	10	0-23	9	0-23	18
46.....	10- 8	18	10-23	15	0-23	21	0-23	10	0-23	9	0-23	19
47.....	17- 9	0	0-23	16	0-23	21	0-23	11	0-23	9	0-23	19
48.....	23-10	6	0-23	17	0-23	22	0-23	11	0-23	10	0-23	19
49.....	6-11	12	0-23	18	0-23	22	0-23	11	0-23	10	0-23	20
50.....	0-23	19	0-23	19	0-23	23	0-23	11	0-23	10	0-23	20
51.....	19-16	1	0-23	20	0-23	23	0-23	12	0-23	10	0-23	21
52.....	2-17	7	0-23	21	0-23	0	0-23	12	8-16	10	0-23	21
(53)....	7-14	12	0-23	21	0-23	0	0-23	12	0-23	11	0-23	21

Table 33.—*For construction of secondary stencils—Continued*

Constituent A	L		M		N				O					
	MK		MN		2MK		ν		μ		2N			
Constituent B	L hours	Differ- ence, hours	M hours	Differ- ence, hours	N hours	Differ- ence, hours	N hours	Differ- ence, hours	O hours	Differ- ence, hours	O hours	Differ- ence, hours		
Page														
1	23-10	-	0	0-23	-	20-7	+	0	0-23	+	0	0-23	+	0
2	20- 8	1	0-23	2	11-23	1	0-23	1	0-23	1	0-23	1	0	0
3	17- 5	2	0-23	4	2-14	2	0-23	1	0-23	1	0-23	1	0	0
4	15- 3	3	0-23	5	17- 6	3	0-23	1	0-23	2	0-23	2	0	0
5	12- 1	4	0-23	7	9-21	4	0-23	2	0-23	2	0-23	2	0	0
6	9-22	5	0-23	8	0-13	5	0-23	2	7- 8	2	0-23	2	0	0
7	7-20	6	0-23	10	15- 4	6	0-23	3	0-23	3	0-23	3	0	0
8	4-17	7	5- 0	11	6-19	7	0-23	3	0-23	3	0-23	3	0	0
9	2-15	8	0-23	13	22-11	8	0-23	3	0-23	4	0-23	4	0	0
10	23-12	9	18- 8	14	13- 2	9	0-23	4	0-23	4	0-23	4	0	0
11	20-10	10	0-23	16	4-18	10	0-23	4	0-23	5	0-23	5	1	
12	18- 7	11	7-15	17	20- 9	11	0-23	5	0-23	5	0-23	5	1	
13	15- 5	12	0-23	19	11- 1	12	0-23	5	0-23	6	0-23	6	1	
14	12- 2	13	19-22	20	2-16	13	2-10	5	0-23	6	0-23	6	1	
15	10- 0	14	0-23	22	17- 2	14	0-23	6	0-23	7	0-23	7	1	
16	7-21	15	0-23	0	9-23	15	0-23	6	0-23	7	0-23	7	1	
17	4-19	16	0-23	1	0-15	16	0-23	7	0-23	8	0-23	8	1	
18	2-17	17	0-23	3	15- 6	17	0-23	7	0-23	8	0-23	8	1	
19	23-14	18	0-23	4	6-21	18	0-23	8	21- 5	8	0-23	8	1	
20	21-12	19	0-23	6	22-13	19	0-23	8	0-23	9	0-23	9	1	
21	18- 9	20	0-23	7	13- 4	20	0-23	8	0-23	9	0-23	9	1	
22	15- 7	21	0-23	9	4-20	21	0-23	9	0-23	10	0-23	10	1	
23	13- 5	22	0-23	10	19-11	22	0-23	9	0-23	10	0-23	10	1	
24	10- 2	23	0-23	12	11- 3	23	0-23	10	0-23	11	0-23	11	1	
25	7- 0	0	0-23	13	2-18	0	0-23	10	0-23	11	0-23	11	1	
26	5-21	1	0-23	15	17-10	1	0-23	10	0-23	12	0-23	12	1	
27	2-19	2	0-23	16	8- 1	2	0-23	11	0-23	12	0-23	12	1	
28	23-16	3	0-23	18	0-16	3	0-23	11	0-23	13	0-23	13	1	
29	21-14	4	6- 0	19	15- 8	4	0-23	12	0-23	13	0-23	13	1	
30	18-11	5	0-23	21	6-23	5	0-23	12	0-23	14	0-23	14	2	
31	15- 9	6	19- 7	22	22-15	6	0-23	12	0-23	14	0-23	14	2	
32	13- 6	7	0-23	0	13- 6	7	0-23	13	11- 4	14	0-23	14	2	
33	10- 4	8	7-15	1	4-22	8	0-23	13	0-23	15	0-23	15	2	
34	8- 1	9	0-23	3	19-13	9	0-23	14	0-23	15	0-23	15	2	
35	5-23	10	20-22	4	11- 5	10	0-23	14	0-23	16	0-23	16	2	
36	2-21	11	0-23	6	2-20	11	2-20	14	0-23	16	0-23	16	2	
37	0-18	12	0-23	8	17-12	12	0-23	15	0-23	17	0-23	17	2	
38	21-16	13	0-23	9	8- 3	13	0-23	15	0-23	17	0-23	17	2	
39	18-13	14	0-23	11	0-18	14	0-23	16	0-23	18	0-23	18	2	
40	16-11	15	0-23	12	15-10	15	0-23	16	0-23	18	0-23	18	2	
41	13- 8	16	0-23	14	6- 1	16	6- 9	16	0-23	19	0-23	19	2	
42	10- 6	17	0-23	15	21-17	17	0-23	17	0-23	19	0-23	19	2	
43	8- 4	18	0-23	17	13- 8	18	0-23	17	0-23	20	0-23	20	2	
44	5- 1	19	0-23	18	4- 0	19	0-23	18	0-23	20	0-23	20	2	
45	3-23	20	0-23	20	19-15	20	0-23	18	1-23	20	0-23	20	2	
46	0-20	21	0-23	21	10- 6	21	0-23	19	0-23	21	0-23	21	2	
47	21-18	22	0-23	23	2-22	22	0-23	19	0-23	21	0-23	21	2	
48	19-15	23	18-16	0	17-13	23	0-23	19	0-23	22	0-23	22	2	
49	16-13	0	0-23	2	8- 5	0	0-23	20	0-23	22	0-23	22	3	
50	13-10	1	6-23	3	0-20	1	0-23	20	0-23	23	0-23	23	3	
51	11- 8	2	0-23	5	15-12	2	0-23	21	0-23	23	0-23	23	3	
52	8- 5	3	19- 7	6	6- 3	3	0-23	21	0-23	0	0-23	0	3	
(53)	0-23	4	0-23	8	0-23	4	0-23	21	0-23	0	0-23	0	3	

Table 33.—*For construction of secondary stencils—Continued*

Constituent A.....		O									
Constituent B.....		P		Q		2Q					
Page	hours	Differ-	hours	O	Differ-	O	Differ-	O	Differ-	O	Differ-
		ence,		hours	ence,	hours	ence,	hours	ence,	hours	ence,
1.....	18- 1	-	2	0-23	-	3	18-22	-	5	23-11	-
2.....	0-23	8	6- 3	9	6- 8	17	9-20	18	6	12-17	7
3.....	18-15	13	18-12	15	18- 6	6	7-17	7	7	21- 5	19
4.....	7-19	18	7-22	21	7-16	18	17- 5	19	6	-----	20
5.....	19-22	23	19- 8	3	19- 1	6	2-14	7	7	15-18	8
6.....	0-23	5	7-18	9	7-11	18	12- 0	19	1	6	20
7.....	19-12	10	19- 3	15	19-21	6	22-10	7	7	11-18	8
8.....	7-16	15	7-13	21	7-19	19	20- 6	20	-----	-----	-----
9.....	19	20	19-23	3	19- 5	7	6-18	8	-----	-----	-----
10.....	8- 5	2	8	9	8-15	19	16- 3	20	4	7	21
11.....	20- 9	7	0-23	16	20- 0	7	1-13	8	14-19	9	-----
12.....	8-13	12	8- 5	22	8-10	19	11-23	20	0- 7	21	-----
13.....	0-23	18	20-15	4	20	7	21- 8	8	9-19	9	-----
14.....	8- 2	23	8- 1	10	8-18	20	19- 7	21	-----	-----	-----
15.....	20- 6	4	20-10	16	20- 4	8	5-17	9	18-19	10	-----
16.....	9	9	9-20	22	9-14	20	15- 2	21	3- 8	22	-----
17.....	21-19	15	21- 6	4	21-23	8	0-12	9	13-20	10	-----
18.....	9-23	20	9-15	10	9	20	10-22	21	23- 8	22	-----
19.....	21- 3	1	21- 1	16	21- 7	9	8-20	10	-----	-----	-----
20.....	0-23	7	9-11	22	9-17	21	18- 6	22	7- 8	23	-----
21.....	21-16	12	0-23	.5	21- 3	9	4-16	10	17-20	11	-----
22.....	10-20	17	10- 8	11	10-12	21	13- 1	22	2- 9	23	-----
23.....	22- 0	22	22-17	17	22	9	23-11	10	12-21	11	-----
24.....	0-23	4	10- 3	23	10-21	22	22- 9	23	-----	-----	-----
25.....	22-13	9	22-13	5	22- 6	10	7-19	11	20-21	12	-----
26.....	10-17	14	10-22	11	10-16	22	17- 5	23	6- 9	0	-----
27.....	22- 6	19	22- 8	17	22- 2	10	3-14	11	15-21	12	-----
28.....	10- 6	1	10-18	23	10-11	22	12- 0	23	1- 9	0	-----
29.....	23-10	6	23- 3	5	23-10	11	11-22	12	-----	-----	-----
30.....	11-14	11	11-13	11	11-19	23	20- 8	0	9-10	1	-----
31.....	0-23	17	23	17	23- 5	11	6-18	12	19-22	13	-----
32.....	11- 3	22	0-23	0	11-15	23	16- 4	0	5-10	10	-----
33.....	23- 7	3	23-20	6	23- 0	11	1-13	12	14-22	13	-----
34.....	11	8	11- 5	12	11-23	0	0-10	1	-----	-----	-----
35.....	0-20	14	0-15	18	0- 9	12	10-21	13	22-23	14	-----
36.....	12- 0	19	12- 1	0	12-18	0	19- 7	1	8-11	2	-----
37.....	0- 4	0	0-10	6	0- 4	12	5-17	13	18-23	14	-----
38.....	0-23	6	12-20	12	12-14	0	15- 2	1	3-11	2	-----
39.....	0-17	11	0- 6	18	0-12	13	13-23	14	-----	-----	-----
40.....	12-21	16	12-15	0	12-22	1	23-11	2	-----	-----	-----
41.....	1	21	1	6	1- 7	13	8-20	14	21- 0	15	-----
42.....	13-10	3	0-23	13	13-17	1	18- 6	2	7-12	3	-----
43.....	1-14	8	1-22	19	1- 3	13	4-16	14	17- 0	15	-----
44.....	13-16	13	13- 8	1	13	1	14- 1	2	2-12	3	-----
45.....	0-23	19	1-17	7	1-11	14	12- 0	15	-----	-----	-----
46.....	13- 7	0	13- 3	13	13-21	2	22- 9	3	10-12	4	-----
47.....	2-11	5	2-13	1	2- 6	14	7-19	15	20- 1	16	-----
48.....	14-15	10	14-22	1	14-16	2	17- 5	3	6-13	4	-----
49.....	2- 0	16	2- 8	7	2	14	3-15	15	16- 1	16	-----
50.....	14- 4	21	14-18	3	14- 0	3	1-13	4	-----	-----	-----
51.....	2- 8	2	2- 3	19	2-10	15	11-23	16	0- 1	17	-----
52.....	0-23	8	0-23	2	14-20	3	21- 8	4	9-13	5	-----
(53).....	4-16	12	4-23	7	4	13	5-16	14	17- 3	15	-----

Table 34.—*For summation of long-period constituents*

ASSIGNMENT OF DAILY SUMS FOR CONSTITUENT MF

Constituent division	Days of series																	
0	1	28	55	82*	110	137	164*	192	219	246	274	301	328	356				
1	2	29	56	84	111	138	166	193	220	248*	275	302	330*	357				
2	3	30	57*	85	112	139	167	194	221	249	276	303	331	358				
3	4	31	59	86	113	141*	168	195	223*	250	277	304*	332	359				
4	5	32	60	87	114	142	169	196	224	251	278	306	333	360				
5	6	34*	61	88	115*	143	170	197*	225	252	279	307	334	361				
6	7	35	62	89	117	144	171	199	226	253	281*	308	335	363				
7	8	36	63	90*	118	145	172	200	227	254	282	309	336	364				
8	9	37	64	92	119	146	174*	201	228	256*	283	310	337*	365				
9	10	38	65	93	120	147	175	202	229	257	284	311	339	366				
10	11	39	67*	94	121	149*	176	203	230*	258	285	312*	340	367				
11	12	40	68	95	122	150	177	204	232	259	286	314	341	368				
12	13	42*	69	96	123*	151	178	205*	233	260	287	315	342	369				
13	14	43	70	97	125	152	179	207	234	261	289*	316	343					
14	15	44	71	98	126	153	180	208	235	262	290	317	344					
15	16	45	72	100*	127	154	182*	209	236	263*	291	318	345*					
16	17	46	73	101	128	155	183	210	237	265	292	319						
17	18	47	75*	102	129	156*	184	211	238*	266	293	320	348					
18	19	48	76	103	130	158	185	212	240	267	294	322*	349					
19	20	49*	77	104	131*	159	186	213	241	268	295	323	350					
20	21	51	78	105	133	160	187	215*	242	269	297*	324	351					
21	22	52	79	106	134	161	188	216	243	270	298	325	352					
22	23	53	80	108*	135	162	189*	217	244	271*	299	326	353					
23		54	81	109	136	163	191	218	245	273	300	327	355*					

ASSIGNMENT OF DAILY SUMS FOR CONSTITUENT MSFs

Constituent division		Days of series												
0		1	30	60*	89	119	148	178	207	237	266	296	325	355
1		2	31	61	90	120	149*	179	208*	238	268*	297	327*	356
2		3	33*	62	92*	121	151	180	210	239	269	298	328	357
3		4	34	63	93	122	152	181*	211	240*	270	300*	329	359*
4		5*	35	65*	94	124*	153	183	212	242	271	301	330	360
5		7	36	66	95	125	154	184	213*	243	272*	302	332*	361
6		8	37*	67	96*	126	156*	185	215	244	274	303	332	362
7		9	39	68	98	127	157	186	216	245	275	304*	334	364*
8		10	40	69*	99	128*	158	188*	217	247*	276	306	335	365
9		12*	41	71	100	130	159	189	218	248	277	307	336*	366
10		13	42	72	101*	131	160*	190	220*	249	279*	308	338	367
11		14	44*	73	103	132	162	191	221	250	280	309	339	368*
12		15	45	74	104	133*	163	192*	222	252*	281	311*	340	-----
13		17*	46	76*	105	135	164	194	223	253	282	312	341	-----
14		18	47	77	106	136	165*	193	224*	254	284*	313	343*	-----
15		19	49*	78	108*	137	167	196	226	255	285	314	344	-----
16		20	50	79	109	138	168	197*	227	256*	286	316*	345	-----
17		21*	51	80*	110	140*	169	199	228	258	287	317	346	-----
18		23	52	82	111	141	170	200	229	259	288*	318	348*	-----
19		24	53*	83	112*	142	172*	201	231*	260	290	319	349	-----
20		25	55	84	114	143	173	202	232	261	291	320*	350	-----
21		26	56	85*	115	144*	174	201*	233	263*	292	322	351	-----
22		23*	57	87	116	146	175	205	234	264	293	323	352*	-----
23		29	58	88	117*	147	176*	206	236*	265	295*	324	354	-----

Table 34.—*For summation of long-period constituents—Continued*

## ASSIGNMENT OF DAILY SUMS FOR CONSTITUENT Mm

Constituent division	Days of series																	
0.....	1	28	56	83	111	138	166	193	221	249*	276	304	331	359				
1.....	2	29	57	84	112	139*	167	195*	222	250	277	305	332	360				
2.....	3	30	58	85*	113	141	168	196	223	251	278	306	333*	361				
3.....	4	32*	59	87	114	142	169	197	224	252	280*	307	335	362				
4.....	5	33	60	88	115	143	170*	198	226*	253	281	308	336	363				
5.....	6	34	61	89	116*	144	172	199	227	254	282	309	337	364*				
6.....	7	35	63*	90	118	145	173	200	228	255	283	311*	338	366				
7.....	9*	36	64	91	119	146	174	201*	229	257*	284	312	339	367				
8.....	10	37	65	92	120	147*	175	203	230	258	285	313	340	368				
9.....	11	38	66	94*	121	149	176	204	231	259	286	314	342*	369				
10.....	12	40*	67	95	122	150	177	205	232*	260	288*	315	343	---				
11.....	13	41	68	96	123	151	178*	206	234	261	289	316	344	---				
12.....	14	42	69	97	125*	152	180	207	235	262	290	317	345	---				
13.....	15*	43	71*	98	126	153	181	208	236	263*	291	319*	346	---				
14.....	17	44	72	99	127	154	182	209*	237	265	292	320	347	---				
15.....	18	45	73	100	128	156*	183	211	238	266	293	321	348	---				
16.....	19	46*	74	102*	129	157	184	212	239	267	294*	322	350*	---				
17.....	20	48	75	103	130	158	185	213	240*	268	296	323	351	---				
18.....	21	49	76	104	131	159	187*	214	242	269	297	324	352	---				
19.....	22	50	77*	105	133*	160	188	215	243	270	298	325*	353	---				
20.....	23*	51	79	106	134	161	189	216	244	271*	299	327	354	---				
21.....	25	52	80	107	135	162	190	218*	245	273	300	328	355	---				
22.....	26	53	81	108*	136	164*	191	219	246	274	301	329	356*	---				
23.....	27	54*	82	110	137	165	192	220	247	275	302*	330	358	---				

## ASSIGNMENT OF DAILY SUMS FOR CONSTITUENT Sa

Constituent division	Days of series	Constituent division		Days of series
		0	1	
0.....	{ 1- 8 359-369	12	13	176-190
1.....	9- 23	14		191-205
2.....	24- 38	15		206-221
3.....	39- 53			222-236
4.....	54- 69	16		237-251
5.....	70- 84	17		252-266
6.....	85- 99	18		267-282
7.....	100-114	19		283-297
		20		298-312
8.....	115-129	21		313-327
9.....	130-145	22		328-342
10.....	146-160	23		343-358
11.....	161-175			

Table 35.—*Products* ( $a \frac{S}{15}$ ) for Form 444

Constituent	Time meridian in hours = S ÷ 15							
	1.000	2.000	3.000	4.000	5.000	5.500	6.000	6.500
	Products, in degrees							
M <sub>2</sub> .....	28.98	57.97	86.95	115.94	144.92	159.41	173.90	188.40
S <sub>2</sub> .....	30.00	60.00	90.00	120.00	150.00	165.00	180.00	195.00
N <sub>2</sub> .....	28.44	56.88	85.32	113.76	142.20	156.42	170.64	184.86
K <sub>1</sub> .....	15.04	30.08	45.12	60.16	75.21	82.73	90.25	97.77
M <sub>4</sub> .....	57.97	115.94	173.90	231.87	289.84	318.83	347.81	16.79
O <sub>1</sub> .....	13.94	27.89	41.83	55.77	69.72	76.69	83.66	90.63
M <sub>5</sub> .....	86.95	173.90	260.86	347.81	74.76	118.24	161.71	205.19
(MK) <sub>3</sub> .....	44.03	88.05	132.08	176.10	220.13	242.14	264.15	286.16
S <sub>4</sub> .....	60.00	120.00	180.00	240.00	300.00	330.00	0.00	30.00
(MN) <sub>4</sub> .....	57.42	114.85	172.27	229.70	287.12	315.83	344.54	13.25
v <sub>2</sub> .....	28.51	57.03	85.54	114.05	142.56	156.82	171.08	185.33
S <sub>6</sub> .....	90.00	180.00	270.00	0.00	90.00	135.00	180.00	225.00
$\mu_2$ .....	27.97	55.94	83.90	111.87	139.84	153.83	167.81	181.79
(2N) <sub>2</sub> .....	27.90	55.79	83.69	111.58	139.48	153.42	167.37	181.32
(OO) <sub>1</sub> .....	16.14	32.28	48.42	64.56	80.70	88.77	96.83	104.90
$\lambda_2$ .....	29.46	58.91	88.37	117.82	147.28	162.01	176.73	191.46
S <sub>1</sub> .....	15.00	30.00	45.00	60.00	75.00	82.50	90.00	97.50
M <sub>1</sub> .....	14.50	28.99	43.49	57.99	72.48	79.73	86.98	94.23
J <sub>1</sub> .....	15.59	31.17	46.76	62.34	77.93	85.72	93.51	101.31
M <sub>M</sub> .....	0.54	1.09	1.63	2.18	2.72	2.99	3.27	3.54
S <sub>sa</sub> .....	0.08	0.16	0.25	0.33	0.41	0.45	0.49	0.53
S <sub>a</sub> .....	0.04	0.08	0.12	0.16	0.21	0.23	0.25	0.27
MS <sub>1</sub> .....	1.02	2.03	3.05	4.06	5.08	5.59	6.10	6.60
M <sub>f</sub> .....	1.10	2.20	3.29	4.39	5.49	6.04	6.59	7.14
$\rho_1$ .....	13.47	26.94	40.41	53.89	67.36	74.09	80.83	87.56
Q <sub>1</sub> .....	13.40	26.80	40.20	53.59	66.99	73.69	80.39	87.09
T <sub>2</sub> .....	29.96	59.92	89.88	119.84	149.79	164.77	179.75	194.73
R <sub>2</sub> .....	30.04	60.08	90.12	120.16	150.21	165.23	180.25	195.27
(2Q) <sub>1</sub> .....	12.85	25.71	38.56	51.42	64.27	70.70	77.13	83.55
P <sub>1</sub> .....	14.96	29.92	44.88	59.84	74.80	82.27	89.75	97.23
(2SM) <sub>2</sub> .....	31.02	62.03	93.05	124.06	155.08	170.59	186.10	201.60
M <sub>3</sub> .....	43.48	86.95	130.43	173.90	217.38	239.12	260.86	282.60
L <sub>2</sub> .....	29.53	59.06	88.59	118.11	147.64	162.41	177.17	191.94
(2MK) <sub>3</sub> .....	42.93	85.85	128.78	171.71	214.64	236.10	257.56	279.03
K <sub>2</sub> .....	30.08	60.16	90.25	120.33	150.41	165.45	180.49	195.53
M <sub>8</sub> .....	115.94	231.87	347.81	103.75	219.68	277.65	335.62	33.59
(MS) <sub>4</sub> .....	58.98	117.97	176.95	235.94	294.92	324.41	353.90	23.40

Table 35.—*Products* ( $a \frac{S}{15}$ ) for Form 444—Continued

Constituent	Time meridian in hours = $S \div 15$							
	7.000	8.000	9.000	10.000	10.500	11.000	11.500	12.000
	Products in degrees							
M <sub>2</sub> .....	202.89	231.87	260.86	289.84	304.33	318.83	333.32	347.81
S <sub>2</sub> .....	210.00	240.00	270.00	300.00	315.00	330.00	345.00	0.00
N <sub>2</sub> .....	199.08	227.52	255.96	284.40	298.62	312.84	327.06	341.28
K <sub>1</sub> .....	105.29	120.33	135.37	150.41	157.93	165.45	172.97	180.49
M <sub>4</sub> .....	45.78	103.75	161.71	219.68	248.67	277.65	306.63	335.62
O <sub>1</sub> .....	97.60	111.54	125.49	139.43	146.40	153.37	160.34	167.32
M <sub>6</sub> .....	248.67	335.62	62.57	149.52	193.00	236.48	279.95	323.43
(MK) <sub>3</sub> .....	308.18	352.20	36.23	80.25	102.26	124.28	146.29	168.30
S <sub>4</sub> .....	60.00	120.00	180.00	240.00	270.00	300.00	330.00	0.00
(MN) <sub>4</sub> .....	41.97	99.39	156.81	214.24	242.95	271.66	300.37	329.09
v <sub>2</sub> .....	199.59	228.10	256.61	285.18	299.38	313.64	327.89	342.15
S <sub>6</sub> .....	270.00	0.00	90.00	180.00	225.00	270.00	315.00	0.00
$\mu_2$ .....	195.78	223.75	251.71	279.68	293.67	307.65	321.63	335.62
(2N) <sub>2</sub> .....	195.27	223.16	251.06	278.95	292.90	306.85	320.80	334.74
(OO) <sub>1</sub> .....	112.97	129.11	145.25	161.39	169.46	177.53	185.60	193.67
$\lambda_2$ .....	206.19	235.65	265.10	294.56	309.28	324.01	338.74	353.47
S <sub>1</sub> .....	105.00	120.00	135.00	150.00	157.50	165.00	172.50	180.00
M <sub>1</sub> .....	101.48	115.97	130.47	144.97	152.22	159.46	166.71	173.96
J <sub>1</sub> .....	109.10	124.68	140.27	155.85	163.65	171.44	179.23	187.03
M <sub>m</sub> .....	3.81	4.35	4.90	5.44	5.72	5.99	6.26	6.53
S <sub>sa</sub> .....	0.57	0.66	0.74	0.82	0.86	0.90	0.94	0.99
S <sub>a</sub> .....	0.29	0.33	0.37	0.41	0.43	0.45	0.47	0.49
MSf.....	7.11	8.13	9.14	10.16	10.67	11.17	11.68	12.19
Mf.....	7.69	8.78	9.88	10.98	11.53	12.08	12.63	13.18
$\rho_1$ .....	94.30	107.77	121.24	134.72	141.45	148.19	154.92	161.66
Q <sub>1</sub> .....	93.79	107.19	120.59	133.99	140.69	147.39	154.08	160.78
T <sub>2</sub> .....	209.71	239.67	269.63	299.59	314.57	329.55	344.53	359.51
R <sub>2</sub> .....	210.29	240.33	270.37	300.41	315.43	330.45	345.47	0.49
(2Q) <sub>1</sub> .....	89.98	102.83	115.69	128.54	134.97	141.40	147.82	154.25
P <sub>1</sub> .....	104.71	119.67	134.63	149.59	157.07	164.55	172.03	179.51
(2SM) <sub>2</sub> .....	217.11	248.13	279.14	310.16	325.67	341.17	356.68	372.19
M <sub>3</sub> .....	304.33	347.81	31.29	74.76	96.50	118.24	139.98	161.71
L <sub>2</sub> .....	206.70	236.23	265.76	295.28	310.05	324.81	339.58	354.34
(2MK) <sub>3</sub> .....	300.49	343.42	26.34	69.27	90.73	112.20	133.66	155.13
K <sub>2</sub> .....	210.57	240.66	270.74	300.82	315.86	330.90	345.94	0.99
M <sub>8</sub> .....	91.55	207.49	323.43	79.36	137.33	195.30	253.27	311.24
(MS) <sub>4</sub> .....	52.89	111.87	170.86	229.84	259.33	288.83	318.32	347.81

Table 36.—Angle differences for Form 445

Constituent	Jan. 1, 0 <sup>h</sup> , to Feb. 1, 0 <sup>h</sup>		Feb. 1, 0 <sup>h</sup> , to Dec. 31, 24 <sup>h</sup>				Jan. 1, 0 <sup>h</sup> , to Dec. 31, 24 <sup>h</sup>			
			Common year		Leap year		Common year		Leap year	
	°	°	°	°	°	°	°	°	°	°
M <sub>2</sub> .....	+324.2	-35.8	+136.6	-223.4	+112.2	-247.8	+100.8	-259.2	+76.4	-283.6
S <sub>2</sub> .....	0	0	0	0	0	0	0	0	0	0
N <sub>2</sub> .....	+279	-81	+93	-267	+56	-304	+12	-348	+335	-25
K <sub>1</sub> .....	+31	-329	+329	-31	+330	-30	0	0	+1	-359
M <sub>4</sub> .....	+288	-72	+274	-86	+225	-135	+202	-158	+153	-207
O <sub>1</sub> .....	+294	-66	+167	-193	+142	-218	+101	-259	+76	-284
M <sub>6</sub> .....	+253	-107	+49	-311	+336	-24	+302	-58	+229	-131
(MK) <sub>3</sub> .....	+355	-5	+106	-254	+82	-278	+101	-259	+77	-283
S <sub>4</sub> .....	0	0	0	0	0	0	0	0	0	0
(MN) <sub>4</sub> .....	+243	-117	+230	-130	+168	-192	+113	-247	+51	-309
v <sub>2</sub> .....	+333	-27	+317	-43	+281	-79	+290	-70	+254	-106
S <sub>6</sub> .....	0	0	0	0	0	0	0	0	0	0
$\mu_2$ .....	+288	-72	+274	-86	+225	-135	+202	-158	+153	-207
(2N) <sub>2</sub> .....	+234	-126	+49	-311	+359	-1	+283	-77	+233	-127
(OO) <sub>1</sub> .....	+127	-233	+131	-229	+159	-201	+258	-102	+286	-74
$\lambda_2$ .....	+315	-45	+316	-44	+303	-57	+271	-89	+258	-102
S <sub>1</sub> .....	0	0	0	0	0	0	0	0	0	0
M <sub>1</sub> .....	+342	-18	+248	-112	+236	-124	+230	-130	+218	-142
J <sub>1</sub> .....	+76	-284	+12	-348	+27	-333	+88	-272	+103	-257
M <sub>m</sub> .....	+45	-315	+44	-316	+57	-303	+89	-271	+102	-258
S <sub>sa</sub> .....	+61	-299	+299	-61	+300	-60	0	0	1	-359
S <sub>a</sub> .....	+31	-329	+329	-31	+330	-30	0	0	+1	-359
MS <sub>f</sub> .....	+36	-324	+223	-137	+248	-112	+259	-101	+284	-76
M <sub>f</sub> .....	+97	-263	+162	-198	+188	-172	+259	-101	+285	-75
$\rho_1$ .....	+303	-57	+348	-12	+311	-49	+291	-69	+254	-106
Q <sub>1</sub> .....	+249	-111	+123	-237	+85	-275	+12	-348	+334	-26
T <sub>2</sub> .....	+329	-31	+31	-329	+30	-330	0	0	+259	-1
R <sub>2</sub> .....	+31	-329	+329	-31	+330	-30	0	0	+1	-359
(2Q) <sub>1</sub> .....	+204	-156	+80	-280	+28	-332	+284	-76	+232	-128
P <sub>1</sub> .....	+329	-31	+31	-329	+30	-330	0	0	+359	-1
(2SM) <sub>2</sub> .....	+36	-324	+223	-137	+248	-112	+259	-101	+284	-76
M <sub>3</sub> .....	+306	-54	+25	-335	+348	-12	+331	-29	+294	-66
L <sub>2</sub> .....	+9	-351	+180	-180	+169	-191	+189	-171	+178	-182
(2MK) <sub>3</sub> .....	+258	-102	+304	-56	+254	-106	+202	-158	+152	-208
K <sub>2</sub> .....	+61	-299	+299	-61	+300	-60	0	+1	+1	-359
M <sub>8</sub> .....	+217	-143	+186	-174	+89	-271	+43	-317	+306	-54
(MS) <sub>4</sub> .....	+324	-36	+137	-223	+112	-248	+101	-259	+76	-284

Table 37.—*U. S. Coast and Geodetic Survey tide-predicting machine No. 2*  
GENERAL GEARS AND CONNECTING SHAFTS

Gears and Shafts	Face or diameter	Number of teeth	Pitch	Period of rotation	Remarks
<i>Inches</i>					
S-1-----	0.56			Dial hours	
G-1-----	0.56	40	24	4	Hand crank shaft for operating machine.
G-2-----	0.50	120	24	4	Spur gear on shaft 1.
G-3-----	0.50	120	24	12	Spur-stud gear.
S-2-----	0.50			12	Spur gear on shaft 2.
				12	Short horizontal shaft.
G-4-----	0.41	72	24	12	Bevel gear on shaft 2.
G-5-----	0.41	72	24	12	Bevel gear on shaft 3.
S-3-----	0.50			12	Diagonal shaft connecting with middle section.
G-6-----	0.38	75	30	12	Bevel gear on shaft 3.
G-7-----	0.38	75	30	12	Bevel gear on shaft 4.
S-4-----	0.38			12	Short vertical shaft through desk top.
G-8-----	0.38	75	30	12	Bevel gear on shaft 4.
G-9-----	0.38	75	30	12	Bevel gear on shaft 5.
S-5-----	0.38			12	Short horizontal shaft.
G-10-----	0.27	75	30	12	Bevel gear on shaft 5.
G-11-----	0.27	75	30	12	Bevel gear on shaft 6.
S-6-----	0.38			12	Main vertical shaft of dial case.
G-12-----	0.17	60	48	12	Releaseable bevel gear on shaft 6.
G-13-----	0.17	120	48	24	Bevel gear on shaft 7.
S-7-----	0.15			24	Intermediate shaft to hour hand.
G-14-----	0.17	84	48	24	Bevel gear on shaft 7.
G-15-----	0.17	84	48	24	Bevel gear on shaft 8.
S-8-----	0.15			24	Hour-hand shaft.
G-16-----	0.17	180	48	12	Releaseable bevel gear on shaft 6.
G-17-----	0.17	60	48	4	Bevel gear on shaft 9.
S-9-----	0.15			4	Intermediate shaft to minute hand.
G-18-----	0.17	240	48	4	Bevel gear on shaft 9.
G-19-----	0.17	60	48	1	Bevel gear on shaft 10.
S-10-----	0.15			1	Minute-hand shaft.
G-20-----	0.17	60	48	12	Releaseable bevel gear on shaft 6.
G-21-----	0.17	120	48	24	Bevel gear on shaft 11.
S-11-----	0.15			24	Intermediate shaft to day dial.
G-22-----		1		24	Worm screw, 0.56 inch diameter, 18 threads to inch on shaft 11.
G-23-----		366		24×366	Worm wheel, 6.47 inch diameter, on shaft 12.
S-12-----	0.31			24×366	Day dial shaft.
G-24-----	0.25	46	40	12	Spur gear at top of shaft 6.
G-25-----	0.25	60	40	-----	Spur-stud gear.
G-26-----	0.25	60	40	-----	Spur-stud gear connected with gear 25 by ratchet wheel and pawl.
G-27-----	0.25	46	40	12	Spur gear at lower end of feeding roller.
G-28-----	0.41	72	24	12	Bevel gear on shaft 3.
G-29-----	0.41	72	24	12	Bevel gear on shaft 13.
S-13-----	0.44			12	Main vertical shaft of middle section.
G-30-----	0.38	110	30	12	Spur gear on shaft 13.
G-31-----	0.38	110	30	12	Spur stud gear.
G-32-----	0.38	110	30	12	Spur stud gear on shaft 14.
S-14-----	0.38			12	Front vertical shaft of rear section.
G-33-----	0.28	75	30	12	Bevel gear on shaft 14.
G-34-----	0.28	75	30	12	Bevel gear on shaft 15.
S-15-----	0.50			12	Main connecting horizontal shaft of rear section.
G-35-----	0.28	75	30	12	Bevel gear on shaft 15.
G-36-----	0.28	75	30	12	Bevel gear on shaft 16.
S-16-----	0.38			12	Rear vertical shaft of rear section.

Table 38.—*U. S. Coast and Geodetic Survey tide-predicting machine No. 2*

## CONSTITUENT GEARS AND MAXIMUM AMPLITUDE SETTINGS

Constituents	Theoretical speed per hour	Teeth in gear wheels				Gear speed per dial hour	Error per year	Maximum amplitude settings of cranks
		Vertical shafts	Intermediate shafts		Crank shafts			
			I	II	III	IV		
	°					°	°	Units
J <sub>1</sub> .....	15.5854433	107	90	52	119	15.5854342	-0.08	1.4
K <sub>1</sub> .....	15.0410686	61	73	51	85	15.0410959	+.24	11.0
K <sub>2</sub> .....	30.0821372	122	80	96	146	30.0821918	+.48	3.9
L <sub>2</sub> .....	29.5284788	104	61	56	97	29.5284773	-.01	2.4
*M <sub>1</sub> .....	14.4920521	103	85	59	148	14.4920509	-.01	1.0
M <sub>2</sub> .....	28.9841042	103	74	59	85	28.9841017	-.02	20.0
M <sub>3</sub> .....	43.4761563	86	62	70	67	43.4761675	+.10	1.4
M <sub>4</sub> .....	57.9682084	118	74	103	85	57.9682035	-.04	4.0
M <sub>6</sub> .....	86.9523126	140	62	86	67	86.9523351	+.20	1.0
M <sub>8</sub> .....	115.936168	118	37	103	85	115.9364070	+.09	0.4
N <sub>2</sub> .....	28.4397296	65	46	53	79	28.4397358	+.05	6.0
2N.....	27.8953548	68	58	46	58	27.8953627	+.07	1.0
O <sub>1</sub> .....	13.9430356	92	89	58	129	13.9430363	+.01	9.0
OO.....	16.1391016	134	131	71	135	16.1391009	-.01	0.8
P <sub>1</sub> .....	14.9589314	91	73	50	125	14.9589041	-.24	4.8
Q <sub>1</sub> .....	13.3986609	84	88	51	109	13.3986656	+.04	3.0
2Q.....	12.8542862	127	114	50	130	12.8542510	-.31	0.6
R <sub>2</sub> .....	30.0410667	85	50	43	73	30.0410959	+.24	0.4
S <sub>1</sub> .....	15.0000000	63	75	50	84	15.0000000	.00	2.0
S <sub>2</sub> .....	30.0000000	70	70	70	70	30.0000000	.00	9.8
S <sub>4</sub> .....	60.0000000	75	45	60	50	60.0000000	.00	1.0
S <sub>5</sub> .....	90.0000000	90	48	80	50	90.0000000	.00	0.4
T <sub>2</sub> .....	29.958933	81	50	45	73	29.9589041	-.24	1.0
λ <sub>2</sub> .....	29.4556254	131	65	57	117	29.4556213	-.04	0.4
μ <sub>2</sub> .....	27.9682084	125	82	74	121	27.9681516	-.50	1.2
v <sub>2</sub> .....	28.5125830	89	69	70	95	28.5125858	+.02	2.0
ρ <sub>1</sub> .....	13.4715144	69	70	41	90	13.4714286	-.75	0.8
M <sub>K</sub> .....	44.0251728	120	81	105	106	44.0251572	-.14	1.9
2MK.....	42.9271398	81	52	79	86	42.9271020	-.33	1.4
MN.....	57.4238338	135	42	53	89	57.4237560	-.68	0.7
MS.....	58.9841042	118	61	62	61	58.9841440	-.35	2.0
2SM.....	31.0158958	69	47	50	71	31.0158825	-.12	1.4
Mf.....	1.0980330	84	45	1	51	1.0980392	+.05	4.0
MSf.....	1.0158958	149	80	1	55	1.0159091	+.12	2.0
Mm.....	0.5443747	93	41	1	125	0.5443902	+.14	3.0
Sa.....	0.0410686	51	{ 149	1	-----{ 120 } 60	0.0410738	+.05	8.0
Ssa.....	0.0821372	51	149	1	125	0.0821477	+.09	3.0

\*Designed for one-half of speed of M<sub>2</sub>.

Table 39.—*Synodic periods of constituents*

## DIURNAL CONSTITUENTS

	J <sub>1</sub>	K <sub>1</sub>	M <sub>1</sub>	O <sub>1</sub>	OO	P <sub>1</sub>	Q <sub>1</sub>	2Q	S <sub>1</sub>
	Days.	Days.	Days.	Days.	Days.	Days.	Days.	Days.	Days.
K <sub>1</sub>	27.555								
M <sub>1</sub>	13.777	27.555							
O <sub>1</sub>	9.133	13.661	27.555						
OO	27.093	13.661	9.133	6.830					
P <sub>1</sub>	23.942	182.621	32.451	14.765	12.710				
Q <sub>1</sub>	6.859	9.133	13.661	27.555	5.474	9.614			
2Q	5.492	6.859	9.133	13.777	4.566	7.127	27.555		
S <sub>1</sub>	25.622	365.243	29.803	14.192	13.168	365.243	9.367	6.991	
P <sub>1</sub>	7.096	9.557	14.632	31.812	5.623	10.085	205.892	24.302	9.814

## SEMI DIURNAL CONSTITUENTS

	K <sub>2</sub>	L <sub>2</sub>	M <sub>2</sub>	N <sub>2</sub>	2N	R <sub>2</sub>	S <sub>2</sub>	T <sub>2</sub>	λ <sub>2</sub>	μ <sub>2</sub>	ν <sub>2</sub>
	Days.	Days.	Days.	Days.	Days.	Days.	Days.	Days.	Days.	Days.	Days.
L <sub>2</sub>	27.093										
M <sub>2</sub>	13.661	27.555									
N <sub>2</sub>	9.133	13.777	27.555								
2N	6.859	9.185	13.777	27.555							
R <sub>2</sub>	365.225	29.263	14.192	9.367	6.991						
S <sub>2</sub>	182.621	31.812	14.765	9.614	7.127	365.259					
T <sub>2</sub>	121.748	34.847	15.387	9.874	7.269	182.630	365.259				
λ <sub>2</sub>	23.942	205.892	31.812	14.765	9.614	25.622	27.555	29.803			
μ <sub>2</sub>	7.096	9.614	14.765	31.812	205.892	7.236	7.383	7.535	10.085		
ν <sub>2</sub>	9.557	14.765	31.812	205.892	24.302	9.814	10.085	10.371	15.906	27.555	
2SM	16.064	10.085	7.383	5.823	4.807	15.387	14.765	14.192	9.614	4.922	5.992

Table 40.—*Day of the common year corresponding to day of month*

[For leap year increase all numbers after February 29 by 1 day]

Day of month.	Jan.	Feb.	Mar.	Apr.	May	June	July	Aug.	Sept.	Oct.	Nov.	Dec.
1	1	32	60	91	121	152	182	213	244	274	305	335
2	2	33	61	92	122	153	183	214	245	275	306	336
3	3	34	62	93	123	154	184	215	246	276	307	337
4	4	35	63	94	124	155	185	216	247	277	308	338
5	5	36	64	95	125	156	186	217	248	278	309	339
6	6	37	65	96	126	157	187	218	249	279	310	340
7	7	38	66	97	127	158	188	219	250	280	311	341
8	8	39	67	98	128	159	189	220	251	281	312	342
9	9	40	68	99	129	160	190	221	252	282	313	343
10	10	41	69	100	130	161	191	222	253	283	314	344
11	11	42	70	101	131	162	192	223	254	284	315	345
12	12	43	71	102	132	163	193	224	255	285	316	346
13	13	44	72	103	133	164	194	225	256	286	317	347
14	14	45	73	104	134	165	195	226	257	287	318	348
15	15,	46	74	105	135	166	196	227	258	288	319	349
16	16	47	75	106	136	167	197	228	259	289	320	350
17	17	48	76	107	137	168	198	229	260	290	321	351
18	18	49	77	108	138	169	199	230	261	291	322	352
19	19	50	78	109	139	170	200	231	262	292	323	353
20	20	51	79	110	140	171	201	232	263	293	324	354
21	21	52	80	111	141	172	202	233	264	294	325	355
22	22	53	81	112	142	173	203	234	265	295	326	356
23	23	54	82	113	143	174	204	235	266	296	327	357
24	24	55	83	114	144	175	205	236	267	297	328	358
25	25	56	84	115	145	176	206	237	268	298	329	359
26	26	57	85	116	146	177	207	238	269	299	330	360
27	27	58	86	117	147	178	208	239	270	300	331	361
28	28	59	87	118	148	179	209	240	271	301	332	362
29	29		88	119	149	180	210	241	272	302	333	363
30	30		89	120	150	181	211	242	273	303	334	364
31	31		90		151		212	243		304		365

Table 41.—Values of  $h$  in formula  $h = (1 + r^2 + 2r \cos x)^{\frac{1}{2}}$

x	r										x
	0.0	0.1	0.2	0.3	0.4	0.5	0.6	0.7	0.8	0.9	
0	1.000	1.100	1.200	1.300	1.400	1.500	1.600	1.700	1.800	1.900	2.000
10	1.000	1.099	1.197	1.296	1.396	1.495	1.594	1.694	1.793	1.893	1.992
20	1.000	1.095	1.190	1.286	1.383	1.480	1.577	1.675	1.773	1.871	1.970
30	1.000	1.088	1.177	1.269	1.361	1.455	1.549	1.644	1.739	1.835	1.932
40	1.000	1.079	1.160	1.245	1.331	1.420	1.510	1.601	1.693	1.786	1.879
50	1.000	1.067	1.139	1.215	1.294	1.376	1.460	1.546	1.634	1.723	1.813
60	1.000	1.054	1.114	1.179	1.249	1.323	1.400	1.480	1.562	1.646	1.732
70	1.000	1.038	1.085	1.138	1.197	1.262	1.331	1.403	1.479	1.557	1.638
80	1.000	1.022	1.053	1.093	1.140	1.193	1.252	1.316	1.385	1.457	1.532
90	1.000	1.005	1.020	1.044	1.077	1.118	1.166	1.221	1.281	1.345	1.414
100	1.000	0.988	0.985	0.993	1.010	1.037	1.073	1.117	1.167	1.224	1.286
110	1.000	0.970	0.950	0.941	0.941	0.953	0.974	1.006	1.045	1.093	1.147
120	1.000	0.954	0.917	0.889	0.872	0.866	0.872	0.889	0.917	0.954	1.000
130	1.000	0.939	0.885	0.839	0.804	0.779	0.767	0.768	0.782	0.808	0.845
140	1.000	0.926	0.856	0.794	0.740	0.696	0.664	0.646	0.644	0.657	0.684
150	1.000	0.915	0.833	0.755	0.684	0.620	0.566	0.527	0.504	0.501	0.518
160	1.000	0.907	0.815	0.725	0.639	0.557	0.482	0.418	0.369	0.344	0.347
170	1.000	0.902	0.804	0.706	0.610	0.515	0.422	0.334	0.254	0.193	0.174
180	1.000	0.900	0.800	0.700	0.600	0.500	0.400	0.300	0.200	0.100	0.000

Table 42.—Values of  $k$  in formula  $k = \tan^{-1} \frac{r \sin x}{1 + r \cos x}$

[When  $\tau$  is between  $180^\circ$  and  $360^\circ$ , tabular values are negative]

## EXPLANATION OF SYMBOLS USED IN THIS BOOK

Although the following list is fairly comprehensive, some of the symbols given may at times be used in the text to represent other quantities not listed below, but such application will be made clear by the context.

- A**    (1) General symbol for a tidal constituent or its amplitude. It is sometimes written with a subscript to indicate the species of the constituent (par. 52).  
 (2) General symbol with an identifying subscript for a constituent term in the development of the lunar tide-producing force (par. 66).  
 (3) The particular tidal constituent being cleared by the elimination process (par. 245).  
 (4) Azimuth of tide-producing body reckoned from the south through the west (par. 80).  
 (5) Azimuth of horizontal component of force in any given direction (par. 85).
- a**    (1) Speed or rate of change in argument of constituent *A*.  
 (2) Mean radius of earth.
- B**    (1) Tidal constituent following constituent *A* in a series.  
 (2) General symbol with an identifying subscript for a constituent term in the development of the solar tide-producing force (par. 117).  
 (3) General symbol for disturbing constituents in elimination process (par. 245).
- b**    Speed or rate of change in argument of constituent *B*.
- C**    (1) Mean constituent coefficient (par. 74).  
 (2) General symbol for coefficients of cosine terms in Fourier series (par. 187).
- c**    Reciprocal of mean value of  $1/d$ .
- $c_1$**     Reciprocal of mean value of  $1/d_1$ .
- D**    Declination of moon or sun.
- d**    Distance from center of earth to center of moon.
- $d_1$**     Distance from center of earth to center of sun.
- E**    (1) Mass of earth.  
 (2) Argument of tidal constituent (same as  $V+u$ ).
- e**    Eccentricity of moon's orbit.
- $e_1$**     Eccentricity of earth's orbit.
- F**    Reduction factor, reciprocal of node factor *f* (par. 78).
- $F_a$**     Horizontal component of tide-producing force in azimuth *A*. When numerals are annexed, the first digit (3 or 4) signifies the power of the parallax of the moon or sun involved in the development and the second digit (0, 1, 2, or 3) indicates the species of the terms included in the group. Thus  $F_{a30}$  represents that part of the horizontal component in azimuth *A* that comprises the long-period terms depending upon the cube of the parallax.
- $F_s$**     South horizontal component of tide-producing force. (See  $F_a$  for explanation of annexed numerals.)
- $F_v$**     Vertical component of tide-producing force. (See  $F_a$  for explanation of annexed numerals.)
- $F_w$**     West horizontal component of tide-producing force. (See  $F_a$  for explanation of annexed numerals.)
- f**    Node factor (par. 77).
- G**    (1) Greenwich epoch or phase lag of a tidal constituent (par. 226).  
 (2) Gear ratio of predicting machine (par. 396).
- g**    (1) Mean acceleration of gravity on earth's surface.  
 (2) Modified epoch of tidal constituent, same as  $\kappa'$  (par. 225).
- H**    Mean amplitude of a tidal constituent (par. 143).

- $H_0$  Mean water level above datum used for tabulation.
- $h$  (1) Mean longitude of sun; also rate of change in same.  
 (2) Height of tide at any time.
- $h_3$  Height of equilibrium tide involving cube of moon's parallax. A second digit in the subscript limits the height to that due to terms of the single species indicated by this digit (pars. 97 and 101).
- $h_4$  Height of equilibrium tide involving 4th power of moon's parallax. A second digit in the subscript has the same significance as in the case of  $h_3$ .
- $I$  Obliquity of lunar orbit with respect to earth's equator.
- $i$  Inclination of lunar orbit to the ecliptic.
- $J_1$  Tidal constituent.
- $j$  Longitude of moon in its orbit reckoned from lunar intersection.
- $K_1, K_2$  Tidal constituents.
- $KJ_2, KP_1, KQ_1$  Tidal constituents.
- $k$  Difference between mean and true longitude of moon (par. 59).
- $L$  Longitude of place; positive for west longitude, negative for east longitude.
- $L_2, LP_1$  Tidal constituents.
- $M$  Mass of moon.
- $M_1, M_2, M_3, M_4, M_6, M_8$  Tidal constituents.
- $Mf$  Tidal constituent.
- $MK_3, 2MK_3, MK_1$  Tidal constituents.
- $Mm$  Tidal constituent.
- $MN_4, 2MN_6, MNS_2$  Tidal constituents.
- $MP_1$  Tidal constituent.
- $2MS_2, MS_4, 2MS_6, 3MS_8, 2(MS)_8$  Tidal constituents.
- $MSf$  Tidal constituent.
- $MSN_6, 2MSN_8$  Tidal constituents.
- $m$  Ratio of mean motion of sun to that of moon (par. 62).
- $N$  Longitude of moon's node; also rate of change in same.
- $N_2, 2N_2, NJ_1$  Tidal constituents.
- $O_1, OO_1$  Tidal constituents.
- $P$  Mean longitude of lunar perigee reckoned from lunar intersection (par. 122).
- $P_1$  Tidal constituent.
- $p$  (1) Mean longitude of lunar perigee; also rate of change in same.  
 (2) Numeral indicating species of constituent, frequently written as the subscript of the constituent symbol. In special case used with long-period constituents to show number of periods in month or year.
- $p_1$  Mean longitude of solar perigee; also rate of change in same.
- $Q$  Term in argument of constituent  $M_1$  (par. 123).
- $Q_a$  Factor in amplitude of constituent  $M_1$  (par. 122).
- $Q_u$  Term in argument of constituent  $M_1$  (par. 122).
- $Q_1, 2Q_1$  Tidal constituents.
- $R$  (1) Amplitude of constituent pertaining to a particular time (par. 143).  
 (2) Term in argument of constituent  $L_2$  (par. 129).
- $R_a$  Factor in amplitude of constituent  $L_2$  (par. 129).
- $R_2, RP_1$  Tidal constituents.
- $r$  Distance of any point from center of earth.
- $S$  (1) Mass of sun.  
 (2) Longitude of time meridian; positive for west longitude, negative for east longitude.  
 (3) General symbol for coefficients of sine terms in Fourier series (par. 187).  
 (4) Working scale factor of predicting machine.
- $S'$  Solar factor  $U_1/U$  (par. 118).
- $S_1, S_2, S_3, S_4, S_6, S_8$  Tidal constituents.
- $Sa$  Tidal constituent.
- $SK_3$  Tidal constituent.
- $2SM_6$  Tidal constituent.

$\text{SO}_1, \text{SO}_3$	Tidal constituents.
$\text{Ssa}$	Tidal constituent.
$s$	Mean longitude of moon: also rate of change in same.
$s'$	True longitude of moon in orbit referred to equinox (par. 59).
$T$	(1) Number of Julian centuries reckoned from Greenwich mean noon, December 31, 1899. (2) Hour angle of mean sun. (3) Time expressed in degrees of constituent reckoned from phase zero of Greenwich argument (par. 439).
$T_2$	Tidal constituent.
$t$	(1) Hour angle of tide-producing body. (2) Time reckoned from beginning of tidal series.
$U$	Basic factor ( $M/E$ ) ( $a/c$ ) <sup>3</sup> .
$U_1$	Factor ( $S/E$ ) ( $a/c_1$ ) <sup>3</sup> .
$u$	Part of constituent argument depending upon variations in obliquity of lunar orbit (par. 71).
$V$	(1) Principal portion of constituent argument (par. 71). (2) Velocity of current (par. 330).
$(V+u)$	Constituent argument at any time.
$(V_o+u)$	Constituent argument at beginning of a tidal series.
$V_g$	Potential due to gravity at earth's surface (par. 96).
$V_3$	Tide-producing potential involving cube of moon's parallax (par. 94).
$V_4$	Tide-producing potential involving 4th power of moon's parallax (par. 94).
$X$	Longitude of observer reckoned in celestial equator from lunar intersection.
$Y$	Latitude of observer. When combined with a subscript consisting of a letter and numerals, it represents the latitude factor to be used with the tidal force component similarly marked (par. 79).
$z$	Geocentric zenith distance of tide-producing body.
$\alpha$ ( <i>Alpha</i> )	General symbol for the initial phase of tidal constituent <i>A</i> .
$\beta$ ( <i>Beta</i> )	Initial phase of constituent <i>B</i> .
$\gamma$ ( <i>Gamma</i> )	Initial phase of constituent <i>C</i> .
$\delta$ ( <i>Delta</i> )	Initial phase of constituent <i>D</i> .
$\epsilon$ ( <i>Epsilon</i> )	Initial phase of constituent <i>E</i> .
$\xi$ ( <i>Zeta</i> )	The explement of the initial phase of a constituent (par. 221).
$\theta_1$ ( <i>Theta</i> )	Tidal constituent, same as $\lambda\theta_1$ .
$\kappa$ ( <i>Kappa</i> )	Local phase lag or epoch of tidal constituent (par. 144).
$\kappa'$	Modified epoch of tidal constituent (par. 225).
$\lambda_2$ ( <i>Lambda</i> )	Tidal constituent.
$\mu$ ( <i>Mu</i> )	Attraction of gravitation between unit masses at unit distance.
$\mu_2$	Tidal constituent, same as $2\text{MS}_2$ .
$\nu$ ( <i>Nu</i> )	Right ascension of lunar intersection (par. 24).
$\nu'$	Term in argument of lunisolar constituent $K_1$ (par. 133).
$2\nu''$	Term in argument of lunisolar constituent $K_2$ (par. 135).
$\nu_2$	Tidal constituent.
$\xi$ ( <i>Xi</i> )	Longitude in moon's orbit of lunar intersection (par. 24).
$\pi$ ( <i>Pi</i> )	An angle of 3.14159 radians or $180^\circ$ .
$\pi_1$	Tidal constituent, same as $\text{TK}_1$ .
$\rho_1$ ( <i>Rho</i> )	Tidal constituent, same as $\nu K_1$ .
$\sigma_1$ ( <i>Sigma</i> )	Tidal constituent, same as $\nu J_1$ .
$\tau$ ( <i>Tau</i> )	Length of series in mean solar hours (par. 248).
$\varphi_1$ ( <i>Phi</i> )	Tidal constituent, same as $\text{KP}_1$ .
$\chi_1$ ( <i>Chi</i> )	Tidal constituent, same as $\text{LP}_1$ .
$\psi_1$ ( <i>Psi</i> )	Tidal constituent, same as $\text{RP}_1$ .
$\omega$ ( <i>Omega</i> )	Obliquity of ecliptic.
$\Upsilon$	Vernal equinox.
$\Omega$	Moon's ascending node.



## INDEX

<b>A</b>	Page	Page	
Adams, J. C.	1	Equilibrium argument	22,
Airy, George B.	1	50, 75, 108, 124, 157, 204	
Amplitude of constituent	2, 49	Equilibrium theory	28
Analysis of high and low waters	100	Equilibrium tide	28, 38
Analysis of monthly sea level	98, 114	Equinox	6
Analysis of observations	49	Eudoxas	1
Analysis of tidal currents	118	Evection	4
Anomalistic month, year	4	Explanation of tables	153
Approximation, degree of	8	Explanation of tidal movement	2
Argument. ( <i>See Equilibrium argument.</i> )		Extreme equilibrium tide	33
Astres fictifs	23	Extreme tide-producing force	13
Astronomical data	3, 153, 162	 <b>F</b>	
Astronomical day	3	Factor F. ( <i>See Reduction factor.</i> )	
Astronomical periods	163	Factor f. ( <i>See Node factor.</i> )	
Astronomical tide	30	Ferrel, William	1, 127
Augmenting factors	71, 91, 157, 228	Forms for analysis of tides	104
 <b>B</b>		Forms for predicting machine	143
Basic factor	24	Fourier series	62
 <b>C</b>		Fourth power of moon's parallax	34
Calendars	4	Fundamental astronomical data	153,
Civil day	3	162	
Coefficients	24	Fundamental formulas	10
Component. ( <i>See Constituent tides.</i> )		 <b>G</b>	
Component of force, horizontal	26	General coefficient	24
Component of force, vertical	15	General explanation, tidal movement	2
Compound tides	47, 167	Gravitational tide	30
Constituent day	3	Greatest equilibrium tide	33
Constituent hour	4	Greatest tide-producing force	13
Constituent tides	2, 16, 87	Greenwich argument	76
Formulas	21, 35, 39	Greenwich epoch	77
Tables	153, 164, 167	Gregorian calendar	4
Currents, analysis	118	 <b>H</b>	
Currents, prediction	147	Harmonic analysis	3, 49, 112
 <b>D</b>		Harmonic constants	3, 49, 143
Darwin, G. H.	1	Harmonic prediction	3, 123
Datum for prediction	124, 144	Harris, Rollin A.	1
Day, several kinds	3	High and low water analysis	100
Day of year, table	309	Historical statement	1
Declinational factor	17	Horizontal component, tide-producing force	26, 37
Degree of approximation	8	Hour, several kinds	4
Development of tide-producing force	10	Hourly heights	104
Diurnal constituents	16	Hydraulic current	148
 <b>E</b>		Hydrographic datum	144
Eccentricity of orbit	4	 <b>I</b>	
Eclipse year	4	Inclination of moon's orbit	6,
Elimination	84, 116, 158, 236	155, 173	
Elliptic factor	24	Inference of constants	78, 114
Epoch of constituent	49, 75		
Equations of moon's motion	19		

J	Page	Page
Julian calendar-----	4	Predicting machine. ( <i>See Tide-predicting machine.</i> )
K		Prediction of tidal currents----- 147
K <sub>1</sub> and K <sub>2</sub> tides-----	44	Prediction of tides----- 123
Kelvin, Lord-----	1, 126	Principal lunar constituents----- 21
L		Principal solar constituents----- 39
Laplace-----	1	R
Latitude-----	6	Record of observations----- 50
Latitude factors-----	17, 24, 154, 168	Reduction factor----- 25, 111, 156, 186
Length of series-----	51	S
Lesser lunar constituents-----	35	Secondary stencils----- 57, 159, 299
Lesser solar constituents-----	40	Semidiurnal constituents----- 16
Lesser tide-producing force-----	34, 40	Settings for tide-predicting machine----- 145, 306
Longitude-----	6	Shallow-water constituents----- 46, 167
Longitude, lunar and solar elements-----	162, 170	Shidy, L. P.----- 53
Long-period constituents-----	16, 87, 302	Sidereal day----- 3
L <sub>2</sub> -tide-----	43, 156, 177, 192	Sidereal hour, month, year----- 4
Lunar constituents-----	21, 35	Solar day----- 3
Lunar day-----	3	Solar factor----- 40
Lunar hour-----	4	Solar hour----- 4
Lunar intersection-----	6	Solar tides----- 39
Lunar node-----	6, 8	South component, tide-producing force----- 26, 37
Lunisolar tides-----	44	Species of constituent----- 16
M		Speed of constituent----- 3, 23
Mean constituent coefficient-----	24	Stationary wave----- 2
Mean longitude-----	7	Stencil sums----- 107
Meteorological tides-----	46	Stencils----- 53, 106, 158, 268
Month, several kinds-----	4	Summarized formulas: Equilibrium tide----- 33 Lesser tide-producing force----- 36 Principal tide-producing force----- 26
Monthly sea-level analysis-----	98, 114	Summation for analysis----- 52
Moon's motion, equations-----	19	Surface of equilibrium----- 30, 32
Moon's node-----	6, 8	Symbols used in book----- 311
Moon's parallax, 4th power-----	34	Synodical month----- 4
M <sub>1</sub> -tide-----	41, 156, 179, 192	Synodic periods of constituents----- 161, 309
N		T
Node, lunar-----	6, 8	Tables----- 162
Node factor-----	25	Explanation----- 153
Compound tides-----	47	Terdiurnal constituents----- 34
Constituent K <sub>1</sub> -----	45	Thomson, Sir William----- 1, 126
Constituent K <sub>2</sub> -----	46	Tidal currents----- 118, 147
Constituent L <sub>2</sub> -----	44	Tidal movement----- 2
Constituent M <sub>1</sub> -----	43	Tide-predicting machine----- 126
Lesser tide-producing force-----	36	Adjustments----- 139
Predictions-----	124	Automatic stopping device----- 135
Table-----	199	Base----- 127
Nodal month-----	4	Constituent cranks----- 130
O		Constituent dials----- 131
Obliquity factor-----	24	Constituent pulleys----- 132
Obliquity of ecliptic-----	6	Constituent sliding frames----- 131
Obliquity of moon's orbit-----	6	Datum of heights----- 141
Observational data-----	50	Day dial----- 128
Overtides-----	47	Dial hour----- 128
P		Dimensions----- 127
Period of constituent-----	3	Doubling gears----- 132
Periods, astronomical-----	163	Forms used----- 143
Phase lag-----	49, 75	Gear speeds----- 129
Phase of constituent-----	2	Gearing----- 128, 160, 307
Poor, Charles Lane-----	1	Graph scale----- 137
Potential-----	30	

Tide-predicting machine—Con.	Page	Tide-predicting machine—Con.	Page
Height formula	126	Summation wheels	133
Height predictions	134	Tide curve	136
Height scale	134, 141	Time dials	128
Height side	128	Time formula	126, 132
High and low water marking device	139	Time prediction	135
Hour marking device	139	Time side	128
Marigram gears	137, 141	Verification of settings	142
Marigram scale	137	Tide-producing force	10
Nonreversing ratchet	136	Tide-producing potential	31
Operation of machine	142	Tropical month, year	4
Paper	136, 142		V
Pens	138, 142	Variation inequality	4
Plane of reference	141	Vernal equinox	6
Positive and negative directions	131	Vertical component, tide-producing force	15, 34
Predicting	142		W
Releasable gears	130	West component, tide-producing force	26, 37
Scale, amplitude settings	132, 140		Y
Scale, height dial	134, 141	Year, several kinds	4
Scale, marigram	137	Young, Thomas	1
Scale, table	138		
Scale, working	135		
Setting machine	140		
Stopping device	135		
Summation chains	133		







